

MANUSCRIPT BOOK 2  
OF  
SRINIVASA RAMANUJAN



(10)

Box M88  
Mss. 2



V

—

—

2	2	144	110880
3	4	160	166320
4	6	168	221760
6	12	180	277200
8	24	192	332640
9	36	200	498960
10	48	216	554400
12	60	224	665280
16	120	240	720720
18	180	256	1081080
20	240	288	1441440
24	360	320	2162160
30	720	336	2882880
32	840	360	3603600
36	1260	384	4324320
40	1680	400	6486480
48	2520	432	7207200
60	5040	448	8648640
64	7560	480	10810800
72	10080	504	14414400
80	15120	512	17297280
8	20160	576	21621600
90	25200	600	32432400
96	27720	640	36756720
100	45360	672	43243200
108	50400	720	61261200
120	55440	768	73513440
128	83160	800	110270160

864	122522400
896	147026880
960	183783600
1008	245044800
1024	294053760
1152	367567200
1200	551350800
1280	698377680
1344	735134400
1440	1102701600
1536	1396755360
1600	2095133040
1680	2205403200
1728	2327925600
1792	2793510720
1920	3491888400
2016	4655851200
2048	5587021440
2304	6983776800
2400	10475665200
2688	13967553600
2880	20951330400
3072	27935107200
3360	41902660800
3456	48886427600
3584	64250746560
3600	7339656400
3840	80310433200
4032	97772875200
4096	128501493120
4320	146659312800

$$\text{If } p^3 + q^3 + r^3 = s^3$$

$$\text{and } \begin{cases} m = (p+q) \sqrt{\frac{p-q}{r+p}} \text{ and} \\ n = (r-p) \sqrt{\frac{r+p}{s-q}} \end{cases}$$

then

$$(pa^2 + ma^2b - rb^2)^3 + (qa^2 - ma^2b + sb^2)^3 + (ra^2 - ma^2b - pb^2)^3 = (sa^2 - ma^2b + qb^2)^3$$

$$\frac{\chi^3(x)}{\chi(x^2)} = 1 + 3x \cdot \frac{\psi(x^2)}{\psi(x)}$$

$$\frac{\chi^5(x)}{\chi(x^5)} = 1 + 5x \cdot \left[ \frac{\psi(-x^5)}{\psi(x)} \right]^2$$

## CHAPTER I

magic squares can be constructed by combining two sets of letters so that the same letter may not appear in a row, a column or a corner, for example if we want to construct a square containing  $m$  rows and  $n$  columns, we should take two sets of  $n$  letters  $A, B, C, D, E$  &c and  $P, Q, R, S, T$  &c combine them as  $A+P, A+Q, A+R$  &c,  $B+P, B+Q, B+R$  &c,  $C+P, C+Q, C+R$  &c &c and arrange them in such a way that any letter, say  $K$ , may not appear in the same row, column or corner.

Now we have algebraically constructed a square and the sum of the figures in  $m$  rows and  $n$  columns will be equal if we give an value to the letters; yet the same figure may likely appear again owing to arithmetical values. This difficulty is removed from the following truths.

$A+M$  differs from  $A+N$  as  $B+M$  from  $B+N$ , as  $C+M$  from  $C+N$  &c.

Cor. 1. If  $A+P, A+Q, A+R$  &c are in A.P. then  $B+P, B+Q, B+R$  &c are also in A.P.

Cor. 2. If the values of  $A+P, A+Q, A+R, A+S$  &c are known, then from the value of  $B+P$ , those of  $B+Q, B+R, B+S$  &c are, also known.

N.B. We should not give separate values to  $A, B, C$  &c and to  $P, Q, R$  &c but we should give values to  $A+P, A+Q, \&c$ .



1. The values of  $A+P, B+P, C+P, D+P$  are 8, 10, 12, 14  
 and the value of  $C+B = 263$  find  $A+B, B+C, C+D, D+A$   
 $A+B=22; B+C=24; C+D=26; D+A=28$

2.  $A+P, A+Q, A+S$  are 5, 7, 11 respectively  
 $B+Q=28; B+R=24; C+P=21; C+S=23$   
 $D+R=10; D+T=21; E+T=23$

2. To construct a square containing three rows  
 let  $a$  be row or a column,  $m$  middle row or  
 column and  $c$  a corner &  $x$  the middle fig.

(ii) If  $a, m$  &  $c$  are different,  
 write  $\frac{1}{3}(m_1 + m_2 + c_1 + c_2 - 5)$  in the middle.  
 3 being the whole sum and supply the  
 other figures.

Sol.  $m_1 + m_2 + c_1 + c_2 = 5 + 2x$

$\therefore x = \frac{1}{2}(m_1 + m_2 + c_1 + c_2 - 5)$

As this is only relation existing between  
 $c_1, c_2, m$  we may supply the other figures  
 as we choose.

(iii) If the columns and rows are equal and  
 the corners different,

write  $\frac{c_1 + c_2 - 2}{3}$  in the middle.

Sol. By I we get  $x = \frac{1}{2}(m_1 + m_2 + c_1 + c_2 - 5)$

but here  $m_1 = m_2 = 2 \therefore 2 + 2 + c_1 + c_2 - 5$

$\therefore x = \frac{c_1 + c_2 - 1}{3}$

When the rows columns and corners are all equal write  $\frac{a}{3}$  in the middle.

Sol.  $a = \frac{c_1 + c_2}{3}$ . But  $c_1 = c_2 = a$ .  $\therefore a = \frac{a+a}{3}$

Cor. 1. The numbers in the two corners and in the middle row and column are in A.P.

Since the last is one-third of the sum the 1st & the 3rd are together twice the 2nd and consequently they are in A.P.

Ex. 1. Construct a square where  $r = m = c = 18$   
 (i)  $r = m = c = 27$  and all  $m, s$  are odd.

6	1	21
7	5	16
8	9	11

25	14	11
5	9	15
7	17	3

2. (A)  $r = m = c = 36$  and all  $m, s$  are even  
 (B)  $r = m = c = 36$  and all  $m, s$  are odd

16	9	11
13	14	9
6	18	12

24	9	3
11	14	11
18	17	11

V.B. The solution fails when the given sum is not a multiple of 3.

4  
 To construct a square for  $A+B+C+P+Q+R$ .

$C+Q$	$A+P$	$B+R$
$A+R$	$B+Q$	$C+P$
$B+P$	$C+R$	$A+Q$

	$\wedge$	$\vee$	
$\wedge$	$\vee$	$\times$	$\wedge$
$\vee$	$\times$	$\wedge$	$\vee$
$\times$	$\wedge$	$\vee$	

N.B. In order that the two corners may satisfy the given conditions  $A, B, C$  must be in A.P. and so also  $P, Q, R$  must be in A.P.

E: A given odd side, and given sum, construct a square.

- (i) for 11
- (ii) for 13

10	9	7
8	6	5
5	11	8

16	5	4
7	11	10
12	5	6

- (3) if the corners are 16, 5, 4 and the sum is 28
- (17) if the corners are 15, 4, 3 the columns 16, 15, 14
- if the corners are 4, 15, 14

16	3	15
4	5	11
8	10	7

7	7	8
8	7	6
7	6	5

5. To construct an oblong containing 3 rows and 4 columns.

$$A+C=2B+3D$$

A	C+D	A+2D	C+3D
B+D	B+4D	B+3D	B
C	A+D	C+2D	A+3D

✓	∧	✓	∧
x	x	x	x
∧	✓	∧	✓

Ex. Construct an oblong (i) with the diagonals  
 (ii) whose sum is 15 and all numbers are odd.

1	7	3	15
11	7	7	7
19	3	11	7

1	35	5	49
31	17	13	7
53	3	37	7

6. To construct a square containing 4 rows and 4 columns.

i. When the corners, columns and rows are all different, arrange the middle four so that the sum may be equal to half the difference between the whole sum and the sum of the corners, the middle rows and the middle columns.

ii. When the rows, columns & corners are equal

A	B	C	D	P	Q	R	S
D	C	B	A	R	S	P	Q

Add these two as  $A+P$ ,  $B+Q$  &c and fill up the other two rows. Or we may construct as the oblong in I 5.

A+P	D+S	C+Q	B+R
C+R	B+Q	A+S	D+P
B+S	C+P	D+R	A+Q
D+Q	A+R	B+P	C+S

$$A+D=B+C, P+S=Q+R$$

A+P	D+Q	D+R	A+S
B+S	C+R	C+Q	B+P
C+S	B+R	B+Q	C+P
D+P	A+Q	A+R	D+S

N.B If  $A+D=B+C$  &  $P+R=Q+S$  the extreme middle four in the 1st sq. also satisfy the given conditions.

Ex. 1. Construct for 36 and 38.

1	13	11	8
12	7	9	13
6	9	16	3
15	4	5	10

1	11	13	7
12	11	1	5
13	1	1	2
12	3	9	16

1	13	11	8
12	7	9	13
6	9	16	3
15	4	5	10

2. Construct two different squares for 66.

1	30	32	8
2	7	2	28
3	23	32	18
31	4	5	26

1	21	11	5
2	14	10	21
3	17	26	17
20	1	23	25

3. Construct two different squares for 60.

20	1	1	1
23	20	27	
17	17	14	

15	3	2
21	20	15
8	5	15

If  $m$  is a multiple of  $n$  then a square of  $m$  does  
 can be formed of different squares of  $n$  rows.  
 Exception: - The central numbers in the squares  
 of 3 rows are not different and consequently the  
 square of 6 rows cannot be formed by the above  
 method; however a regular square of 6 rows can  
 be formed by making the corners of the three-  
 rowed squares different.  
 If  $m$  is a multiple of 4 it may also be constructed  
 - ed as in I 6 (ii) second square.

16	15	14	13	12	11	10	9
7	6	5	4	3	2	1	0
17	16	15	14	13	12	11	10
8	7	6	5	4	3	2	1
18	17	16	15	14	13	12	11
9	8	7	6	5	4	3	2
19	18	17	16	15	14	13	12
10	9	8	7	6	5	4	3
20	19	18	17	16	15	14	13
11	10	9	8	7	6	5	4
21	20	19	18	17	16	15	14
12	11	10	9	8	7	6	5
22	21	20	19	18	17	16	15
13	12	11	10	9	8	7	6
23	22	21	20	19	18	17	16
14	13	12	11	10	9	8	7
24	23	22	21	20	19	18	17
15	14	13	12	11	10	9	8
25	24	23	22	21	20	19	18

16	15	14	13	12	11	10	9
7	6	5	4	3	2	1	0
17	16	15	14	13	12	11	10
8	7	6	5	4	3	2	1
18	17	16	15	14	13	12	11
9	8	7	6	5	4	3	2
19	18	17	16	15	14	13	12
10	9	8	7	6	5	4	3
20	19	18	17	16	15	14	13
11	10	9	8	7	6	5	4
21	20	19	18	17	16	15	14
12	11	10	9	8	7	6	5
22	21	20	19	18	17	16	15
13	12	11	10	9	8	7	6
23	22	21	20	19	18	17	16
14	13	12	11	10	9	8	7
24	23	22	21	20	19	18	17
15	14	13	12	11	10	9	8
25	24	23	22	21	20	19	18

8. To construct a square of odd rows & columns
- |                            |                            |
|----------------------------|----------------------------|
| A, B, C, D, E, F, G, H & C | P, Q, R, S, T, U, V, W & G |
| A, B, C, D, E, F & C       | R, S, T, U, V, W, & C      |
| A, B, C, D & C             | T, U, V, W & C             |
| A, B & C                   | V, W, & C                  |

Thus arranging the letters and adding the two sets  
 a square of any number of odds rows can be  
 formed and we can find many ways of con-  
 structing a square and the peculiarities are  
 common to all the odd squares.

A+P	E+R	D+T	C+Q	B+S
C+T	B+Q	A+S	E+P	D+R
E+S	D+P	C+R	B+T	A+Q
B+R	A+T	E+Q	D+S	C+P
D+Q	C+S	B+P	A+R	E+T

D+Q	E+S	A+P	B+R	C+T
E+R	A+T	B+Q	C+S	D+T
A+S	B+P	C+R	D+T	E+Q
B+T	C+Q	D+S	E+P	A+R
C+P	D+R	E+T	A+Q	B+S

N.B. In the 2nd square  $A+B+D+E$  must be equal

Ex. Construct a square with sides 65 and 65

17	34	1	8	15
23	3	7	16	14
4	6	13	20	22
10	12	11	25	21
11	18	25	2	9

1	2	3	4	5
6	7	8	9	10
11	12	13	14	15
16	17	18	19	20
21	22	23	24	25

2. Construct a seven row square for 170 x 170

1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31	32	33	34	35
36	37	38	39	40	41	42
43	44	45	46	47	48	49

1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31	32	33	34	35
36	37	38	39	40	41	42
43	44	45	46	47	48	49

# CHAPTER II

$$\frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \frac{1}{n+4} + \dots + \frac{1}{2n}$$

$$= \frac{n}{2n+1} + \frac{1}{2^2-2} + \frac{1}{4^2-4} + \frac{1}{6^2-6} + \dots + \frac{1}{(2n)^2-2n}$$

*Sol.*  $(2n^2-2n) = 2n(n-1) = 2n \left( \frac{n-1}{n} \right) = 2(n-1)$

$\frac{1}{2(n-1)} = \frac{1}{2} \left( \frac{1}{n-1} - \frac{1}{n} \right)$

$\frac{1}{4(n-2)} = \frac{1}{4} \left( \frac{1}{n-2} - \frac{1}{n} \right)$

$\frac{1}{6(n-3)} = \frac{1}{6} \left( \frac{1}{n-3} - \frac{1}{n} \right)$

$\dots$

$\frac{1}{2n(n-1)} = \frac{1}{2n} \left( \frac{1}{n-1} - \frac{1}{n} \right)$

$$= 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{2n-1} - \frac{1}{2n}$$

*Cor.*  $2 \log_2 2 = 1 + \frac{2}{2^3-2} + \frac{2}{4^3-4} + \frac{2}{6^3-6} + \dots$

*Sol.* By comparing  $\log_2 2$  on the above series

$2 \log_2 2 = \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots$

*Ex.* Show that  $\frac{n-1}{n+1} + \frac{n-2}{n+2} + \dots + \frac{n-n}{n+n}$

$$= 2n \left\{ \frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{3 \cdot 4 \cdot 5} + \frac{1}{5 \cdot 6 \cdot 7} + \dots + \frac{1}{(2n-1) 2n (2n+1)} \right\}$$

$$- \frac{n}{2n+1}$$

*Sol.* we have by the 1.

$\frac{1}{1 \cdot 2 \cdot 3} = \frac{1}{2} \left( \frac{1}{1 \cdot 2} - \frac{1}{2 \cdot 3} \right)$

$\frac{1}{3 \cdot 4 \cdot 5} = \frac{1}{4} \left( \frac{1}{3 \cdot 4} - \frac{1}{4 \cdot 5} \right)$

$\dots$

$\frac{1}{(2n-1) 2n (2n+1)} = \frac{1}{2n} \left( \frac{1}{(2n-1) 2n} - \frac{1}{2n(2n+1)} \right)$

By an ar: have  $\frac{1}{1 \cdot 2 \cdot 3} + \frac{1}{3 \cdot 4 \cdot 5} + \dots + \frac{1}{(2n-1) 2n (2n+1)} =$

$\frac{1}{2n} \left( \frac{1}{1 \cdot 2} + \frac{1}{3 \cdot 4} + \dots + \frac{1}{(2n-1) 2n} \right) - \frac{1}{2n(2n+1)}$

Subtracting 1 from each term on the left and

on from the right we get the result.



$$2. \frac{1}{n+1} + \frac{1}{n+2} + \frac{1}{n+3} + \dots + \frac{1}{3n+1}$$

$$\approx 1 + \frac{2}{3^2-3} + \frac{2}{6^2-6} + \frac{2}{9^2-9} + \dots + \frac{2}{(3n)^2-3n}$$

Sol. As in II 1.

$$\text{Cor. } \log 3 = 1 + \frac{2}{3^2-3} + \frac{2}{6^2-6} + \frac{2}{9^2-9} + \dots$$

Sol. Let  $x = \frac{1}{3}$  then  $\log x = \dots$  when  $x=3$   
 putting  $x=3$  in R.S. =  $\log 3$   
 it gives  $\frac{dx}{x} = \log 3$ .

$$3. \tan^{-1} \frac{1}{n+1} + \tan^{-1} \frac{1}{n+2} + \dots + \tan^{-1} \frac{1}{3n+1}$$

$$= \tan^{-1} 1 + \tan^{-1} \frac{10}{5.8} + \tan^{-1} \frac{20}{14.85} + \dots + \tan^{-1} \frac{1800}{(3n+2)(9n)}$$

Sol. As in II 2.

$$4. \left\{ \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n} \right\} + \left\{ \frac{1}{2n+1} + \frac{1}{2n+3} + \dots + \frac{1}{4n+1} \right\}$$

$$= 1 + \frac{2}{4^2-4} + \frac{2}{8^2-8} + \dots + \frac{2}{(4n)^2-4n}$$

$$= \left( 1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{4n+1} \right) + \frac{1}{2} \left( 1 - \frac{1}{2} + \frac{1}{3} - \dots - \frac{1}{2n} \right)$$

Sol. By proceeding as in II 1, R.S. =  $\frac{1}{n+1} - \frac{1}{2} = \frac{1}{2(n+1)}$

$$= \frac{1}{2} = \frac{1}{2} = \frac{1}{2(n+1)} = \frac{1}{2n} - \frac{1}{2} \left( \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n} \right)$$

$$= \left( \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{4n+1} \right) - \left( \frac{1}{2n+2} + \frac{1}{2n+4} + \dots + \frac{1}{4n} \right)$$

$$= \left( \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{4n+1} \right) + \left( \frac{1}{2n+1} + \frac{1}{2n+3} + \dots + \frac{1}{4n+1} \right)$$

$$\text{Again } = \frac{1}{2(n+1)} - \frac{1}{2} = \frac{1}{2n} - \frac{1}{2} = \frac{1}{2n} = \frac{1}{2(n+1)} + \frac{1}{2}$$

$$+ \frac{1}{2} = \frac{1}{2} - \frac{1}{2} = \frac{1}{2n} = \left( 1 + \frac{1}{2} + \dots + \frac{1}{4n+1} \right) - \left( 1 + \frac{1}{2} + \dots + \frac{1}{2n} \right)$$

$$+ \frac{1}{2} \left( 1 + \frac{1}{2} + \frac{1}{3} + \dots - \frac{1}{2n} \right) - \left( \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{2n} \right)$$

$$= \left( 1 - \frac{1}{2} + \frac{1}{3} - \dots + \frac{1}{4n+1} \right) + \frac{1}{2} \left( 1 - \frac{1}{2} + \frac{1}{3} - \dots - \frac{1}{2n} \right)$$

Cor.  $\frac{3}{2} \log_2 2 = 1 + \frac{2}{4^2-4} + \frac{2}{8^2-8} + \frac{2}{12^2-12} + \dots$

5.  $\frac{1}{3} (\frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{2n}) + (\frac{1}{2n+1} + \frac{1}{2n+3} + \dots + \frac{1}{4n+1})$   
 $= 1 + \frac{2}{6^2-6} + \frac{2}{12^2-12} + \frac{2}{18^2-18} + \dots + \frac{2}{(6n)^2-6n}$

sol. by preceding question II. the sum is  $\frac{2}{6n+1} - \frac{1}{3} = \frac{2}{3n} - \frac{1}{3} = \frac{2n}{3n} - \frac{1}{3} = \frac{2n-1}{3n} = \frac{2}{3} - \frac{1}{3n}$

Cor.  $\frac{1}{2} \log_2 3 + \frac{1}{3} \log_2 4 = 1 + \frac{2}{6^2-6} + \frac{2}{12^2-12} + \frac{2}{18^2-18} + \dots$

N.B.  $1 + \frac{2}{a^2-a} + \frac{2}{(2a)^2-2a} + \dots + \frac{2}{(an)^2-an}$

cannot be expressed as in II for all values of a except 2, 3, 4 and 6 though it can be summed up for all values of a when n becomes infinite. see chapter

Ex. 1.  $\log_2 2 = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots$

2.  $\log_2 4 = 1 - \frac{1}{2^2-2} + \frac{2}{2^2-2} - \frac{2}{2^2-2} + \dots$

3.  $\log_2 1 + \frac{2}{2^2-2} + \frac{2}{2^2-2} + \dots + \frac{2}{(2n)^2-2n}$

$= \{ 1 + \frac{2}{2^2-2} + \frac{2}{2^2-2} + \dots + \frac{2}{(2n)^2-2n} \} + \dots$

$\{ \frac{1}{2} + \frac{1}{2^2-2} + \frac{1}{2^2-2} + \dots + \frac{1}{(2n)^2-2n} \} + \dots$

$= \frac{1}{2} + \frac{1}{2^2-2} + \frac{1}{2^2-2} + \dots + \frac{1}{(2n)^2-2n} + \dots$

$= \frac{1}{2} + \frac{1}{2^2-2} + \frac{1}{2^2-2} + \dots + \frac{1}{(2n)^2-2n} + \dots$

5.  $\frac{3^2-3}{2^2-2} + \frac{2}{2^2-2} + \frac{2}{12^2-12} + \dots = \frac{1}{2} \log_2 5 - \frac{1}{2} \log_2 2$

6.  $\frac{4}{3} \log_2 2 = 1 - \frac{2}{3^2-3} + \frac{2}{6^2-6} - \frac{2}{9^2-9} + \dots$



$$7. \left\{ 1 + \frac{2}{3^2-1} + \frac{2}{5^2-1} + \frac{2}{7^2-1} + \dots + \frac{2}{(2n+1)^2-1} \right\}$$

$$= \left\{ 1 + \frac{1}{3-1} + \frac{1}{5-1} + \dots + \frac{1}{2n+1} \right\}$$

$$+ \left\{ 1 + \frac{1}{3+1} + \frac{1}{5+1} + \dots + \frac{1}{2n+1} \right\}$$

$$= \frac{2}{(2n+1)(6n+2)}$$

$$8. \tan^2 \frac{1}{2} + \tan^2 \frac{1}{3} + \dots + \tan^2 \frac{1}{13}$$

$$= \frac{1}{2} + 2 \tan^2 \frac{1}{4} + \tan^2 \frac{1}{29} + \tan^2 \frac{3}{23} + \tan^2 \frac{4}{15}$$

$$9. 2 \left( \tan^2 \frac{1}{2n} + \tan^2 \frac{1}{2n+1} + \dots + \tan^2 \frac{1}{2n+1} \right)$$

$$= \tan^2 \frac{2n}{2n+1} + \tan^2 \frac{2}{11} + \tan^2 \frac{1}{139} + \tan^2 \frac{1}{607}$$

$$+ \dots + \tan^2 \frac{2n}{2n+1} + \dots$$

$$2 \left( \tan^2 \frac{1}{2n} + \tan^2 \frac{1}{2n+1} + \dots + \tan^2 \frac{1}{2n+1} \right)$$

$$10. \tan^2 \frac{1}{n+1} + \tan^2 \frac{1}{n+2} + \dots + \tan^2 \frac{1}{2n}$$

$$= \tan^2 \frac{1}{2n+1} + \tan^2 \frac{1}{2n+3} + \dots + \tan^2 \frac{1}{2n+1}$$

$$= \frac{1}{2} + \tan^2 \frac{2}{57} + \tan^2 \frac{18}{577} + \dots + \tan^2 \frac{9n}{2n^2+2n+1}$$

$$= \tan^2 \frac{6}{157} + \tan^2 \frac{9}{107} + \dots + \tan^2 \frac{3n}{15n^2+2n+1}$$

6. If  $A_n = 3^n(n + \frac{1}{2}) - \frac{1}{2}$ , then

$$\frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{A_n}$$

$$= \left\{ 1 + \frac{1}{3^2-3} + \frac{1}{13-6} + \frac{1}{9^2-9} + \dots + \frac{2}{(3n)^2-3n} \right\}$$

$$+ (n-1) \left\{ \frac{2}{(3A_0+3)^3 - (3A_0+3)} + \frac{2}{(3A_0+6)^3 - (3A_0+6)} + \dots + \frac{2}{(3A_1)^3 - 3A_1} \right\}$$

$$+ (n-2) \left\{ \frac{2}{(3A_1+3)^3 - (3A_1+3)} + \frac{2}{(3A_1+6)^3 - (3A_1+6)} + \dots + \frac{2}{(3A_2)^3 - 3A_2} \right\}$$

+ &c to n terms.

By (1) we have

$$\frac{1}{2^2} + \frac{1}{2^3} + \dots + \frac{1}{2^{n-1}} + \frac{1}{2^n} + \frac{1}{2^{n+1}} + \dots$$

$$\frac{1}{3^2} + \frac{1}{3^3} + \dots + \frac{1}{3^{n-1}} + \frac{1}{3^n} + \frac{1}{3^{n+1}} + \dots$$

By (2) we have  $\frac{1}{3^2} = 1 + \frac{1}{3^2} + \dots + \frac{1}{3^{n-1}}$  (symmetrical series)  
 repeating this n times and then adding up  
 all the series we can get the result.

$$\text{Cor. } 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{2^n - 2}$$

$$= n + (n-1) \left( \frac{2}{3^2-3} \right) + (n-2) \left( \frac{2}{6^2-6} + \frac{2}{9^2-9} + \frac{2}{12^2-12} \right)$$

$$+ (n-3) \left( \frac{2}{15^2-15} + \frac{2}{18^2-18} + \dots + \frac{2}{37^2-37} \right) + \&c$$

to n terms.

N.B. The above theorems are very useful in finding  $\sum \frac{1}{n}$ . If  $a_1$  &  $a_n$  are very great &  $a_1, a_2, a_3, \&c$  are in A.P., then the approximate value of  $\frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n} = \frac{2n}{a_1 + a_n}$ .

$$\text{Ex. 1. } 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{100}$$

$$= 2 + \frac{1}{2} + \frac{1}{100} + \frac{1}{100} + \frac{1}{100}$$

$$\therefore 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{100} = 7\frac{1}{2} \text{ very nearly}$$

$$7. \tan^{-1} \frac{2}{(n+1)^2} + \tan^{-1} \frac{2}{(n+3)^2} + \tan^{-1} \frac{2}{(n+5)^2} + \dots \text{to } n \text{ terms}$$

$$= \tan^{-1} \frac{2n}{n^2 + 2n + 1}$$

Sol.  $\tan^{-1} \frac{2}{n} - \tan^{-1} \frac{2}{n+2} = \tan^{-1} \frac{2}{(n+1)^2}$   
 $\therefore \text{L.S.} = \tan^{-1} \frac{2}{n} - \tan^{-1} \frac{2}{n+2n} = \tan^{-1} \frac{2}{n^2 + 2n + 1}$

Cor.  $\tan^{-1} \frac{2}{(n+1)^2} + \tan^{-1} \frac{2}{(n+3)^2} + \tan^{-1} \frac{2}{(n+5)^2} + \dots = \tan^{-1} \frac{2}{n}$

Ex. 1.  $\tan^{-1} \frac{2}{(n+1)^2} + \tan^{-1} \frac{2}{(n+1)^2} + \tan^{-1} \frac{2}{(n+3)^2} + \dots = \tan^{-1} \frac{2n}{n^2 + 2n + 1}$

Sol.  $\tan^{-1} \frac{2}{(n+1)^2} + \tan^{-1} \frac{2}{(n+1)^2} + \dots = \tan^{-1} \frac{2}{n}$

$\therefore \tan^{-1} \frac{2}{(n+2)^2} + \tan^{-1} \frac{2}{(n+4)^2} + \dots = \tan^{-1} \frac{2}{n+2}$

$\therefore \tan^{-1} \frac{2}{(n+4)^2} + \tan^{-1} \frac{2}{(n+6)^2} + \dots = \tan^{-1} \frac{2}{n+4}$

N. B. If  $n < \frac{\sqrt{5}-1}{2}$  add  $\pi$  to R.S.

1.  $\tan^{-1} \frac{2}{(n+1)^2} = \tan^{-1} \frac{2}{(n+1)^2} + \tan^{-1} \frac{2}{(n+3)^2} - \dots = \tan^{-1} \frac{2}{n+1}$

2.  $\tan^{-1} \frac{2}{2(n+1)^2} + \tan^{-1} \frac{2}{2(n+3)^2} + \tan^{-1} \frac{2}{2(n+5)^2} + \dots = \tan^{-1} \frac{2}{n+1}$

3.  $\frac{\pi}{2} = \tan^{-1} \frac{1}{1} + \tan^{-1} \frac{2}{2} + \tan^{-1} \frac{3}{3} + \dots$

4.  $\frac{\pi}{4} = \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{2} + \dots = \tan^{-1} \frac{1}{1} = \tan^{-1} 1$

5.  $\frac{\pi}{8} = \tan^{-1} \frac{1}{(1+\sqrt{2})^2} + \tan^{-1} \frac{1}{(1+\sqrt{2})^2} + \dots$

6.  $\frac{\pi}{8} = \tan^{-1} \frac{1}{(1+\sqrt{2})^2} + \tan^{-1} \frac{1}{(1+\sqrt{2})^2} + \dots$

7.  $\frac{\pi}{8} = \tan^{-1} \frac{1}{(1+\sqrt{2})^2} + \tan^{-1} \frac{1}{(1+\sqrt{2})^2} + \dots$

8.  $\frac{\pi}{8} = \tan^{-1} \frac{1}{1} + \tan^{-1} \frac{1}{2} + \tan^{-1} \frac{1}{2} + \dots$

8. If  $\alpha, \beta, \gamma, \delta$  &c are the roots of the equation  $f(x) = 0$ , then  
 $f(x) = f(0) (1 - \frac{x}{\alpha})(1 - \frac{x}{\beta})(1 - \frac{x}{\gamma})$  &c. Only if the text given  
 can be  $f(x) = f(x) (\frac{1}{x-\alpha} + \frac{1}{x-\beta} + \frac{1}{x-\gamma} + \dots)$  at the end of the  
 note book is true

2.  $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma} + \frac{1}{\delta} + \dots = - \frac{f'(0)}{f(0)}$

9. i)  $\frac{\sin x}{x} = (1 - \frac{x^2}{\pi^2})(1 - \frac{x^2}{4\pi^2})(1 - \frac{x^2}{9\pi^2})$  &c

ii)  $\cos x = (1 - \frac{4x^2}{\pi^2})(1 - \frac{4x^2}{9\pi^2})(1 - \frac{4x^2}{25\pi^2})$  &c

d. The roots of the equation  $\frac{\sin x}{x} = 0$  are  
 $\pm \pi, \pm 2\pi, \pm 3\pi$  &c, and those of  $\cos x = 0$   
 are  $\pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \pm \frac{5\pi}{2}$  &c. Applying the  
 above theorem we get the result.

1.  $\frac{e^x - e^{-x}}{2x} = (1 + \frac{x^2}{\pi^2})(1 + \frac{x^2}{4\pi^2})(1 + \frac{x^2}{9\pi^2})$  &c

2.  $\frac{e^x + e^{-x}}{2} = (1 + \frac{x^2}{\pi^2})(1 + \frac{x^2}{4\pi^2})(1 + \frac{x^2}{9\pi^2})$  &c

3. change  $x$  to  $\frac{x}{2}$  in the above result

3.  $\cos \frac{x}{4} + \sin \frac{x}{4} = (1 + \frac{x}{\pi})(1 - \frac{x}{3\pi})(1 + \frac{x}{5\pi})(1 - \frac{x}{7\pi})$  &c

4.  $\frac{\sin(x+a)}{\sin a} = (1 + \frac{x}{a})(1 - \frac{x}{\pi-a})(1 + \frac{x}{\pi+a})(1 - \frac{x}{2\pi-a})$  &c

Ex. 1.  $\frac{\cos(x+a)}{\cos a} = (1 + \frac{x}{\pi+a})(1 - \frac{x}{\pi-a})(1 + \frac{x}{\pi+a})$  &c

2.  $1 + \frac{\sin x}{\sin a} = (1 + \frac{x}{a})(1 + \frac{x}{\pi-a})(1 - \frac{x}{\pi+a})$  &c

N.B If we know the value of  $(1+a_1x)(1+a_2x)(1+a_3x) \dots$   
 then it is possible to find  $(1+a_1x^n)(1+a_2x^n) \dots$

10.  $\cot x = \frac{1}{x} - \frac{1}{\pi-x} + \frac{1}{\pi+x} - \frac{1}{2\pi-x} + \frac{1}{2\pi+x} - \dots$

Sol. Equate the coeff. of  $x$  in  $\Pi$  of both.

Case 1.  $\tan x = \frac{1}{\frac{\pi}{2}-x} - \frac{1}{\frac{\pi}{2}+x} + \frac{1}{\frac{3\pi}{2}-x} - \frac{1}{\frac{3\pi}{2}+x} + \dots$

2.  $\operatorname{cosec} x = \frac{1}{x} + \frac{1}{\pi-x} - \frac{1}{\pi+x} - \frac{1}{2\pi-x} + \dots$

3.  $\sec x = \frac{1}{\frac{\pi}{2}-x} + \frac{1}{\frac{\pi}{2}+x} - \frac{1}{\frac{3\pi}{2}-x} - \frac{1}{\frac{3\pi}{2}+x} + \dots$

Sol.  $\tan x = \cot(\frac{\pi}{2}-x)$ ;  $\operatorname{cosec} x = \frac{1}{\sin(\frac{\pi}{2}-x)}$   
 and  $\sec x = \operatorname{cosec}(\frac{\pi}{2}-x)$ . Apply the same rule.

11.  $\tan^{-1} \frac{x}{a} - \tan^{-1} \frac{x}{\pi a} + \tan^{-1} \frac{x}{\pi+a} - \tan^{-1} \frac{x}{2\pi+a} + \dots$

$= \tan^{-1} \left( \frac{e^x - e^{-x}}{e^x + e^{-x}} \cot a \right)$

Sol. L.S =  $\frac{1}{2i} \log \left\{ \frac{1 + \frac{x}{\pi a}}{1 - \frac{x}{\pi a}} \cdot \frac{1 - \frac{x}{\pi+a}}{1 + \frac{x}{\pi+a}} \dots \right\}$

Case 4.

Cor. 1.  $\tan^{-1} \frac{x}{a} + \tan^{-1} \frac{x}{\pi a} - \tan^{-1} \frac{x}{\pi+a} - \tan^{-1} \frac{x}{2\pi+a} + \dots$

$= \tan^{-1} \left( \frac{e^x - e^{-x}}{2} \operatorname{cosec} a \right)$

2.  $\tan^{-1} \frac{x}{1} - \tan^{-1} \frac{x}{3} + \tan^{-1} \frac{x}{5} - \dots = \tan^{-1} \left( \frac{e^{\frac{\pi x}{2}} - 1}{e^{\pi x} + 1} \right)$

3.  $\tan^{-1} \frac{x}{1} + \tan^{-1} \frac{x}{3} - \tan^{-1} \frac{x}{5} - \dots = \tan^{-1} \left( \frac{e^{\frac{\pi x}{2}} - 1}{\sqrt{2} e^{\frac{\pi x}{2}}} \right)$

Ex. 1.  $\tan^{-1} \frac{e^x - 1}{e^x + 1} - \tan^{-1} \frac{e^{-x} - 1}{e^{-x} + 1} + \tan^{-1} \frac{e^{2x} - 1}{e^{2x} + 1} - 200 + 3$   
 $= \tan^{-1} (\tanh x \tanh 2x)$

2.  $\tan^{-1} \frac{e^x - 1}{e^x + 1} + \tan^{-1} \frac{e^{-x} - 1}{e^{-x} + 1} - \tan^{-1} \frac{e^{2x} - 1}{e^{2x} + 1} - 200$   
 $= \tan^{-1} \left( \frac{\sinh x}{\cosh x} \right)$

3.  $(1 + \frac{1}{2^2})(1 + \frac{1}{3^2})(1 + \frac{1}{4^2})(1 + \frac{1}{5^2}) \dots = \frac{1}{\pi} \operatorname{cosec}(\pi \cos \frac{\pi}{8})$

Sol.  $(1 + \frac{1}{2^2}) = (1 + \frac{1}{2})(1 - \frac{1}{2} + \frac{1}{2^2}) \dots$   
 $= (1 + \frac{1}{2})(1 - \frac{1}{2})^2 \left\{ 1 + \frac{3}{(2^2-1)^2} \right\}$

$\therefore L.S = \left(\frac{1}{2}\right)^2 \cdot \frac{4}{1} \cdot \left(\frac{3}{4}\right)^2 \cdot \frac{2}{3} \cdot \left(\frac{5}{6}\right)^2 \cdot \frac{6}{5} \dots \times (1 + \frac{1}{2^2})(1 + \frac{1}{3^2})(1 + \frac{1}{4^2}) \dots$   
 $= \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{3}{2} \cdot \frac{3}{2} \cdot \frac{5}{4} \cdot \frac{5}{4} \dots (1 + \frac{1}{2^2})(1 + \frac{1}{3^2})(1 + \frac{1}{4^2}) \dots$   
 $= \frac{1}{\pi} \operatorname{cosec} \frac{\pi \sqrt{3}}{2}$

4.  $(1 - \frac{1}{2^2})(1 - \frac{1}{3^2})(1 - \frac{1}{4^2})(1 - \frac{1}{5^2}) \dots = \frac{\operatorname{Cosh}(\pi \cos \frac{\pi}{8})}{3\pi}$

Sol.  $(1 - \frac{1}{2^2}) = (1 - \frac{1}{2})(1 + \frac{1}{2} + \frac{1}{2^2}) \dots$   
 $= (1 - \frac{1}{2})(1 + \frac{1}{2n})^2 \left\{ 1 + \frac{3}{(2n+1)^2} \right\}$  & proceed as before

12. To find convergents to a root of the eq.:

$1 = A_1 x + A_2 x^2 + A_3 x^3 + A_4 x^4 + \dots$

If  $P_n = A_1 P_{n-1} + A_2 P_{n-2} + A_3 P_{n-3} + \dots + A_{n-1} P_1$  and

$P_1 = 1$ , then  $\frac{P_n}{P_{n+1}}$  approaches  $x$  when  $n$  becomes greater and greater.



E.g. 1.  $x+x^2=1$

$x = \frac{0}{1}, \frac{1}{1} \mid \frac{1}{2}, \frac{2}{3}, \frac{3}{5}, \frac{5}{8}, \frac{8}{13}, \frac{13}{21}, \dots$

2.  $x+x^2+x^3=1$

$x = \frac{0}{1}, \frac{1}{1}, \frac{1}{2} \mid \frac{2}{3}, \frac{4}{7}, \frac{13}{13}, \frac{13}{24}, \frac{24}{44}, \dots$

3.  $x+x^3=1$

$x = \frac{0}{1}, \frac{1}{1}, \frac{1}{1} \mid \frac{1}{2}, \frac{4}{3}, \frac{3}{4}, \frac{1}{6}, \frac{6}{9}, \frac{9}{13}, \frac{13}{19}, \dots$

4.  $2x+x^2+x^3=1$

$x = \frac{0}{1}, \frac{1}{2}, \frac{2}{5} \mid \frac{5}{13}, \frac{13}{33}, \frac{33}{74}, \frac{74}{174}, \dots$

N.B. If  $\frac{p}{q}$  &  $\frac{r}{s}$  are two consecutive convergents to  $x$ , then we may take  $\frac{mp+nr}{mq+ns}$  in a suitable manner equivalent to  $x$ .

Ex. 1. Find convergents to  $\log 2$ .

Let  $\log 2 = x$ , then  $e^x = 2$

$\therefore 1 = x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$

$\therefore x = \frac{0}{1}, \frac{1}{1} \mid \frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{3}{8}, \dots$

$= \frac{1}{3}, \frac{9}{10}, \frac{52}{75}, \frac{375}{541}, \dots$

2. If  $e^{-x} = 2$ , show that the convergents to  $x$

are  $\frac{1}{2}, \frac{1}{7}, \frac{21}{67}, \frac{148}{261}, \dots$

Sol.  $1 = 2x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \dots$

$$1. \text{ If } P_0 + P_1 x + P_2 x^2 + P_3 x^3 + \dots = e^x (Q_0 + Q_1 x + Q_2 x^2 + \dots)$$

$$\text{then } P_0 f^{(0)} + P_1 f^{(1)} + P_2 f^{(2)} + P_3 f^{(3)} + P_4 f^{(4)} + \dots$$

$$= Q_0 f^{(0)} + Q_1 f^{(1)} + Q_2 f^{(2)} + Q_3 f^{(3)} + Q_4 f^{(4)} + \dots$$

Sol. The coeff<sup>s</sup> of  $f^{(n)}$  in both sides are the same as those of  $x^n$  in both sides of the first equation which are equal.

$$\text{Coef. 1. } P_0 + n P_1 x + n(n-1) P_2 x^2 + n(n-1)(n-2) P_3 x^3 + \dots$$

$$= Q_0 (1+x)^n + n Q_1 x (1+x)^{n-1} + n(n-1) Q_2 x^2 (1+x)^{n-2} + \dots$$

Sol. using  $(1+x)^n$  for  $f(x)$  in the above theorem we get

$$f^{(0)} = 1, f^{(1)} = nx, f^{(2)} = n(n-1)x^2 \dots \text{ and}$$

$$f^{(n)} = (1+x)^n, f^{(n-1)} = n(1+x)^{n-1}, f^{(n-2)} = n(n-1)(1+x)^{n-2} \dots$$

Coef. 2. If  $\phi(x) = e^x \psi(x)$ , then

$$\phi(x) f^{(0)} + \frac{\phi'(x) f^{(1)}}{1} + \frac{\phi''(x) f^{(2)}}{2} + \frac{\phi'''(x) f^{(3)}}{6} + \dots$$

$$= \psi(x) f^{(0)} + \frac{\psi'(x) f^{(1)}}{1} + \frac{\psi''(x) f^{(2)}}{2} + \frac{\psi'''(x) f^{(3)}}{6} + \dots$$

Sol. write  $\frac{\phi^{(n)}(x)}{n!}$  for  $P_n$  &  $\psi^{(n)}(x)$  for  $Q_n$  in III/1

$$2. \frac{x}{x!} + \frac{x^2}{(n+1)!} + \frac{x^3}{(n+2)!} + \frac{x^4}{(n+3)!} + \dots$$

$$= e^x \left\{ \frac{x}{x} - \frac{x^2}{x(n+1)} + \frac{x^3}{x!n(n+1)(n+2)} - \dots \right\}$$

$$\text{Sol. 1. } \frac{x}{x!} = \frac{1}{(x-1)!} \left\{ \frac{x^n}{n!} + \frac{x^{n+1}}{(n+1)!} + \frac{x^{n+2}}{(n+2)!} + \dots \right\}$$

$$= \frac{1}{x-1} \left\{ e^{x-1} \right\} = e^x \left\{ \frac{x}{x} - \frac{x^2}{x(n+1)} + \frac{x^3}{x!n(n+1)(n+2)} - \dots \right\}$$

$$\text{Sol. 2. Let } \phi(x) = \frac{x}{x!} + \frac{x^2}{(n+1)!} + \frac{x^3}{(n+2)!} + \dots$$

then,  $n\phi(x) = x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$   
 and  $x\phi(x+1) = \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$

$\therefore n\phi(x) + x\phi(x+1) = x + x^2 + \frac{x^3}{2!} + \dots = x e^x$

$\therefore \phi(x) = e^{-x} \frac{x}{n} - \frac{x}{n} \phi(x+1) = e^{-x} \frac{x}{n} - e^{-x} \frac{x^2}{n(n+1)} + \frac{x^3}{n(n+1)} \phi(x+2)$   
 $\&c \&c$

Cor. 1.  $\frac{f(x)}{n} + \frac{f'(x)}{(n+1)2!} + \frac{f''(x)}{(n+2)3!} + \dots$   
 $= \frac{f(x)}{n} - \frac{f'(x)}{n(n+1)} + \frac{f''(x)}{n(n+1)(n+2)} - \dots$

Cor. 2.  $\frac{x}{1!} + (1+\frac{1}{2})\frac{x^2}{2!} + (1+\frac{1}{2}+\frac{1}{3})\frac{x^3}{3!} + \dots$   
 $= e^x (\frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} - \dots)$

Sol. By III (1) we have  $\frac{x^2}{(n+1)2!} + \frac{x^3}{(n+2)3!} + \dots$

$= \frac{x^2}{(n+1)2!} + \frac{x^3}{(n+2)3!} + \dots$

Equating the coeff of  $x^3$  on both sides we get the result.

3. If  $\frac{1^n}{1!}x + \frac{2^n}{2!}x^2 + \frac{3^n}{3!}x^3 + \dots = e^x f(x)$ , then

$\frac{x}{n+1} - \frac{x^2}{(n+1)(n+2)} + \frac{x^3}{(n+1)(n+2)(n+3)} - \dots$   
 $= \frac{f(x)}{n} - \frac{f'(x)}{n^2} + \frac{f''(x)}{n^3} - \frac{f'''(x)}{n^4} + \dots$

Sol. By III 2 we have  $\frac{x}{n+1} + \frac{x^2}{(n+2)2!} + \frac{x^3}{(n+3)3!} + \dots$

$= e^{-x} \left\{ \frac{x}{n+1} - \frac{x^2}{(n+1)(n+2)} + \frac{x^3}{(n+1)(n+2)(n+3)} - \dots \right\}$

$= \frac{1}{n} \left( \frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right) - \frac{1}{n^2} \left( \frac{x^2}{1!} + \frac{x^3}{2!} + \dots \right)$

$+ \frac{1}{n^3} \left( \frac{x^3}{1!} + \frac{x^4}{2!} + \dots \right) - \dots$

$= e^{-x} \left\{ \frac{f(x)}{n} - \frac{f'(x)}{n^2} + \frac{f''(x)}{n^3} - \dots \right\}$

2.  $e^{ax} = 1 + \frac{ax}{1} f(x) + \frac{a^2}{2} f'(x) + \frac{a^3}{6} f''(x) + \dots$   
 Sol.  $e^{ax} = 1 + x a + \frac{x^2}{2} a^2 + \frac{x^3}{6} a^3 + \dots$  The coeff.  
 of  $a^n$  is  $\frac{f(x)}{n!} \left\{ \frac{1}{1} x + \frac{2}{2} x^2 + \frac{3}{6} x^3 + \dots \right\} = \frac{e^x}{n!} f(x)$

$\therefore e^{ax} = e^x \left\{ 1 + \frac{ax}{1} f(x) + \frac{a^2}{2} f'(x) + \frac{a^3}{6} f''(x) + \dots \right\}$   
 5.  $f(x) = x \left\{ 1 + n f(x) + \frac{n(n-1)}{2} f'(x) + \frac{n(n-1)(n-2)}{6} f''(x) + \dots \right\}$

Sol. Differentiating both sides in III, w.r.t. res-  
 pect to  $a$ ,  $x e^a e^{-ax} = f(x) + \frac{a}{1} f'(x) + \frac{a^2}{2} f''(x) + \dots$   
 $= x e^a \left\{ 1 + \frac{a}{1} f(x) + \frac{a^2}{2} f'(x) + \dots \right\}$ . By matching the  
 coeff. of  $a^n$  we get the result.

con. The above result may be written thus

$f(x), f'(x), f''(x), f'''(x), \dots, f^{(n)}(x)$

$\left. \begin{matrix} a_0 & b_0 & c_0 & d_0 \\ & a_1 & b_1 & c_1 \\ & & a_2 & b_2 \\ & & & a_3 \end{matrix} \right\}$

These are successive diff.  $a_n$  being equal to  $x f^{(n)}(x)$ .

6. If  $f(x) = \phi_1^{(n)} x + \phi_2^{(n)} x^2 + \phi_3^{(n)} x^3 + \dots + \phi_{n+1}^{(n)} x^{n+1}$   
 then  $\frac{\phi_1^{(n)}}{1} + \frac{\phi_2^{(n)}}{2} + \frac{\phi_3^{(n)}}{3} + \dots = \frac{n^n}{(n-1)}$

Sol.  $e^x f(x) = e^x \left\{ \phi_1^{(n)} x + \phi_2^{(n)} x^2 + \dots + \phi_{n+1}^{(n)} x^{n+1} \right\}$   
 But  $e^x f(x) = \frac{1}{1} x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$   
 Equating the coeff. of  $x^n$  in both sides we get the result.

7.  $\phi_{n+1}^{(n)} \frac{1}{n} = (n+1) - n \cdot n + \frac{n(n-1)}{2} (n-1) - \frac{n(n-1)(n-2)}{6} (n-2) + \dots$

Sol.  $f(x) = \phi_1(n)x + \phi_2(n)x^2 + \phi_3(n)x^3 + \dots$

$= e^{-x} \left\{ \frac{1}{10}x + \frac{7}{11}x^2 + \frac{5}{12}x^3 + \dots \right\}$

Equating the coeffts of  $x^{n+1}$  we can get the result

8.  $\phi_n(n+1) = n \phi_n(n) + \phi_n(n)$

Sol.  $\phi_n(n+1) = \frac{1}{(n+1)!} \left\{ n^{n+1} - (n-1)(n-2)^{n+1} + \frac{(n-1)(n-2)(n-3)^{n+1}}{1!} - \dots \right\}$

$\therefore \phi_n(n+1) - \phi_n(n) = \frac{1}{(n+1)!} \left\{ n \cdot n^n - n(n-1)(n-1)^n + \frac{n(n-1)(n-2) \cdot n^{n-1}}{1!} - \dots - n \cdot n^n \right\} = n \phi_n(n)$

Cor. The above theorem may be written thus  
Write under each term the product of the coefft. and the index of x of that term together with the coefft. of the preceding one.

$f_0(x) = x$

$f_1(x) = x + x^2$

$f_2(x) = x + 3x^2 + x^3$

$f_3(x) = x + 7x^2 + 6x^3 + x^4$

$f_4(x) = x + 15x^2 + 25x^3 + 10x^4 + x^5$

$f_5(x) = x + 31x^2 + 90x^3 + 65x^4 + 15x^5 + x^6$

$f_6(x) = x + 63x^2 + 301x^3 + 350x^4 + 140x^5 + 21x^6 + x^7$

Ex. 1. If  $\frac{a_1}{n+1} + \frac{a_2}{(n+1)(n+2)} + \frac{a_3}{(n+1)(n+2)(n+3)} - \dots - \frac{a_n}{n!} = \frac{F(n)}{n!}$

Show that  $F(n) = \phi_1(n)a_1 + \phi_2(n)a_2 + \phi_3(n)a_3 + \dots$

2. Show that  $\phi_{n+1}(n)$  is the coefft. of  $\frac{x^n}{n!}$  in  $\frac{e^x}{1-x} (e^x - 1)$

Sol. By III 7 we have  $\phi_{n+1}(n) = \frac{1}{(n+1)!} \left\{ (n+1)^{n+1} - \frac{a_1(n+1)}{1!} (n-1)^{n+1} - \dots \right\}$

$= \frac{(n+1)^{n+1}}{1!} - \frac{a_1(n+1)(n-1)^{n+1}}{1!} - \dots$

= the coeff. of  $x^n$  in  $\{e^{x(n+1)}\} = e^{x(n+1)} = e^{nx} + \frac{n(n-1)}{1!} e^{x(n-1)} + \dots$

= that of  $\frac{x^n}{1!}$  in  $e^x (e^x)^n$

3. 
$$\frac{d f_{n+1}(x)}{dx} = n f_{n-1}(x) + \frac{n(n-1)}{1!} f_{n-3}(x) + \frac{n(n-1)(n-2)}{1!} f_{n-4}(x) + \dots$$

Sol. Differentiating both sides in III, with respect to  $x$  and proceeding as in III, by differentiating the result with respect to  $x$  and equating the coeffs. we get the result.

4. 
$$\int f_n(x) dx + \frac{1}{2} f_n(x) = \frac{f_{n+1}(x)}{n+1} + \frac{\beta_2}{1!} x f_n(x) - \beta_4 \frac{n(n+1)(n-2)}{1!} f_{n-3}(x) + \beta_6 \frac{n(n+1)(n-1)(n-3)(n-4)}{1!} f_{n-5}(x) - \dots$$

Sol. Integrating both sides in III, with respect to  $x$  we have  $\frac{1}{x} \{1 + \frac{\beta_2}{1!} f_1(x) + \frac{\beta_4}{1!} f_3(x) + \frac{\beta_6}{1!} f_5(x) + \dots\}$

$$= \frac{1}{x} + 2 + \frac{3}{x} \{1 + \frac{\beta_2}{1!} f_1(x) + \frac{\beta_4}{1!} f_3(x) + \dots\} + \dots$$

Equating the coeffs of  $x^0$

5. i. If  $\frac{x^n}{10} + \frac{x^n}{4} + \frac{3x^n}{12} + \frac{4x^n}{15} + \dots = e A_n$

show that  $A_0=1, A_1=2, A_2=5, A_3=16, A_4=52, A_5=203$

$A_6=877, A_7=4140, A_8=21147 \dots$

ii. If  $-\frac{x^n}{10} + \frac{x^n}{4} - \frac{3x^n}{12} + \frac{4x^n}{15} - \dots = \frac{A_n}{e}$  show that

$A_0=-1, A_1=0, A_2=-1, A_3=1, A_4=-2, A_5=-9, A_6=-9, A_7=50, A_8=267 \dots$

Sol. 2.  $1+1, 5 = 1+2+2, 15 = 1+2+3+2+5, 52 = 1+4+6+6+16$   
 $+6+6+4+5+15 \dots$  Similarly for the final one.

24

N.B.	1	2	5	15	52	203	877	-1	0	1	-2	-9	-9	50
		1	3	10	37	151	674		1	1	-3	-7	0	59
			2	7	27	114	523			0	-1	-3	-4	57
				5	20	87	409				-1	-2	-1	52
					15	67	322					-1	1	41
						52	235						2	29
							203							18
														9

6. *Illustration*

$$(i) \frac{x^2}{10} + \frac{x^3}{10} + \frac{x^4}{12} + \frac{x^5}{12} + \dots = 3\left(\frac{x^2}{10} + \frac{x^3}{10} + \frac{x^4}{12} + \dots\right)$$

$$(ii) \frac{x^2(x^2+1)}{10} + \frac{x^3(x^2+1)}{12} + \frac{x^4(x^2+1)}{12} + \dots = 4x\left(\frac{x^2}{10} + \frac{x^3}{10} + \frac{x^4}{12} + \dots\right)$$

$$(iii) \frac{x^2}{10} - \frac{x^3}{10} + \frac{x^4}{12} - \frac{x^5}{12} + \dots = \frac{x^2}{10} - \frac{x^3}{10} + \frac{x^4}{12} - \frac{x^5}{12} + \dots$$

$$(iv) \frac{x^6}{10} - \frac{x^6}{10} + \frac{x^6}{12} - \frac{x^6}{12} + \dots = \frac{x^6}{10} - \frac{x^6}{10} + \frac{x^6}{12} - \frac{x^6}{12} + \dots$$

$$(v) \frac{x^2(x^2+1)(x^2+1)}{10} - \frac{x^3(x^2+1)(x^2+1)}{12} + \frac{x^4(x^2+1)(x^2+1)}{12} - \dots$$

$$= x\left(\frac{x^2}{10} - \frac{x^3}{10} + \frac{x^4}{12} - \frac{x^5}{12} + \dots\right)$$

9.  $\int (a+b)^n \frac{x}{10} + (a+3b)^n \frac{x^2}{12} + (a+3b)^n \frac{x^3}{12} + \dots = e^x f(x)$

then i.  $\frac{F_0(x)}{n} - \frac{F_1(x)}{n^2} + \frac{F_2(x)}{n^3} - \dots$

$$= \frac{x}{n+a+b} - \frac{x^2}{(n+a+b)(n+a+3b)} + \frac{x^3}{(n+a+b)(n+a+3b)(n+a+5b)} - \dots$$

ii.  $F_0(x) + \gamma F_1(x) + \frac{\gamma^2}{12} F_2(x) + \frac{\gamma^3}{12} F_3(x) + \dots = x e^{\gamma(a+b)} e^{x^2}$

iii.  $F_{n+1}(x) - (a+b)F_n(x) = bx \left\{ F_n(x) + \frac{n}{12} F_{n-1}(x) + \frac{n(n-1)}{12} F_{n-2}(x) + \dots \right\}$

iv.  $\int F_n(x) = \phi_1(n)x + \phi_2(n)x^2 + \phi_3(n)x^3 + \dots$ , then

$$\frac{\phi_1(n)}{10} + \frac{\phi_2(n)}{12} + \frac{\phi_3(n)}{12} + \dots = \frac{(a+3b)^n}{10}$$

N.B.  $\int F_{n+1}(x) - (a+b)F_n(x) = \psi_n(x)$ , then

$$\psi_0(x), \psi_1(x), \psi_2(x), \psi_3(x), \psi_4(x)$$

$a_1$	$b_1$	$c_1$	$d_1$
$a_2$	$b_2$	$c_2$	
$a_3$	$b_3$	$c_3$	
$a_4$	$b_4$	$c_4$	

These are successive diff. of 6 times the previous term being subtracted from each term and  $a_n$  being equal to  $b \times F_n(x)$ .

$$V. \phi_n^{(n-1)}(x) = (a+nb)^n - \frac{n}{1} (a+n-1)b^n + \frac{(n-1)(n-2)}{1 \cdot 2} (a+n-2)b^n - \frac{(n-1)(n-2)(n-3)}{1 \cdot 2 \cdot 3} (a+n-3)b^n + \dots$$

$$vi. \phi_n^{(n+1)} = (a+nb)\phi_n^{(n)} + b\phi_{n+1}^{(n)}$$

N.B. Write under each term the product of  $a+nb$   $n$  being the index of  $x$ , and the coefft. of  $x$  of that term together with  $b$  times the coefft. of the preceding one

$$F_0(x) = x$$

$$F_1(x) = (a+b)x + b x^2$$

$$F_2(x) = (a+b)^2 x + b(2a+3b)x^2 + b^2 x^3$$

$$F_3(x) = (a+b)^3 x + b\{3(a+b)(a+2b) + b^2\} x^2 + 3b^2(a+2b)x^3 + b^3 x^4$$

$$F_4(x) = (a+b)^4 x + b\{2(a+b)(a+2b) + b^2\}(2a+3b)x^2 + b^2\{6(a+2b)^2 + b^2\} x^3 + 3b^2(2a+5b)x^4 + b^4 x^5$$

vii.  $\phi_n^{(n)}$  is the coefft. of  $\frac{x^n}{n!}$  in  $\frac{e^{x(a+b)}}{n!} (e^{bx})^n$

$$Ex. i. \frac{1^3+1^2}{1^6} = \frac{3^3+3^2}{1^6} + \frac{5^3+5^2}{1^6} - \dots = 0$$

$$ii. \frac{1^4}{1^6} + \frac{2^4}{1^6} + \frac{3^4}{1^6} + \frac{4^4}{1^6} + \dots = 2 \left( \frac{1^4}{1^6} + \frac{3^4}{1^6} + \frac{5^4}{1^6} + \dots \right)$$

$$iii. \frac{1^5+1^4}{1^6} - \frac{2^5+2^4}{1^6} + \frac{3^5+3^4}{1^6} - \dots = \frac{1^5}{1^6} - \frac{3^5}{1^6} + \frac{5^5}{1^6} - \dots$$

$$iv. \frac{1^6}{1^6} - \frac{2^6}{1^6} + \frac{3^6}{1^6} - \frac{4^6}{1^6} + \dots = \left(1 - \frac{1}{1^6} + \frac{1}{2^6} - \frac{1}{3^6} + \dots\right)$$



10.  $\phi(x) + \frac{x}{1!} \phi'(1) + \frac{x^2}{2!} \phi''(2) + \frac{x^3}{3!} \phi'''(3) + \dots = e^x \phi_\infty(x)$   
 where  $\phi_n(x) = \phi_{n-1}(x) + \frac{a}{1!n} \phi_{n-1}'(x) + \frac{a^2}{2!(1!)^2} \phi_{n-1}''(x) + \dots$   
 $\frac{a^3}{3!(1!)^3} \phi_{n-1}'''(x) + \dots$  ad. inf. where  $\phi_1(x) = \phi(x)$ ,  
 $\phi''(x)$  is the  $n$ th diff. coeff. of  $\phi(x)$  and  $a$  is  
 ultimately made equal to  $x$ .

Sol.  $\phi(x) + \frac{x}{1!} \phi'(1) + \frac{x^2}{2!} \phi''(2) + \frac{x^3}{3!} \phi'''(3) + \dots$   
 $= e^x \left\{ \phi(1) + \frac{\phi'(1)}{1!} x + \frac{\phi''(2)}{2!} x^2 + \frac{\phi'''(3)}{3!} x^3 + \dots \right\}$   
 $= e^x \left[ \phi(1) + \frac{x}{2} \phi''(2) + \left\{ \frac{x}{1} \phi'(1) + \frac{x^2}{2} \phi''(2) \right\} + \right.$   
 $\left. \left\{ \frac{x^2}{2!} \phi''(2) + \frac{x^2}{1!2} \phi'(1) + \frac{x^3}{3!} \phi'''(3) \right\} + \right.$   
 $\left. \left\{ \frac{x}{1!3} \phi'(1) + \frac{5}{144} x^2 \phi''(2) + \frac{x^3}{4!} \phi'''(3) + \frac{x^4}{3!2} \phi^{(4)}(4) \right\} \right.$   
 $\left. + \left\{ \frac{x}{1!4} \phi'(1) + \frac{x^2}{2!4} \phi''(2) + \frac{7x^3}{576} \phi'''(3) + \frac{2x^4}{27!} \phi^{(4)}(4) + \frac{x^5}{1040} \phi^{(5)}(5) \right\} \right.$   
 $\left. + \dots \dots \dots \right]$

Collecting the last terms, the last but one terms  
 we can get the result.

Cor. If  $x$  is great and  $\phi''(x)$  can be neglected, then  
 $e^{-x} \left\{ \phi(1) + \frac{x}{1!} \phi'(1) + \frac{x^2}{2!} \phi''(2) + \dots \right\} = \phi(x) + \frac{x}{2} \phi''(x)$   
 very nearly.

As in the above solution neglecting the third  
 and the other terms we get  $\phi(1) = \frac{x}{1} \phi'(1) + \frac{x^2}{2} \phi''(2) + \dots$   
 $= e^x \left\{ \phi(1) + \frac{x}{2} \phi''(2) \right\}$

Ex. 1. Show that  $\log_e \left( \frac{x}{11} \sqrt{1} + \frac{x^2}{12} \sqrt{2} + \frac{x^3}{13} \sqrt{3} + \dots \right)$   
 $= x + \frac{1}{2} \log_e x - \frac{1}{8x} - \frac{1}{16x^2}$  very nearly.

2.  $e^{-x} \left( \frac{x}{11} \log_e 2 + \frac{x^2}{12} \log_e 3 + \frac{x^3}{13} \log_e 4 + \dots \right)$   
 $= \log_e x + \frac{1}{2x} + \frac{1}{12x^2}$  very nearly.

3.  $\log_e \left\{ \phi(10) + \frac{100}{11} \phi(11) + \frac{100^2}{12} \phi(12) + \frac{100^3}{13} \phi(13) + \dots \right\}$   
 $= 100 + \log_e \frac{\phi(110) + \phi(90)}{2}$  nearly.

4. Show that  $\frac{x}{11} + \frac{x^2}{12} \left(1 + \frac{1}{2}\right) + \frac{x^3}{13} \left(1 + \frac{1}{2} + \frac{1}{3}\right) + \dots$   
 $= e^x (c + \log_e x)$  very nearly where  $c$  is the constant of the series  $1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{x}$ .

11. If  $e^{A_1 x + A_2 \frac{x^2}{2} + A_3 \frac{x^3}{3} + \dots}$   
 $= P_0 + P_1 x + P_2 x^2 + P_3 x^3 + \dots$

then  $P_n = A_1 P_{n-1} + A_2 P_{n-2} + A_3 P_{n-3} + \dots$  to  $n$  terms  
 where  $P_0 = 1$ .

*Sol. - Take logarithm of both sides then differentiate then equate the coeffs.*

Cor. If  $P_n = a_1^n + a_2^n + a_3^n + \dots + a_n^n$  and  $P_n$  denotes the sum of the products of  $a_1, a_2, a_3, \dots, a_n$  taken  $r$  at a time, then

$n P_n = S_1 P_{n-1} - S_2 P_{n-2} + S_3 P_{n-3} - \dots$  where  $P_0 = 1$ .

*Sol. Apply the above theorem in  $(1-x)(1+a_1 x)(1+a_2 x) \dots$*

28. 12. If  $n^2 + \frac{(n+1)^{2+1}}{a^{1+1}} + \frac{(n+2)^{2+2}}{a^{1+2}} + \frac{(n+3)^{2+3}}{a^{1+3}} + \dots = F_n(n)$ , then

$$F_{n+1}(n) = n F_n(n) + \frac{1}{a} F_{n+1}(n).$$

$$\begin{aligned} \text{Sol. } F_{n+1}(n) &= n^{2+1} + \frac{(n+1)^{2+1}}{a^{1+1}} + \frac{(n+2)^{2+2}}{a^{1+2}} + \dots \\ &= n \left\{ n^2 + \frac{(n+1)^{2+1}}{a^{1+1}} + \frac{(n+2)^{2+2}}{a^{1+2}} + \dots \right\} \\ &\quad + \frac{1}{a} \left\{ (n+1)^{2+1} + \frac{(n+2)^{2+2}}{a^{1+1}} + \frac{(n+3)^{2+3}}{a^{1+2}} + \dots \right\} \\ &= n F_n(n) + \frac{1}{a} F_{n+1}(n) \end{aligned}$$

We see from this identity that if we are able to find the sum for one value of  $n$  we can sum up the series for all values of  $n$ .

N.B.  $F_n(n)$  is convergent when  $a > e$  or  $a < e$  according as  $n$  is positive or not.

13. If  $x = a \log_e x$ , then  $\frac{x^n}{n} = F_{-1}(n)$ .

$$\text{Sol. Let } f(x) = \left(1 + \frac{x}{a} + \frac{x(x+1)}{a^2} + \frac{x(x+1)(x+2)}{a^3} + \dots\right)$$

multiplying up by  $(x)$  we get

$$x f(x) = \left\{ \frac{x^n}{n} \right\} \dots \text{ Let } f(1) = 2, \text{ then } \frac{x^n}{n} = \frac{f(x)}{x}$$

$$\text{Let } x=1, \text{ when } n=0, = \frac{1}{a} + \frac{1}{a^2} + \frac{1}{a^3} + \dots$$

$$= \frac{1}{a} \left(1 + \frac{1}{a} + \frac{1}{a^2} + \dots\right) = \frac{1}{a} f(1) = \frac{2}{a}$$

$$\therefore \frac{x^n}{n}, \text{ when } n=0 = \frac{2}{a} \text{ or } x = a \log_e x.$$

N.B. The minimum value of  $\frac{x}{\log_e x} = e$ ; if  $a = e$ ,  $f(x) = e^{-x}$   $f(x)$  is convergent if  $a > e$  & divergent if  $a < e$ .

$$e^{ax} = 1 + \frac{x}{1} e^{ax} + \frac{x(x+2n)}{e^{2n} 2!} + \frac{x(x+2n)^2}{e^{3n} 3!} + \dots$$

If  $ayx^p - x^q + 1 = 0$ , then

$$= 1 + \frac{n}{1} a + \frac{n(n+2p-q)}{2} a^2 + \frac{n(n+3p-q)(n+3p-2q)}{6} a^3$$

$$+ \frac{n(n+4p-q)(n+4p-2q)(n+4p-3q)}{24} a^4 + \dots$$

Corollary of Th. 13.

Cor. 1.  $\left(\frac{2}{1+\sqrt{1-4x}}\right)^n = 1 + nx + \frac{n(n+3)}{2} x^2 + \frac{n(n+4)(n+5)}{6} x^3$

$$+ \frac{n(n+5)(n+6)(n+7)}{24} x^4 + \dots$$

Cor. 2.  $(x+\sqrt{1+x^2})^n = 1 + \frac{n}{1} x + \frac{n^2}{2} x^2 + \frac{n(n^2-1)}{6} x^3$

$$+ \frac{n^2(n^2-4)}{24} x^4 + \frac{n(n^2-1)(n^2-9)}{120} x^5 + \dots$$

15.  $1 + \frac{1}{1} e^{-(1+\frac{x^2}{2})} + \frac{3}{2} e^{-2(1+\frac{x^2}{2})} + \frac{4^2}{12} e^{-3(1+\frac{x^2}{2})}$

$$+ \frac{5^3}{24} e^{-4(1+\frac{x^2}{2})} + \dots$$

$$= e^{-1-x} + \frac{x^2}{3} - \frac{x^3}{36} - \frac{x^4}{270} - \dots + \frac{11}{1080} x^4$$

$$= e^{1+\frac{x^2}{2}} \left(1 - x + \frac{x^2}{3} - \frac{x^3}{36} - \frac{x^4}{270} - \dots\right) + \frac{11}{1080} x^4$$

Let  $e^n = 1 + \frac{1}{1} ne^{-n} + \frac{n^2}{2} e^{-2n} + \dots$

$$e^{n+1} = 1 + \frac{1}{1} e^{-(n+1)} + \frac{1}{2} e^{-2(n+1)} + \dots$$

Let  $n + a^n = 1 + \frac{x^2}{3}$ . Solve the equation and find  $a$ .

N.B. This result is useful in finding the numerical value of  $F_n(n)$  when  $a$  approaches  $e$ .

$$E. 1. z^n = 1 + \frac{m}{z^n} + \frac{m(m+2n-1)}{z^{2n}} + \frac{m(m+1)(m+2)(m+3-1)}{z^{3n}} + \dots$$

2. Find  $x^x$  when  $\frac{(\log x)^m}{x} = a$ . Sol. Let  $x^m = y$ .

3. Find  $x$  in terms of  $a$  in each of the following

i.  $x^a = e^{\pm x}$  Sol.  $a \log x = \pm x$ .

ii.  $x^a = a^{\pm x}$ ; Sol.  $a \log x = \pm x \log a$ .  $\frac{x}{\log x} = \pm \frac{a}{\log a}$ .

iii.  $x = a e^{\pm x}$ ; Sol. Let  $x = \log y$ , then  $\log y = a e^{\pm \log y}$ .

iv.  $x = a^{\pm x}$ ; Sol. Let  $x = \log y$ , then  $\log y = a^{\pm \log y}$ .

v.  $x^{\pm x} = a$ ; Sol. Let  $x = \log y$ , then  $y = a^{\pm \log y}$ .

vi.  $x e^{\pm x} = a$ ; Sol. Let  $x = \log y$ , then  $\log y = a y^{\pm 1}$ .

vii.  $e^{\pm x} = a$ ; Sol. Let  $x = \log y$ , then  $e^{\pm \log y} = a$ .

viii.  $x \pm \log x = a$ ; Sol. Let  $x = \log y$ , then  $\log y \pm \log y = a$ .

4. Use the above to find the values of the following for numerical values of  $e$ .

i.  $x^{\pm x} = \sqrt{e}$ , then  $x^{\pm x} = \sqrt{e}$ .

ii.  $x \pm e^{\pm x} = \sqrt{e}$ , then  $x \pm e^{\pm x} = \sqrt{e}$ .

iii.  $\log_e \{ x \log_e [x \log_e (x \text{ mod } i)] \}$

iv.  $\pm \log_e \{ x \pm \log_e [x \pm \log_e (x \pm \sqrt{e})] \}$ .

16. Writing  $x^n \phi_n(x)$  for  $F_n(x)$ , we have

$$\phi_{n+1}(x) - \log_e x \phi_n(x) = n \phi_n(x).$$

Case I If  $a$  is positive

$$\text{Let } \phi_n(x) = \frac{\Psi_1(a, n)}{(1 - \log_e x)^{a+1}} + \frac{\Psi_2(a, n)}{(1 - \log_e x)^{a+2}} + \dots + \frac{\Psi_{n+1}(a, n)}{(1 - \log_e x)^{a+n+1}}$$

$$\text{Then } n \Psi_t(a, n) + \Psi_{t-1}(a+1, n+1) = \Psi_t(a+1, n+1) + \Psi_{t-1}(a, n)$$

Case II If  $a$  is negative, the terms in R.S continue as far as the term independent of  $(1 - \log_e x)$ .

$$\phi_{-1}(n) = \frac{1}{n}$$

$$\phi_{-2}(n) = \frac{1 - \log_e x}{n(n+1)} + \frac{1}{n^2(n+1)}$$

$$\phi_{-3}(n) = \frac{(1 - \log_e x)^2}{n(n+1)(n+2)} + \frac{(3n+2)(1 - \log_e x)}{n^2(n+1)^2(n+2)} + \frac{3n+2}{n^3(n+1)^3(n+2)}$$

$$\phi_0(n) = \frac{1}{1 - \log_e x}$$

$$\phi_1(n) = \frac{n-1}{(1 - \log_e x)^2} + \frac{1}{(1 - \log_e x)^3}$$

$$\phi_2(n) = \frac{(n-1)(n-2)}{(1 - \log_e x)^3} + \frac{(n-1)(n-2)(\frac{1}{n-2} + \frac{2}{n-1})}{(1 - \log_e x)^4} + \frac{1 \cdot 3}{(1 - \log_e x)^5}$$

$$\phi_3(n) = \frac{(n-1)(n-2)(n-3)}{(1 - \log_e x)^4} + \frac{(n-1)(n-2)(n-3)(\frac{1}{n-3} + \frac{2}{n-2} + \frac{3}{n-1})}{(1 - \log_e x)^5} + \frac{15n-35}{(1 - \log_e x)^6} + \frac{1 \cdot 3 \cdot 5}{(1 - \log_e x)^7}$$

Cor. 1.  $e^x = (1 - x) \left\{ 1 + \frac{x+n}{e^{2n}} + \frac{(x+2n)^2}{e^{4n}} + \frac{(x+3n)^3}{e^{6n}} + \dots \right\}$

2.  $\Psi_1(a, n) + \Psi_2(a, n) + \Psi_3(a, n) + \dots$  as far as the terms cease to continue in  $\phi_n(n) = n^a$ .

3d. L.S. =  $\phi_n(n)$  when  $x=1$  L.S.  $F_n(n)$  when  $x \neq 1$ , L.S.  $F_n(n)$  when  $a = \infty = n^a$ .

17. To expand  $x^m$  in ascending powers of  $h$  when  $x^x = a^a e^h$ .

32. Let  $\frac{x-a}{a} = \frac{A_1}{1} \cdot \frac{h}{a} - \frac{A_2}{2!} \cdot \left(\frac{h}{a}\right)^2 + \frac{A_3}{3!} \left(\frac{h}{a}\right)^3 - \dots$  &  $n = \frac{1}{1 + \log \frac{x}{a}}$

then  $A_n - n(n-2)A_{n-1} = n \left\{ n A_1 A_{n-1} + \frac{n(n-1)}{2} A_2 A_{n-2} + \frac{n(n-1)(n-2)}{6} A_3 A_{n-3} + \dots \right\}$  the last term being

$\frac{1}{2!} A_{n-1}$  or  $\frac{1}{2 \left(\frac{h}{a}\right)^2} A_2$  according as  $n$  is odd or even

$A_1 = n$   
 $A_2 = n^3$

$A_3 = 3n^5 + n^4$

$A_4 = 15n^7 + 10n^6 + 3n^5$

$A_5 = 105n^9 + 105n^8 + 40n^7 + 6n^6$

$A_6 = 945n^{11} + 1260n^{10} + 700n^9 + 196n^8 + 21n^7$

$A_7 = 10395n^{13} + 17225n^{12} + 12600n^{11} + 5068n^{10} + 1148n^9 + 120n^8$

Multiply the power and the coeff. & write under each term the sum of this product and  $(n-1)$  times the coeff. of the preceding term where  $n$  is the suffix of  $A$ .

N.B. For  $\frac{a}{x}$  take  $(n+1)$  times the coeff.; for  $\log \frac{x}{a}$  take  $n$  times the coeff. and generally for  $\left(\frac{x}{a}\right)^m$  take  $(n-m)$  times the coeff.

Ex. 1. Show that the sum of the coeff. of  $A_n = (n-1)!$

*sol.* Put for  $a$ , then  $x^x = e^h$ .

Let  $x = y$ , then  $y^y = a e^{-h}$  or  $\log y = \frac{h}{y}$ .

$y = x = 1 + h - \frac{1}{2} h^2 + \frac{1}{6} h^3 - \frac{1}{24} h^4 + \dots$

the sum of the coeff. of  $A_n = (n-1)!$

2. To expand  $x$  in ascending powers of  $h$  when  $x^x = e^h \log a$ .

*sol.* Let  $x = y$ ; then  $y^y = e^{-h} \left(\frac{1}{a}\right)^{\frac{1}{y}}$ .

Let  $F_1(x) = e^x - 1$ ,  $F_2(x) = e^{e^x} - 1$

$F_3(x) = e^{e^{e^x}} - 1$ ,  $F_4(x) = e^{e^{e^{e^x}}} - 1$

Let us try to find the expansion in powers of  $x$  and in ascending powers of  $x$

1. Let  $e^{-1} = F_n(x) = x \phi_1(x) + x^2 \phi_2(x) + x^3 \phi_3(x) + \dots$   
 $= f_0(x) + x f_1(x) + x^2 f_2(x) + x^3 f_3(x) + \dots$

then  $\log_2 [1 + \log_2 \{1 + \log_2 (1 + \dots + \log_2 (1+x))\}]$   $\log_2$  taken  $n$  times

$= F_{-n}(x) = x \phi_1(-x) + x^2 \phi_2(-x) + x^3 \phi_3(-x) + \dots$   
 $= f_0(x) - x f_1(x) + x^2 f_2(x) - x^3 f_3(x) + \dots$

Sol. we have  $e^{F_n(x)} = F_{n+1}(x)$ ;  $\therefore F_{n+1}(x) = \log_2 \{1 + F_n(x)\}$

$\therefore F_0(x) = x$ ;  $\therefore F_1(x) = \log_2(1+x)$ ;  $\therefore F_2(x) = \log_2 \{1 + \log_2(1+x)\}$

Cor.  $F_0(x) = x$  and  $f_0(x) = x$ .

2.  $\frac{d F_n(x)}{dx} \div \frac{d F_{n-1}(x)}{dx} = 1 + F_n(x)$ .

Sol.  $F_n(x) = \log_2 \{1 + F_{n-1}(x)\}$ ; differentiating both sides with respect to  $x$  we have  $\frac{d F_n(x)}{dx} = \{1 + F_{n-1}(x)\} \frac{d F_{n-1}(x)}{dx}$

Cor. 1.  $\frac{d F_n(x)}{dx} = \{1 + F_{n-1}(x)\} \{1 + F_{n-2}(x)\} \dots \{1 + F_0(x)\}$

Sol.  $F_n'(x) = \{1 + F_{n-1}(x)\} F_{n-1}'(x) = \{1 + F_{n-1}(x)\} \{1 + F_{n-2}(x)\} F_{n-2}'(x) = \dots = \{1 + F_0(x)\} F_0'(x) = \{1 + F_0(x)\} F_0'(x) = \dots =$



$$= \{1 + F_1(x)\} \{1 + F_2(x)\} \dots \{1 + F_n(x)\} + F'_0(x)$$

$$B_1 x - F_1(x) = x$$

$$n \phi_1(x) \phi_2(x) \dots \phi_{n-1}(x) + (n-1) \phi_1(x) \phi_2(x) \dots \phi_{n-2}(x) + \dots + (x-1) \phi_1(x) \phi_2(x) \dots \phi_{n-1}(x)$$

Here equate the coeff<sup>s</sup> of  $x^n$

$$3. \frac{df(x)}{dx} = x - \frac{1}{2} f_1(x) + B_2 f_2(x) - B_4 f_4(x) + B_6 f_6(x) - \dots$$

$$\text{Sol. } F'_n(x) = \{1 + F_1(x)\} \{1 + F_2(x)\} \dots \{1 + F_n(x)\}$$

$$\therefore 1 + n \frac{df(x)}{dx} + \dots = e^{F_0(x) + F_1(x) + \dots + F_n(x)}$$

$$\therefore \log \left\{ 1 + n \frac{df(x)}{dx} + \dots \right\} = F_0(x) + F_1(x) + \dots + n \text{ terms}$$

$$= \psi(x) + \int F_0(x) dx = \frac{1}{2} F_2(x) + \frac{B_2}{13} \frac{dF_2(x)}{dx} + \frac{B_4}{13} \frac{d^2 F_2(x)}{dx^2} + \dots$$

where  $\psi(x)$  is a function of  $x$  independent of  $x$ .

Equating the coeff<sup>s</sup> of  $x$  we get the result:

$$\text{Cor. } \psi(x) = \int_0^x \frac{x - \frac{df(x)}{dx}}{f_1(x)} dx$$

$$\text{Sol. since when } n=0, \log \left\{ 1 + n \frac{df(x)}{dx} + \dots \right\} = 0$$

$$\psi(x) = \frac{x}{2} - \frac{B_2}{2} f_1(x) + \frac{B_4}{6} f_1(x) - \log f_1(x) + \dots$$

$$\psi(x) = \frac{x}{2} - \frac{B_2}{2} f_1(x) + \frac{B_4}{6} f_1(x) - \log f_1(x) + \dots$$

$$\psi(x) f_1(x) = \frac{x}{2} f_1(x) - \frac{B_2}{2} f_1(x) f_1(x) + \frac{B_4}{6} f_1(x) f_1(x) - \dots$$

$$= \frac{x}{2} f_1(x) - \frac{B_2}{2} f_1(x) + \frac{B_4}{6} f_1(x) - \dots \text{ by II. 6.}$$

$$= x - f_1(x) \therefore \psi'(x) = \frac{x - f_1(x)}{f_1(x)}$$

$$6. \frac{d(F_n(x))}{dn} = f_1(x) \frac{dF_n(x)}{dx} \text{ and hence } n f_n'(x) = f_1(x) \frac{d f_n(x)}{dx}$$

Sol. In III write  $F_n(x) = f_n(x)$ ; then  $F_n(x) = F_n \{ F_n(x) \}$ .

$$\text{But } F_n(x) = F_n(x) + n \frac{dF_n(x)}{dx} + \frac{n^2}{2} \frac{d^2 F_n(x)}{dx^2} + \dots$$

$$\text{and } F_n \{ F_n(x) \} = F_n(x) + n f_1 \{ F_n(x) \} + n^2 f_2 \{ F_n(x) \} + \dots$$

Equating the coeff<sup>s</sup> of  $n$  we have  $\frac{dF_n(x)}{dx} = f_1 \{ F_n(x) \}$

Let  $F_n(1) = y$  and  $F_n'(1) = z$ , then we have

$$\frac{dy}{dx} = f_1(y) \quad \text{or} \quad \frac{dz}{dx} = f_1'(y) \frac{dz}{dy}$$

$$\frac{dF_n(x)}{dx} = f_1(x) \frac{dF_n(x)}{dy} \quad \text{Equating the coeff<sup>s</sup> of}$$

$n$  we have  $n f_n'(x) = f_1(x) f_{n-1}'(x)$ .

Cor 1. If  $f_n(x) = \left(\frac{x}{2}\right)^n \{ \psi_1^{(n)} x - \psi_2^{(n)} x^2 + \psi_3^{(n)} x^3 - \dots \}$

$$i. \quad n \psi_n^{(n)} = n \psi_1^{(n-1)} \psi_n^{(1)} + (n+1) \psi_2^{(n-1)} \psi_{n-1}^{(1)} + (n+2) \psi_3^{(n-1)} \psi_{n-2}^{(1)} + \dots$$

$$ii. \quad \phi_n(x) = n \pi \left\{ \frac{\psi_1^{(n-1)}}{n} - \frac{\psi_2^{(n-1)}}{n^2} + \frac{\psi_3^{(n-1)}}{n^3} - \dots \right\} x$$

Sol.  $n f_n'(x) = f_n(x) f_n'(x)$ ; here equate the coeff<sup>s</sup> of like powers of  $x$ .  $f_n'(x)$  is the coeff<sup>s</sup> of  $x^n$  in  $F_n(x)$  by I expansion. Again find the coeff<sup>s</sup> of  $x^n$  by II expansion and equate the two results.

$$\text{Cor 2. } (n+1) \psi_n^{(1)} = \frac{1}{2} \psi_{n-1}^{(1)} + \frac{B_2}{2} \psi_{n-2}^{(2)} - \frac{B_4}{24} \psi_{n-4}^{(4)} + \frac{B_6}{25} \psi_{n-6}^{(6)} - \frac{B_8}{27} \psi_{n-8}^{(8)} + \dots$$

Sol. Equate the coeff<sup>ts</sup> of  $x^n$  in IV 3.

$$5. f_1(x) = (1+x) f_2 \{ \log_2(1+x) \}$$

Sol. In IV 1. write  $\log_2(1+x)$  for  $x$ ; then  $F_{n-1}(x) =$

$$\log_2(1+x) + a f_1 \{ \log_2(1+x) \} + a^2 f_2 \{ \log_2(1+x) \} + \dots$$

$$\therefore e^{F_n(x)} = (1+x) e^{a f_1 \{ \log_2(1+x) \} + a^2 f_2 \{ \log_2(1+x) \} + \dots}$$

$$\text{But } e^{F_n(x)} = 1 + F_n(x) = 1 + x + a f_1(x) + a^2 f_2(x) + \dots$$

Equating the coeff<sup>ts</sup> of  $x$ :  $1 + x + a f_1(x) + a^2 f_2(x) + \dots = (1+x) f_2 \{ \log_2(1+x) \}$

6. i. The sum of the coeff<sup>ts</sup> in  $\phi_n(n)$  without the signs is  $\frac{1}{n}$  and with signs =  $\frac{1}{2n}$ .

Sol.  $F_1(x) = e^x - 1$  and  $F_2(x) = \log_2(1+x)$ ; equate the coeff<sup>ts</sup>.

$$\text{ii. } \psi_1(n) = 1; \quad \psi_2(n-1) = \frac{n}{8} \left( \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{n} \right);$$

$$\psi_3(n-2) = \frac{n(n-1)}{72} \left\{ \left( \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} \right)^2 - \left( \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \right) - \frac{1}{n} + \frac{1}{2} \right\}$$

$$\phi_1(2n) = 1$$

$$\phi_2(2n) = n$$

$$\phi_3(2n) = n^2 - \frac{n}{8}$$

$$\phi_4(2n) = n^3 - \frac{5n^2}{12} + \frac{n}{24}$$

$$\phi_5(2n) = n^4 - \frac{13}{18} n^3 + \frac{n^2}{6} - \frac{n}{90}$$

$$\phi_6(2n) = n^5 - \frac{77}{72} n^4 + \frac{89}{216} n^3 - \frac{91}{1440} n^2 + \frac{11n}{4320}$$

$$\phi_7(2n) = n^6 - \frac{32}{20} n^5 + \frac{175}{216} n^4 - \frac{149}{720} n^3 + \frac{9(n^2)}{4320} - \frac{n}{2360}$$

$$\phi_8(2n) = n^8; \quad \phi_1(n) = n; \quad \phi_2(n) = n(n - \frac{1}{2}); \quad \phi_3(n) = n(n - \frac{1}{2})(n - \frac{1}{4})$$

$$f_1(x) = \frac{x^2}{2} - \frac{x^3}{12} + \frac{x^4}{48} - \frac{x^5}{180} + \frac{11x^6}{8640} - \frac{x^7}{6720}$$

If  $\frac{x}{1-x} = y$  and  $1-x = z$ , then

$$F_{2n}(x) = y + \frac{y^2}{6} \log_e z + \frac{y^3}{72} \left\{ (\log_e z)^2 + (1 - \log_e z)^2 - z \right\} + \dots$$

Sol. Apply IV 6 ii in IV 1.

Ex. 1.  $f_1(x) f_1''(x) = f_1'(x) - f_2(x) + 3B_2 f_3'(x) - 5B_4 f_5'(x) + \dots$

Sol. From IV 3 we have  $f_1'(x) = x^{-1/2} f_1(x) + B_2 f_3(x) - B_4 f_5(x) + \dots$  differentiate both sides and multiplying the result by  $f_1(x)$  we have

$$f_1(x) f_1''(x) = f_1'(x) - \frac{1}{2} f_1(x) f_1'(x) + B_2 f_1(x) f_3'(x) - B_4 f_1(x) f_5'(x) + \dots$$

$$= f_1'(x) - f_1(x) + 3B_2 f_3(x) - 5B_4 f_5(x) + \dots \text{ by IV 4.}$$

2.  $\frac{1}{2} F_{2n}\left(\frac{1}{2}\right) = \frac{1}{2n} - \frac{\log_e 2}{2n^2} + \frac{(1/2 + \log_e 2)^2}{9n^3} - \dots$

Sol. Put  $x = \frac{1}{2}$  in IV 7.

8. i.  $\leq \frac{1}{1} + \leq \frac{1}{2} + \leq \frac{1}{3} + \dots + \leq \frac{1}{x} = (x+1) \leq \frac{1}{x} - x.$

ii.  $(\leq 4)^2 + (\leq \frac{1}{2})^2 + (\leq \frac{1}{3})^2 + \dots + (\leq \frac{1}{x})^2 = (x+1)(\leq \frac{1}{x})^2 - (2x+1)(\leq \frac{1}{x}) + \dots$

iii.  $(\leq \frac{1}{1})^3 + (\leq \frac{1}{2})^3 + (\leq \frac{1}{3})^3 + \dots + (\leq \frac{1}{x})^3 = (x+1)(\leq \frac{1}{x})^3 - 3(x+\frac{1}{2})(\leq \frac{1}{x})^2 + 3(2x+1)(\leq \frac{1}{x}) - 6x + \frac{1}{2}(\frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{x}).$

8. If two functions of  $x$  be equal, then a general theorem can be formed by simply writing  $\phi(x)$  instead of  $x^n$  in the original theorem.

Sol. Put  $x=1$  and multiply it by  $f(x)$ , then change  $x$  to  $x, x^2, x^3, x^4$  etc and multiply  $\frac{f(x)}{1}, \frac{f(x)}{x}, \frac{f(x)}{x^2}, \dots$  etc respectively and add up all the results. Then instead of  $x^n$  we have  $f(x^n)$  for positive  $n$ .

well as negative values, of  $n$ . Changing  $f(x)$  to  $\phi(x)$  we can get the result.

Ex. G. 1 We know that  $\tan^{-1} x + \tan^{-1} \frac{1}{x} = \frac{\pi}{2}$ . The theorem states a general theorem from this identity can be formed as follows:—

$$\frac{\phi(1) + \phi(-1)}{1} - \frac{\phi(3) + \phi(-3)}{3} + \frac{\phi(5) + \phi(-5)}{5} - \dots = \frac{\pi}{2} \phi'(0)$$

$$\text{Sol. } f(0) (\tan^{-1} 1 + \tan^{-1} 1) = \frac{\pi}{2} f(0)$$

$$\frac{f'(0)}{1} (\tan^{-1} x + \tan^{-1} \frac{1}{x}) = \frac{\pi}{2} \cdot \frac{f'(0)}{1}$$

$$\frac{f''(0)}{1^2} (\tan^{-1} x^2 + \tan^{-1} \frac{1}{x^2}) = \frac{\pi}{2} \cdot \frac{f''(0)}{1^2}$$

$$\dots \dots \dots \dots \dots \dots$$

Adding up all the results we have

$$\frac{f(1) + f(\frac{1}{2})}{1} - \frac{f(3) + f(\frac{1}{3})}{3} + \frac{f(5) + f(\frac{1}{5})}{5} - \dots$$

$$= \frac{\pi}{2} f'(0) \dots \dots \dots = \phi'(0), \text{ then } f(1) = \phi(1)$$

$$\frac{f(1) + f(\frac{1}{2})}{1} - \frac{f(3) + f(\frac{1}{3})}{3} + \dots = \frac{\pi}{2} \phi'(0)$$

2. Similarly we can derive from  $\frac{x}{1+x} + \frac{1}{x+1} = 1$   
 $\{ \phi(1) + \phi(-1) \} - \{ \phi(2) + \phi(-2) \} + \{ \phi(3) + \phi(-3) \} - \dots = \phi'(0)$

N. B. 1. If  $\phi(x)$  be substituted for  $x^n$ ,  $\phi'(0)$  must be substituted for  $\log x$ ,  $\phi''(0)$  for  $(\log x)^2$  &c.

N. B. 2. If an infinite number of terms vanish it may assume the form  $0 \times \infty$  and have a definite value. This error in case of a function of  $x$  is a function of  $e^{-x}$  which rapidly decreases

as  $x$  increases

$$3. \frac{\phi(0) - \phi(1)}{1} - \frac{\phi(1) - \phi(2)}{2} + \frac{\phi(2) - \phi(3)}{3} - \dots = \phi'(0).$$

Sol.  $\log_2(1+x) = \log_2(1+\frac{x}{2}) = \log_2 x$  Apply IV 9.

$$4. \frac{\phi(1)}{1!} - \frac{\phi(2)}{2!} + \frac{\phi(3)}{3!} - \dots = c\phi(0) + \phi'(0) \text{ nearly}$$

where  $c$  is the constant of  $\approx \frac{1}{2}$

Sol. change  $\phi(x)$  to  $\phi(\frac{x}{2})$  in the above result.  $\phi(1), \phi(2), \dots$  vanish.

$$\text{Cor. 1. } \frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} - \dots = c + \log_e x \text{ nearly.}$$

Here the error lies between  $\frac{e^{-x}}{x}$  &  $\frac{e^{-x}}{1+x}$ .

2. If  $x$  becomes greater and greater

$$\left(\frac{x}{1!}\right)^n - \frac{1}{2} \cdot \left(\frac{x^2}{2!}\right)^n + \frac{1}{3} \cdot \left(\frac{x^3}{3!}\right)^n - \dots = n \left(\frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} - \dots\right)$$

$$10. \phi(0) + \frac{x}{1!} \phi(1) + \frac{x(x-1)}{2!} \phi(2) + \dots$$

$$= \phi(x) + \frac{x}{1!} \phi(x-1) + \frac{x(x-1)}{2!} \phi(x-2) + \dots$$

$$\text{Sol. } 1 + \frac{x}{1!} + \frac{x(x-1)}{2!} x^2 + \dots = x^n + \frac{x}{1!} x^{n-1} + \dots$$

Apply IV 9.

Cor. If  $x=0$ , the value of the generating function of the series  $x^n \phi(0) + \frac{x}{1!} x^{n-1} \phi(1) + \frac{x(x-1)}{2!} x^{n-2} \phi(2) + \dots = \phi(x)$ .

Ex. 1. when  $x=0 = 0$

$$\frac{\phi(0)}{x} - \frac{\phi(1)}{x^2} + \frac{\phi(2)}{x^3} - \dots = \phi(0)$$

Ex. 2. Let  $\phi(x) = \frac{1}{2} \sin \frac{\pi x}{2}$ , then  $\phi(x) = \frac{\pi}{2}$

∴ When  $x=0$   $\frac{1}{x} - \frac{1}{2x^2} + \frac{1}{3x^3} - \dots = \frac{\pi}{2}$  which is same as saying  $\tan^{-1} \infty = \frac{\pi}{2}$ .

2. If  $x=0$ , then  $\frac{1}{x} - \frac{16}{2x^2} + \frac{16}{x^3} - \dots = \infty$   
Here L.S. =  $\frac{1}{x+1} - \frac{12}{x+3} - \frac{12}{x+5} - \dots$  when  $x=0$   
 $= \frac{1}{1} - \frac{12}{3} - \frac{12}{5} - \dots$

$= 1 + \frac{1}{2} + \frac{1}{4} + \dots = \infty$ .

∴ L.S. =  $\frac{x^2}{1} + \frac{x^2}{16} (1 + \frac{1}{2}) + \frac{x^2}{12} (1 + \frac{1}{2} + \frac{1}{3}) + \dots - e^{x(1+\frac{1}{2}+\dots)}$   
 $\rightarrow \infty$  when  $x=0$ .

3. If  $x=0$ , then  $x^x + \frac{x}{11} x^{-1} + x(1-0)2^{-1} + \dots = \ln x$  for all values of  $x$

to If  $x=0$ , show that

$\frac{1}{x} - \frac{16}{2x^2} + \frac{16}{x^3} - \frac{16}{2x^4} + \dots = \frac{\pi}{2}$ .

Sol. Write  $\sum_{n=1}^{\infty} (-1)^{n+1} \sin \frac{\pi x}{2}$  in ex. 1. Then  $f(x) = \sum_{n=1}^{\infty} \frac{\sin \pi x}{2^n}$ , when  $x=0$ ,  $\frac{\pi}{2}$ .

N.B. Thus we are able to find exact values when  $x=0$ , though the generating function may be too difficult to find. The generating function in ex. 1.

$= \frac{\pi}{2} \cos x + (c + \log 2) \sin x - \left\{ \frac{x}{11} - \frac{x^3}{16} (1 + \frac{1}{2} + \frac{1}{3}) + \frac{x^5}{12} (1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}) - \dots \right\}$   
 $= \frac{\pi}{2}$  when  $x=0$

11.  $\int_0^{\infty} e^{-x} x^n dx = \Gamma(n+1)$  and hence

$$\int_0^{\infty} x^{n-1} \{ \phi(0) - \frac{x}{\Gamma} \phi(1) + \frac{x^2}{\Gamma^2} \phi(2) - \dots \} dx = \Gamma(n) \phi(-n)$$

soln  $\int_0^{\infty} e^{-x} x^n dx = e^{-x} \{ -x^n + n x^{n-1} + n(n-1) x^{n-2} + \dots \}$   
when  $x = \infty = 0$  by IBP etc.

$$10) \int_0^{\infty} e^{-x} x^{-1} dx = \Gamma(0)$$

$$\frac{f'(0)}{\Gamma} \int_0^{\infty} e^{-x} x^{-2} dx = \frac{\Gamma(-1)}{\Gamma^2} f'(0)$$

$$\frac{f''(0)}{\Gamma^2} \int_0^{\infty} e^{-x} x^{-3} dx = \frac{\Gamma(-2)}{\Gamma^3} \frac{f''(0)}{\Gamma}$$

and so on.

Adding up all the results we have

$$\int_0^{\infty} x^{n-1} \{ f(x) - \frac{x}{\Gamma} f(1) + \frac{x^2}{\Gamma^2} f(2) - \dots \} dx = \Gamma(n) f(-n)$$

Let  $f(x) = \phi(x)$  then  $f(-n) = \phi(-n)$ .

Cor 1.  $\int_0^{\infty} x^{n-1} \{ \phi(0) - x \phi(1) + x^2 \phi(2) - \dots \} dx = \frac{\Gamma(n) \phi(-n)}{\sin \pi n}$

Cor 2.  $\int_0^{\infty} x^{n-1} \{ \phi(0) - \frac{x^2}{\Gamma} \phi(2) + \frac{x^4}{\Gamma^2} \phi(4) - \dots \} dx = \frac{\Gamma(n) \phi(-n)}{\Gamma \cos \frac{\pi n}{2}}$

Cor 3.  $\int_0^{\infty} \{ \phi(0) - \frac{x}{\Gamma} \phi(1) + \frac{x^2}{\Gamma^2} \phi(2) - \dots \} \cos \pi x dx = \phi(-1) - \pi^2 \phi(-3) + \pi^4 \phi(-5) - \dots$

Cor 4.  $\int_0^{\infty} \{ \phi(0) - x^2 \phi(2) + x^4 \phi(4) - \dots \} \cos \pi x dx = \frac{\pi}{2} \{ \phi(-1) - \frac{\pi}{\Gamma} \phi(-2) + \frac{\pi^2}{\Gamma^2} \phi(-3) - \frac{\pi^3}{\Gamma^3} \phi(-4) + \dots \}$



49. ~~12~~  
 Ex 5.  $\int_0^1 x^m (1-x)^n dx = \frac{\Gamma(m+1) \Gamma(n+1)}{\Gamma(m+n+1)}$

*sol. change  $x$  to  $\frac{z}{1+z}$  and apply I.I.*

12. From IV II Cor 3 & 4 we see that

if  $\int_0^\infty \phi(x) \cos nx dx = \psi(n)$ , then

(i)  $\int_0^\infty \psi(x) \cos nx dx = \frac{\pi}{2} \phi(n)$ .

(ii)  $\int_0^\infty \psi^2(x) dx = \frac{\pi}{2} \int_0^\infty \phi^2(x) dx$ .

13. (i)  $\int_0^{\frac{\pi}{2}} \sin^m x \cos^n x dx = \frac{\Gamma\left(\frac{m+1}{2}\right) \Gamma\left(\frac{n+1}{2}\right)}{2 \Gamma\left(\frac{m+n+1}{2}\right)}$

(ii)  $\int_0^{\frac{\pi}{2}} \cos^m x \cos nx dx = \frac{\pi \Gamma^m}{2^{m+1} \Gamma\left(\frac{m+n}{2}\right) \Gamma\left(\frac{m-n}{2}\right)}$

14.  $(1 + \frac{x^6}{16})(1 + \frac{x^6}{2^4})(1 + \frac{x^6}{3^2})(1 + \frac{x^6}{4^2}) \dots$   
 $= \frac{\sinh 2\pi x - 2 \sinh \pi x \cos \pi x \sqrt{3}}{4\pi^3 x^3}$

*Sol. L.S = (1 + x^6/16)(1 + x^6/2^4)(1 + x^6/3^2)(1 + x^6/4^2) ...*  
 $\times (1 + \frac{x^6}{5^2}) \dots$   
 $\times (1 + \frac{x^6}{7^2}) \dots$

15.  $e^x \left( \frac{x}{1!} - \frac{x^2}{2!} + \frac{x^3}{3!} - \frac{x^4}{4!} + \dots \right)$   
 $= \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} \left(1 + \frac{1}{3}\right) + \frac{x^4}{4!} \left(1 + \frac{1}{3}\right) + \frac{x^5}{5!} \left(1 + \frac{1}{3} + \frac{1}{5}\right) + \dots$

*Sol. L.S =*  $e^x \int_0^1 \frac{1 - e^{-zx}}{2z} dz = \int_0^1 \frac{e^x - e^{x(1-2z)}}{2z} dz = R.S$

i. If  $f(x+h) - f(x) = h \phi'(x)$ , then  
 $f(x) = \phi(x) - \frac{h}{2} \phi'(x) + \frac{B_2}{2!} h^2 \phi''(x) - \frac{B_4}{4!} h^4 \phi^{IV}(x) + \dots$

ii. If  $f(x+h) + f(x) = h \phi'(x)$ , then  
 $f(x) = \frac{h}{2} \phi'(x) - (2^2-1) B_2 \frac{h^2}{2!} \phi''(x) + (2^4-1) B_4 \frac{h^4}{4!} \phi^{IV}(x) - \dots$

Sol. If we write  $e^x$  for  $\phi(x)$ , we see that the coeff.<sup>s</sup> in R.S. are the same as those in the expansion of  $\frac{h}{e^{h/2}}$  and  $\frac{h}{e^{h/2}}$  respectively.

2. If  $F_n(x) = \phi(x) - \frac{n-1}{n+1} \{ \phi(x+h) + \phi(x-h) \} + \frac{(n-1)(n-3)}{(n+1)(n+3)} \times \{ \phi(x+2h) + \phi(x-2h) \} - \frac{(n-1)(n-3)(n-5)}{(n+1)(n+3)(n+5)} \{ \phi(x+3h) + \phi(x-3h) \} + \dots$

then,

i. If  $f(x+h) - f(x-h) = 2h \phi'(x)$ , then

$$f(x) = F_1(x) + \frac{1}{3} F_3(x) + \frac{1}{5} F_5(x) + \frac{1}{7} F_7(x) + \dots$$

ii. If  $f(x+h) + f(x-h) = 2\phi(x)$ , then

$$f(x) = F_1(x) + \frac{1}{2} F_3(x) + \frac{1 \cdot 3}{2^2} F_5(x) + \frac{1 \cdot 3 \cdot 5}{2^3} F_7(x) + \dots$$

3. If  $f(x+h) + p f(x) = \phi(x)$ , then

$$f(x) = \frac{\phi(x) \psi_0(p)}{p+1} - \frac{h}{2} \frac{\phi'(x) \psi_1(p)}{(p+1)^2} + \frac{h^2}{2!} \frac{\phi''(x) \psi_2(p)}{(p+1)^3}$$

- &c. where  $\psi(p)$  can be found from the expansion

$$\frac{1}{e^{x+p}} = \frac{\psi_0(p)}{p+1} - \frac{x}{2} \frac{\psi_1(p)}{(p+1)^2} + \frac{x^2}{2!} \frac{\psi_2(p)}{(p+1)^3} - \dots$$

Sol. let  $\phi(x) = e^x$ , then  $\frac{e^x}{e^{h+p}} = f(x)$ .

$$4. 1^n - 2^n p + 3^n p^2 - 4^n p^3 + 5^n p^4 - \dots = \frac{\psi_n(p)}{(p+1)^{n+1}}$$

$$\text{Sol. } \frac{1}{e^x + p} = e^{-x} - p e^{-2x} + p^2 e^{-3x} - p^3 e^{-4x} + \dots$$

equate the coeff<sup>ts</sup> of  $x^n$ .

$$5. \psi_0(p) - \frac{\pi}{\Gamma} \frac{\psi_1(p)}{p+1} + \frac{\pi(n-1)}{\Gamma^2} \frac{\psi_2(p)}{(p+1)^2} - \dots + (-1)^n \frac{\psi_n(p)}{(p+1)^{n+1}}$$

$$= (-1)^{n+1} \frac{p \psi_n(p)}{(p+1)^n}$$

Sol. Multiply both sides in  $\nabla. 3.$  by  $e^x + p$ ; then the coeff<sup>ts</sup> of  $x^n = 0$ .

$$6. \text{ If } \psi_n(p) = F_1(n) - p F_2(n) + p^2 F_3(n) - p^3 F_4(n) + \dots$$

$$+ (-1)^{n+1} F_n(n) p^{n-1} \text{ then i. } F_{n-2}(n) = F_{n+1}(n),$$

$$\text{ii. } F_n(n-1) + n F_{n+1}(n-1) + \frac{n(n+1)}{\Gamma^2} F_{n+2}(n-1) + \dots$$

$$+ \frac{1}{\Gamma} \frac{n+2-1}{\Gamma} F_0(n-1) = n^n$$

Sol. equate the coeff<sup>ts</sup> of  $p^{n-1}$  in  $\nabla. 4.$

$$\text{iii. } F_n(n-1) = n^{n-1} - \frac{\pi}{\Gamma} (n-1)^{n-1} + \frac{n(n-1)}{\Gamma^2} (n-2)^{n-1} - \dots$$

to  $n+1$  terms.

Sol. multiply both sides in  $\nabla. 4$  by  $(p+1)^{n+1}$  and equate the coeff<sup>ts</sup> of  $p^{n-1}$ .

7.  $\psi_n(x-1)$  is the integral part of

$$\frac{x^{n+1}}{1-x} \left\{ \frac{\Gamma x}{(\log \frac{1}{1-x})^{n+1}} - \frac{B_{n+1}}{n+1} \sin \frac{\pi x}{2} \right\}$$

Sol.  $e^{-x} + e^{-2x} + e^{-3x} + \dots = \frac{1}{e^x} = \frac{1}{x} - \dots$

Differentiating  $n$  times we have

$$1^n e^{-x} + 2^n e^{-2x} + 3^n e^{-3x} + \dots = \frac{(-1)^n}{x^{n+1}} \pm \dots$$

Writing  $\log \frac{1}{1-x}$  for  $x$  we have

$$1^n (1-x) + 2^n (1-x)^2 + 3^n (1-x)^3 + \dots = \frac{(-1)^n}{(\log \frac{1}{1-x})^{n+1}} \pm \dots$$

Apply  $\Sigma$  4.

$$8. \psi_n(-1) = 1^n; \psi_n(1) = 2^{n+1} (2^{n+1} - 1) \frac{\beta_{n+1}}{n+1} \sin \frac{\pi n}{2} \cdot \psi_0(b) = 1$$

$$\psi_1(b) = 1$$

$$\psi_2(b) = 1 - b$$

$$\psi_3(b) = 1 - 4b + b^2$$

$$\psi_4(b) = 1 - 11b + 11b^2 - b^3$$

$$\psi_5(b) = 1 - 26b + 66b^2 - 26b^3 + b^4$$

$$\psi_6(b) = 1 - 57b + 302b^2 - 302b^3 + 57b^4 - b^5$$

$$\psi_7(b) = 1 - 120b + 1191b^2 - 2416b^3 + 1191b^4 - 120b^5 + b^6$$

Write under each term the sum of the product of its coeff<sup>t</sup> and the no. of terms from the left & the product of the coeff<sup>t</sup> of the preceding term and its no. of terms from above.

Cor. 1.  $f(x)$  is the term independent of  $n$  in

$$\frac{\phi(x) + \frac{1}{n} \phi'(x) + \frac{1}{n^2} \phi''(x) + \frac{1}{n^3} \phi'''(x) + \dots}{e^{xh} + b}$$

2. If  $n \neq a$ , then  $F_n(n-1)$  is the coeff<sup>t</sup> of  $\frac{x^{n-1}}{n!}$  in  $e^{x(n-m)} (e^x - 1)^a$

3.  $\psi_n(b)$  is divisible by  $1-b$  if  $n$  is even

$$4. \frac{p + \cos x}{1 + 2p \cos x + p^2} = \frac{\psi_0(p)}{p+1} - \frac{x^2}{2!} \cdot \frac{\psi_2(p)}{(p+1)^2} + \frac{x^4}{4!} \cdot \frac{\psi_4(p)}{(p+1)^4} - \dots$$

$$5. \frac{\sin x}{1 + 2p \cos x + p^2} = \frac{x}{1!} \cdot \frac{\psi_1(p)}{(p+1)^2} - \frac{x^3}{3!} \cdot \frac{\psi_3(p)}{(p+1)^4} + \frac{x^5}{5!} \cdot \frac{\psi_5(p)}{(p+1)^6} - \dots$$

$$6. \text{If } 1^n (S_2 - 1) - 2^n (S_3 - 1) + 3^n (S_4 - 1) - \dots = \cos n\pi$$

$$- \frac{B_{n+1}}{n+1} (2^{n+1} - 1) \sin \frac{\pi n}{2} + A_1 S_2 - A_2 S_3 + A_3 S_4 - \dots$$

where  $S_n = \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots$ , then

$$i. A_n + n A_{n-1} + \frac{n(n-1)}{2!} A_{n-2} + \dots + A_0 = n^n.$$

$$ii. A_n = n^n - n(n-1)^n + \frac{n(n-1)}{2!} (n-2)^n - \dots$$

$$iii. \frac{A_n}{n!} \text{ is the coefft. of } x^n \text{ in } (e^x - 1)^n.$$

$$iv. \psi_n(p-1) = A_n - p A_{n-1} + p^2 A_{n-2} - \dots \text{ to } n \text{ terms}$$

$$\text{Ex. 1. } \frac{15^5}{2} + \frac{2^5}{2^2} + \frac{3^5}{2^3} + \dots = 1082.$$

$$2. \frac{15^5}{3} + \frac{2^5}{3^2} + \frac{3^5}{3^3} + \dots = 68 \frac{1}{2}.$$

$$9. \frac{x}{e^x - 1} = 1 - \frac{x}{2} + B_2 \frac{x^2}{2!} - B_4 \frac{x^4}{4!} + B_6 \frac{x^6}{6!} - \dots$$

where  $B_n$  can be found from

$$10. \frac{x}{e^x + 1} = \frac{x}{2} - B_2 \frac{x^2}{2!} (2^2 - 1) + B_4 \frac{x^4}{4!} (2^4 - 1) - \dots$$

$$\text{Sol. } \frac{x}{e^x + 1} = \frac{x}{e^x - 1} - \frac{2x}{e^{2x} - 1}.$$

$$11. \log \frac{x}{e^x - 1} = -\frac{x}{2} - B_2 \frac{x^2}{2!} + B_4 \frac{x^4}{4!} - \dots$$

$$\text{Sol. } \log(e^x - 1) = \int \frac{e^x}{e^x - 1} dx.$$

47

$$12. \log \frac{x}{e^{2x}+1} = -\frac{x}{2} - B_2 \frac{x^3}{2 \cdot 2!} (2^2-1) + B_4 \frac{x^5}{4 \cdot 4!} (2^4-1) - \dots$$

Sol.  $\log(e^{2x}+1) = \log(e^{2x}-1) - \log(e^x-1)$

Ex. If  $P, Q, R, S$  etc be so small that  $\frac{1}{2}$  of the sum of their cubes may be neglected, then

1. If  $e^P + e^Q + e^R = 2 + e^{P+Q+R}$  then

$$\left(\frac{1}{P} + \frac{1}{Q} + \frac{1}{R}\right) + \frac{1}{12}(P+Q+R) = -\frac{1}{2}$$

2. If  $e^{P+Q+R+S} = \frac{e^P + e^Q + e^R + e^S - 2}{e^{-P} + e^{-Q} + e^{-R} + e^{-S} - 2}$ , then

$$\left(\frac{1}{P} + \frac{1}{Q} + \frac{1}{R} + \frac{1}{S}\right) + \frac{1}{12}(P+Q+R+S) = 0$$

3. If  $2e^{P+Q+R+S+T} =$

$$\frac{(e^P + e^Q + e^R + e^S + e^T - 2)^2 - (e^{2P} + e^{2Q} + e^{2R} + e^{2S} + e^{2T} - 2)}{e^{-P} + e^{-Q} + e^{-R} + e^{-S} + e^{-T} - 2}$$

then  $\left(\frac{1}{P} + \frac{1}{Q} + \frac{1}{R} + \frac{1}{S} + \frac{1}{T}\right) + \frac{1}{12}(P+Q+R+S+T) = \frac{1}{2}$

13.  $x \cot x = 1 - B_2 \frac{(2x)^2}{2!} - B_4 \frac{(2x)^4}{4!} - B_6 \frac{(2x)^6}{6!} - \dots$

Sol. Change  $x$  to  $x/2$  in  $\nabla 9$ .

14.  $x \operatorname{cosec} x = 1 + B_2 \frac{x^2(2^2-2)}{2!} + B_4 \frac{x^4(2^4-2)}{4!} + \dots$

Sol.  $\operatorname{cosec} x = \cot \frac{x}{2} - \cot x$

15.  $x \tan x = B_2 \frac{2^2-1}{2!} (2x)^2 + B_4 \frac{2^4-1}{4!} (2x)^4 + \dots$

Sol.  $\tan x = \cot x - 2 \cot 2x$

$$16. \log_e \frac{x}{\sin x} = B_2 \frac{(2x)^2}{2!} + B_4 \frac{(2x)^4}{4!} + B_6 \frac{(2x)^6}{6!} + \dots$$

$$\text{Sol. } \log \sin x = \int \cot x \, dx.$$

$$17. \log_e \sec x = B_2 \frac{2^2-1}{2!} (2x)^2 + B_4 \frac{2^4-1}{4!} (2x)^4 + \dots$$

$$\text{Sol. } \log \sec x = \int \tan x \, dx.$$

N.B. 1. From the nature of the coeff<sup>ts</sup>, we see that  $B_0 = -1$ .

$$2. \frac{B_n}{B_{n-2}} = \frac{n(n-1)}{4\pi^2} \text{ nearly if } n \text{ is great.}$$

Sol. Since  $\cot \pi$  is  $-\infty$ ,  $B_{n-2} \frac{(2\pi)^{n-2}}{(n-2)!} \div$

$$B_n \frac{(2\pi)^n}{n!} = 1 \text{ nearly if } n \text{ is great.}$$

Similarly we can prove that

$$3. \frac{B_n}{B_{n-4}} = \frac{1}{(2\pi)^4} \cdot \frac{1}{(n-4)!} \text{ nearly if } n \text{ is great.}$$

$$18. (2n+1) B_{2n} = 2 B_2 B_{2n-2} \frac{2n(2n-1)}{2!} + 2 B_4 B_{2n-4} +$$

$$\frac{2n(2n-1)(2n-3)}{4!} + \dots \text{ the last term being}$$

$$2 B_{n-1} B_{n+1} \frac{1 \cdot 2n}{(n-1)! (n+1)!} \text{ or } (B_n)^2 \frac{1 \cdot 2n}{(n!)^2} \text{ according as}$$

$n$  is odd or even.

$$\text{Sol. } \cot^2 x = -\left(1 + \frac{d \cot x}{dx}\right); \text{ equate the coeff<sup>ts</sup> of } x^{2n-2}.$$

49

$$B_0 = -1; B_2 = \frac{1}{6}; B_4 = \frac{1}{30}; B_6 = \frac{1}{42}; B_8 = \frac{1}{30}; B_{10} = \frac{5}{66}$$

$$B_{12} = \frac{691}{2730}; B_{14} = \frac{7}{6}; B_{16} = \frac{3617}{510}; B_{18} = \frac{43867}{798}; B_{20} = \frac{174611}{330}$$

$$B_{22} = \frac{854513}{138}; B_{24} = \frac{236364091}{2730}; B_{26} = \frac{8553103}{6}; B_{28} = \frac{23749461029}{870}; B_{30} = \frac{8615841276005}{14322}$$

$$B_{32} = \frac{7709321041217}{510}; B_{34} = \frac{2577687858367}{6}; B_{36} = \frac{26315271553053477373}{1919190}; B_{38} = \frac{2929993913841559}{6}; \&c B_{\infty} = \infty$$

19. If  $n$  be an even integer,

- i.  $B_n$  is a fraction &  $2(2^n - 1) B_n$  is an integer
- ii. The numerator of  $B_n$  in its lowest terms is divisible by the greatest odd measure of  $n$  prime to  $(2^n - 1)$  and the quotient is a prime number the denominator
- iii. The denominator of  $B_n$  is the continued product of prime numbers next to the factors of  $n$  including unity and the number itself.





0	2	5	5	7	11	13	17	19	23	29	31	37	41	43	47	53	59	61	67
0	71	73	79	83	89	97	1	3	7	9	13	27	31	37	39	49	51	57	63
1	67	73	79	81	91	93	97	99	11	23	27	29	33	39	41	51	57	63	69
2	71	77	81	83	93	7	11	13	17	31	37	47	49	53	59	67	73	79	83
3	89	97	1	9	19	21	31	33	39	43	49	57	61	63	67	79	87	91	99
5	3	9	21	23	41	47	57	63	69	71	77	87	93	99	1	7	13	17	19
6	31	41	43	47	53	59	61	73	77	83	91	1	9	19	27	33	39	43	51
7	57	61	69	73	87	97	9	11	21	23	27	29	39	53	57	59	63	77	81
8	83	87	7	11	19	29	37	41	47	53	67	71	77	83	91	97	9	13	19
10	21	31	33	39	49	51	61	63	69	87	91	93	97	3	9	17	23	29	51
11	53	63	71	81	87	93	1	13	17	23	29	31	37	49	59	77	79	83	89
12	91	97	1	3	7	19	21	27	61	67	73	81	99	9	23	27	29	33	39
14	47	51	53	59	71	81	83	87	89	93	99	11	23	31	43	49	53	57	67
15	71	79	83	77	1	7	9	13	19	21	27	37	57	63	67	69	93	97	99
17	9	21	23	35	41	47	53	59	77	83	87	89	1	11	23	31	47	61	67
18	71	73	77	79	89	1	7	13	31	33	49	51	73	79	87	93	97	99	3
20	11	17	27	29	39	53	63	69	81	83	87	89	99	11	13	29	31	37	41
21	43	53	61	79	3	7	13	21	37	39	43	51	67	69	73	81	87	93	97
23	9	11	33	39	41	47	57	57	71	77	81	83	89	93	97	11	17	23	37
24	41	47	59	67	73	77	3	21	31	37	43	49	57	57	79	91	93	9	17
26	21	33	47	57	59	63	71	77	83	87	89	93	97	7	11	13	19	29	31
27	41	49	53	67	77	89	93	97	1	3	19	33	37	43	51	57	61	77	87
28	97	3	9	17	27	39	53	57	63	69	71	79	1	7	19	23	37	41	49
30	61	67	79	83	89	9	19	21	37	53	67	69	81	87	91	3	9	17	21
32	29	57	53	57	59	71	99	1	7	13	19	23	29	31	43	47	49	61	71
33	73	89	91	7	13	33	49	57	61	63	67	69	91	99	11	17	27	49	33
35	39	41	47	57	57	71	81	83	93	7	13	17	23	31	37	43	59	71	73
36	77	91	97	1	9	19	27	33	39	61	67	69	79	93	97	3	21	23	33
38	47	51	53	63	77	81	89	7	11	17	19	23	29	31	43	47	67	97	1
50	3	7	13	19	21	27	49	51	57	73	79	91	93	99	11	27	49	33	39
61	53	57	59	77	1	11	17	19	29	31	41	43	53	59	69	71	73	83	89
72	97	27	37	39	49	67	63	73	91	97	9	21	23	41	47	51	57	63	71
44	83	93	7	13	17	19	23	47	49	61	67	83	91	97	3	21	27	37	43
26	77	81	87	83	73	79	97	3	21	23	29	33	51	59	73	87	89	93	99
68	1	13	17	31	37	71	77	89	3	9	17	31	33	37	43	51	57	67	69

22. If  $\sec x = E_1 + \frac{x^2}{12} E_3 + \frac{x^4}{16} E_5 + \frac{x^6}{16} E_7 + \dots$  and consequently  $\frac{e^x + e^{-x}}{2} = E_1 - \frac{x^2}{12} E_3 + \frac{x^4}{16} E_5 - \dots$ , then

$$\frac{B_{2n}}{2^n} 2^{2n} (2^{2n} - 1) = 2 E_1 E_{2n-1} + 2 E_3 E_{2n-3} \frac{(2n-1)(2n-3)}{12} + \dots$$

the last term being  $2 E_{n-1} E_{n+1} \frac{(n-2)}{(n-1)!}$

$(E_n)^2 \frac{B_{n-1}}{(n-1)!}$  according as  $n$  is even or odd.

$E_1 = 1; E_3 = 1; E_5 = 5; E_7 = 61; E_9 = 1385; E_{11} = 50521$

$E_{13} = 2702765; E_{15} = 199360981$  &c &c  $E_{\infty} = \infty$

Sol.  $\frac{d \tan x}{dx} = \sec^2 x$ ; equate the coeff. of  $x^{n-2}$

23. i.  $\frac{1}{1-x^2} + \frac{1}{2^2-x^2} + \frac{1}{3^2-x^2} + \dots = \frac{1}{2x^2} - \frac{\pi}{2x} \cot \frac{\pi x}{2}$

ii.  $\frac{1}{1-x^2} + \frac{1}{3^2-x^2} + \frac{1}{5^2-x^2} + \dots = \frac{\pi}{4x} \tan \frac{\pi x}{2}$

iii.  $\frac{1}{1-x^2} - \frac{1}{2^2-x^2} + \frac{1}{3^2-x^2} - \dots = \frac{\pi}{2x} \operatorname{cosec} \pi x - \frac{1}{2x^2}$

iv.  $\frac{1}{1-x^2} - \frac{3}{3^2-x^2} + \frac{5}{5^2-x^2} - \dots = \frac{\pi}{6} \sec \frac{\pi x}{2}$

Sol. change  $x$  to  $\pi x$  in II 10.

24. i.  $\frac{1}{1+x^2} + \frac{1}{2^2+x^2} + \frac{1}{3^2+x^2} + \dots = \frac{\pi}{2x} \cdot \frac{e^{\pi x} + e^{-\pi x}}{e^{\pi x} - e^{-\pi x}} - \frac{1}{2x^2}$

ii.  $\frac{1}{1+x^2} + \frac{1}{3^2+x^2} + \frac{1}{5^2+x^2} + \dots = \frac{\pi}{4x} \cdot \frac{e^{\pi x} - 1}{e^{\pi x} + 1}$

iii.  $\frac{1}{1+x^2} - \frac{1}{2^2+x^2} + \frac{1}{3^2+x^2} - \dots = \frac{1}{2x^2} - \frac{\pi}{2(e^{\pi/2} - e^{-\pi/2})}$

iv.  $\frac{1}{1+x^2} - \frac{3}{3^2+x^2} + \frac{5}{5^2+x^2} - \dots = \frac{\pi}{e^{\pi/2} + e^{-\pi/2}}$

Sol. change  $x$  to  $\pi x$  in V. 13

N.B. 1. If  $x$  be of the form  $4m+1$ ,  $E_n$  ends in 5 and is 1 if of the form  $4m+3$ .  $E_n - 1$  is always divisible by 4 if  $n$  be any positive integer.

2.  $\frac{E_n}{E_{n+2}} = \frac{4n(n+1)}{\pi^2}$  nearly if  $n$  is great.  
 Sol.  $\sec \frac{\pi}{2} = \infty$ ;  $\therefore \frac{(\frac{\pi}{2})^{n+1}}{\Gamma(n+1)} E_{n+2} \div \frac{(\frac{\pi}{2})^{n-1}}{\Gamma(n-1)} E_n = \frac{1}{2}$  nearly if  $n$  is great. Similarly we can prove that

3.  $\frac{E_{n+2k+1}}{E_{n+1}} = \left(\frac{2}{\pi}\right)^{2k} \frac{\Gamma(n+2k)}{\Gamma(n)}$  (if  $n$  is great.) nearly.

25. i.  $\frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots = \frac{(2\pi)^n}{2\Gamma(n)} B_n = S_n$   
 ii.  $\frac{1}{1^n} + \frac{1}{3^n} + \frac{1}{5^n} + \dots = \frac{(2^{n-1})\pi^n}{2\Gamma(n)} B_n = S_n$   
 iii.  $\frac{1}{1^n} - \frac{1}{2^n} + \frac{1}{3^n} - \dots = \frac{(2^{n-1})\pi^n}{2\Gamma(n)} B_n = S_n'$   
 iv.  $\frac{1}{1^n} - \frac{1}{3^n} + \frac{1}{5^n} - \frac{1}{7^n} + \dots = \frac{\pi^n}{2^{n+1}\Gamma(n)} E_n = S_n''$

N.B. From  $\S$  18 and 22 we know the values of  $B_n$  and  $E_n$  only for even and odd integers; but from 25 for all positive values of  $n$ . For the values of  $B_n$  &  $E_n$  if  $n$  be negative see chap. and to find  $\Gamma(x)$  for all values of  $n$  see chap.

Cor 1. If  $x$  be a positive quantity not less than one the values of  $B_n$  are known from  $\S$  25. i & ii.

E.G.  $B_1 = \infty$ ;  $B_{1\frac{1}{2}} = \frac{3}{4\pi\sqrt{2}} S_{1\frac{1}{2}}$ ;  $B_3 = \frac{3}{2\pi^3} S_3$  &c.  
 2. If  $x$  be not a negative quantity the values of  $B_n$  and  $E_n$  are known from 25. iii & iv.

E.G.  $B_0 = -1$ ;  $B_{\frac{1}{2}} = -(1 + \frac{1}{\sqrt{2}})(\frac{1}{\sqrt{1}} - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} - \dots)$   
 $E_0 = \infty$ ;  $E_{\frac{1}{2}} = 2\sqrt{2}(\frac{1}{\sqrt{1}} - \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{5}} - \dots)$   
 $E_2 = \frac{8}{\pi^2}(\frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots)$  &c &c

$$26. \frac{1}{(a+b)^n} + \frac{1}{(a+2b)^n} + \frac{1}{(a+3b)^n} + \dots = \frac{1}{b(n-1)a^{n-1}} - \frac{1}{2an} \\ + B_2 \frac{n}{1^2} \cdot \frac{1}{a^{n+1}} - B_4 \frac{n(n+1)(n+2)}{1^4} \cdot \frac{1^3}{a^{n+3}} + \dots$$

From this we can sum up the reciprocals of powers of all numbers in A.P. approximately.  
Sol. Let L.S. =  $\phi(a)$ , then  $\phi(a-b) - \phi(a) = \frac{1}{a^n}$  by  $\Delta$ .

$$N.B. S_n = \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots + \frac{1}{(n-1)^n} + \frac{1}{2n^n} \\ + \frac{1}{(n-1)^{n-1}} + B_2 \frac{n}{1^2} \cdot \frac{1}{n^{n+1}} - B_4 \frac{n(n+1)(n+2)}{1^4} \cdot \frac{1}{n^{n+3}} + \dots$$

$$S_2 = 1.6449320668$$

$$S_3 = 1.2020569031$$

$$S_4 = 1.0823232337$$

$$S_5 = 1.0369277551$$

$$S_6 = 1.0173430620$$

$$S_7 = 1.0083492774$$

$$S_8 = 1.0040773562$$

$$S_9 = 1.0020083928$$

$$S_{10} = 1.0009945781$$

$$\frac{1}{B_1} = 0; \frac{1}{B_2} = 6.$$

$$\frac{1}{B_3} = 17.19624.$$

$$\frac{1}{B_4} = 30; \frac{1}{B_5} = 39.34953$$

$$\frac{1}{B_6} = 42; \frac{1}{B_7} = 38.03538$$

$$\frac{1}{B_8} = 36; \frac{1}{B_9} = 20.98719$$

$$\frac{1}{B_{10}} = 13.2$$

Cor. 1.  $n S_{n+1} = 1$  if  $n=0$  and  $S_{n+1} - \frac{1}{n} = .577$  nearly  
sol. write  $n+1$  for  $n$  and  $1$  for  $a$  in the above theorem;  
then we have  $S_{n+1} - \frac{1}{n} = \frac{1}{2} + B_2 \frac{n+1}{1^2} - \dots$   
 $= \frac{1}{2} + \frac{1}{2} - \frac{1}{20} + \dots = .577$  nearly when  $n$  varies.

2.  $\pi n B_{n+1} = 1$  when  $n=0$ .

Sol.  $n S_{n+1} = \frac{(2\pi)^n}{1^{n+1}} \pi n B_{n+1} = 1$  when  $n=0$

i.e.  $\pi n B_{n+1} = 1$  when  $n$  approaches 0.

$$\frac{1}{1-a_1} \cdot \frac{1}{1-a_3} \cdot \frac{1}{1-a_5} \cdot \frac{1}{1-a_7} \cdot \frac{1}{1-a_{11}} \cdot \frac{1}{1-a_{13}} \cdot \frac{1}{1-a_{17}} \dots$$

where 2, 3, 5, 7 &c are prime numbers.

=  $1 + a_2 + a_3 + a_2 a_2 + a_5 + a_2 a_3 + a_7 + a_2 a_2 a_2 + \dots$   
 where the Suffixes are natural numbers resolved into prime numbers.

28.  $(1 - \frac{1}{2^n})(1 - \frac{1}{3^n})(1 - \frac{1}{5^n})(1 - \frac{1}{7^n}) \dots = \frac{1}{S_m}$

Sol. Write  $\frac{1}{p^n}$  for  $a_p$  in 27. Similarly writing  $x^p$  for  $a_p$  we can get,

29. 
$$\frac{1}{(1-x^2)(1-x^3)(1-x^5)(1-x^7)(1-x^{11})(1-x^{13})(1-x^{17}) \dots}$$

$$= 1 + \frac{x^2}{1-x} + \frac{x^2+3}{(1-x)(1-x^2)} + \frac{x^2+3+5}{(1-x)(1-x^2)(1-x^3)}$$

$$+ \frac{x^2+3+5+7}{(1-x)(1-x^2)(1-x^3)(1-x^4)} + \dots$$

Cor. 1.  $(1 + \frac{1}{2^n})(1 + \frac{1}{3^n})(1 + \frac{1}{5^n}) \dots = \frac{S_m}{S_{2m}}$

2.  $\frac{2^n+1}{2^n-1} \cdot \frac{3^n+1}{3^n-1} \cdot \frac{5^n+1}{5^n-1} \dots = \frac{(S_m)^2}{S_{2m}}$

3.  $\frac{1}{2^n} + \frac{1}{3^n} + \frac{1}{5^n} + \frac{1}{7^n} + \frac{1}{8^n} + \frac{1}{11^n} + \frac{1}{12^n} + \dots$

where 2, 3, 5, 7 &c are natural numbers

Containing an odd number of prime factors

=  $\frac{(S_m)^2 - S_{2m}}{2 S_m}$  where  $S_m = \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{5^n} + \dots$

Sol. Invert both sides in 2<sup>o</sup> and coal. and find the difference after applying 2<sup>7</sup>.

Ex. 1. i.  $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$

ii.  $\frac{1}{1^3} - \frac{1}{3^3} + \frac{1}{5^3} - \dots = \frac{\pi^3}{32}$

iii.  $\frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots = \frac{\pi^4}{96}$

2. If 2, 3, 5, 7 &c be prime no., s.

i.  $\frac{2^2+1}{2^2-1} \cdot \frac{3^2+1}{3^2-1} \cdot \frac{5^2+1}{5^2-1} \dots = \frac{5}{2}$

ii.  $(1 + \frac{1}{2^4})(1 + \frac{1}{3^4})(1 + \frac{1}{5^4}) \dots = \frac{105}{\pi^4}$

3. If 2, 3, 5, 7, 8 &c be natural numbers containing an odd no. of prime factors.

i.  $\frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \frac{1}{8^2} + \dots = \frac{\pi^2}{20}$

ii.  $\frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \frac{1}{8^4} + \dots = \frac{\pi^4}{1260}$

Cor 4.  $\frac{3^n}{3^n+1} \cdot \frac{5^n}{5^n-1} \cdot \frac{7^n}{7^n+1} \cdot \frac{11^n}{11^n+1} \dots \frac{P^n}{P^n - \sin \frac{\pi n}{2}} \dots$  ad inf.

where P is a prime number.

$= \frac{1}{1^n} - \frac{1}{3^n} + \frac{1}{5^n} - \frac{1}{7^n} + \dots$

Cor 5.  $\frac{\log 1}{1^n} + \frac{\log 2}{2^n} + \frac{\log 3}{3^n} + \dots = \frac{\log 2}{2^n-1} + \frac{\log 3}{3^n-1} + \dots$

$\frac{\log 5}{5^n} + \dots$  where 2, 3, 5, 7 are prime numbers.

Sol. Differentiate both sides in 2<sup>o</sup>.

Ex.  $\frac{1}{2} \sin \frac{4\pi}{2} + \frac{1}{3} \sin \frac{3\pi}{2} + \frac{1}{5} \sin \frac{5\pi}{2} + \frac{1}{7} \sin \frac{7\pi}{2} + \dots$   
is a Convergent Series, 2, 3, 5 being prime no.s.

30.  $(1+a_2)(1+a_3)(1+a_5)(1+a_7)(1+a_{11}) \dots$   
 $= 1 + a_2 + a_3 + a_5 + a_7 + a_{11} + a_2 a_3 + a_2 a_5 + a_2 a_7 + a_2 a_{11} + a_3 a_5 + a_3 a_7 + a_3 a_{11} + a_5 a_7 + a_5 a_{11} + a_7 a_{11} + \dots$   
where the Suffixes are natural no.s resolved into prime factors no two of which are alike.

Cor 1.  $\frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \frac{1}{5^n} + \frac{1}{6^n} + \dots = \frac{S_n}{S_{2n}}$

2.  $\frac{1}{2^n} + \frac{1}{3^n} + \frac{1}{5^n} + \frac{1}{7^n} + \frac{1}{11^n} + \frac{1}{13^n} + \frac{1}{17^n} + \frac{1}{19^n} + \frac{1}{23^n}$   
 $+ \frac{1}{29^n} + \frac{1}{31^n} + \frac{1}{37^n} + \dots = \frac{(S_n)^2 - S_{2n}}{2 S_n S_{2n}}$

where 2, 3, 5, 7 &c are natural no.s containing an odd no. of prime factors no two of which are alike

3.  $\frac{1}{4^n} + \frac{1}{8^n} + \frac{1}{9^n} + \frac{1}{12^n} + \dots = \frac{S_n (S_{2n} - 1)}{S_{2n}}$

where 4, 8, 9, 12 &c are Composite numbers containing at least two equal prime numbers

Cor. 1. The sum of the reciprocals of all prime numbers is infinite.

Sol. Putting  $n=1$  in VI 28, we have



$$\frac{2}{2-1} \cdot \frac{3}{3-1} \cdot \frac{5}{5-1} \cdot \frac{7}{7-1} \dots = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots$$

$$\therefore \log \frac{2}{2-1} + \log \frac{3}{3-1} + \log \frac{5}{5-1} + \dots = \log (1 + \frac{1}{2} + \frac{1}{3} + \dots)$$

i.e.  $\frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots$  + a finite quantity =  $\infty$

i.e. The sum of the reciprocals of all prime nos. =  $\infty$

2. If 2, 3, 5, 7, ... be primes, then when  $n$  vanishes

$(\log n + \frac{1}{2^{n+1}} + \frac{1}{3^{n+1}} + \frac{1}{5^{n+1}} + \dots)$  is finite.

Sol. Changing  $n$  to  $n+1$  in Ex 2.8, we have

$$\left(1 - \frac{1}{2^{n+1}}\right) \left(1 - \frac{1}{3^{n+1}}\right) \left(1 - \frac{1}{5^{n+1}}\right) \dots = S_{n+1}$$

$$\therefore \log \left(1 - \frac{1}{2^{n+1}}\right) + \log \left(1 - \frac{1}{3^{n+1}}\right) + \log \left(1 - \frac{1}{5^{n+1}}\right) + \dots = -\log S_{n+1}$$

=  $-\log n$  when  $n$  is very small.

$$\therefore \log n + \frac{1}{2^{n+1}} + \frac{1}{3^{n+1}} + \frac{1}{5^{n+1}} + \dots = -.312 \text{ nearly when } n \rightarrow \infty$$

3. If  $P_n$  be the  $n$ th prime number, then

$\frac{P_n}{n} - \log n$  is finite if  $n$  is infinite.

Sol. Let  $S_n$  be the sum of  $n$  prime numbers.

$$\text{Then } S_2 = 5; S_3 = 17; S_4 = 41; S_5 = 77; S_6 = 129$$

$$P_2 = 3; P_3 = 5; P_4 = 7; P_5 = 11; P_6 = 13; P_7 = 17; P_8 = 19; P_9 = 23; P_{10} = 29$$

$$\therefore \frac{P_n + n + 1}{S_{2n}} = 1 \text{ if } n \text{ is very great.}$$

$$\therefore \frac{P_n}{n} - \log n \text{ is finite if } n \rightarrow \infty$$

Let  $f(x) + f(2x) + f(3x) + f(4x) + \dots + f(\infty) = \phi(x)$ , then  
 $\phi(x) = c + \int f(x) dx + \frac{1}{2} f(x) + \frac{B_2}{2} f'(x) - \frac{B_4}{4} f'''(x) +$   
 $\frac{B_6}{6} f^{(5)}(x) - \frac{B_8}{8} f^{(7)}(x) + \dots$

Sol.  $\phi(x) - \phi(x-1) = f(x)$ ; apply V.

N. B. By giving any value to  $x$ ,  $c$  can be found.

R. S. is not a terminating series except in some special cases. Consequently no constant can be found in  $\frac{1}{2} f(x) + \frac{B_2}{2} f'(x) - \frac{B_4}{4} f'''(x) + \dots$  except

in those special cases. If R. S. be a terminating series, it must be some integral function of  $x$ . In this case there is no possibility of a constant

(according to the ordinary sense) in  $\phi(x)$ ; for  $\phi(1) = f(1) + \phi(0)$ ; But  $\phi(1) = f(1)$ .  $\therefore \phi(0)$  is always

whether  $\phi(x)$  is rational or irrational.  $\therefore$  When  $\phi(x)$  is a rational integral function of  $x$  it must be divisible and hence no constant but

0 can exist. The algebraic constant of a series is the constant obtained by completing

the remaining part in the above theorem. We can substitute this constant which is like the center of gravity of a body instead of its dis-

urgent infinite series.

E.G. The constant of the series  $1+1+1+\dots = -\frac{1}{2}$ ; for  
 the sum to  $x$  terms  $= x = c + \int 1 dx + \frac{1}{2}$ .  $\therefore c = -\frac{1}{2}$   
 We may also find the constant thus:-

$$c = 1+1+3+4+\dots$$

$$\therefore 4c = 4 + 8 + \dots$$

$$\therefore -3c = 1-2+3-4+\dots = \frac{1}{(1+1)^2} = \frac{1}{4}$$

$$\therefore c = -\frac{1}{12}$$

$$2. \phi(x) + \sum_{n=0}^{\infty} \frac{B_n}{L^n} f^{(n)}(x) \cos \frac{\pi x}{2} = 0$$

Sol. Let  $\frac{B_n}{L^n} \psi^{(n)}$  be the coeff. of  $f^{(n)}(x)$ , then we

$$\text{see } \psi(0) = 1, \psi(2) = -1, \psi(4) = 1, \psi(6) = -1, \dots$$

$$\psi(3) = 0, \psi(5) = 0, \psi(7) = 0; \frac{B_1}{L} \psi(1) = \frac{1}{2}; \text{ but } B_1 = 0$$

$\therefore \psi(1) = 0$ . Again by  $\nabla$  26 cor 2. we have

$$\pi(n-1)B_n = 1 \text{ when } n \neq 1 \quad \therefore \frac{B_n \psi^{(n)}}{L^n} = \frac{\pi(n-1)B_n}{L^n} \frac{\psi^{(n)}}{\pi(n-1)}$$

$$= \frac{1}{2} \text{ when } n=1, \text{ i.e. } \frac{\psi^{(n)}}{\pi(n-1)} = \frac{1}{2} \text{ when } n \neq 1.$$

$$\therefore \psi^{(n)} = -\cos \frac{\pi x}{2}$$

3. The sum to a negative number of terms is  
 the sum with the sign changed, calculated  
 backwards from the term previous to the  
 first to the given number of terms with the  
 positive sign instead of negative.

$$\text{Sol. } \phi(x) = f(1) + f(2) + \dots + f(x+n)$$

$$- f(1+n) - f(2+n) - \dots - f(x+n)$$

change  $x$  to  $-x$  and put  $n = x$ , then we have

$$\phi(-x) = \phi(0) - \{f(0) + f(1) + f(2) + \dots + f(-x+1)\} \quad \text{ii}$$

but  $\phi(0) = 0$ .

E.G.  $1 + 2 + 3 + \dots$  to  $-5$  terms  
 $= -(0 - 1 - 2 - 3 - 4) = 10$

i. For finding the sum to a fractional number of terms assume the sum to be true always and if there is any difficulty in finding  $\phi(x)$ , take  $n$  any integer you choose, find  $\phi(n+x)$  and then subtract  $\{f(0+x) + f(1+x) + \dots + f(n+x)\}$  from the result

$$\text{ii. } \phi(h) = \phi(n) - \{f(0+h) + f(1+h) + \dots + f(n+h)\} \\ + h f(n) + \frac{x^2 h}{2} f'(n) + \frac{x^3 h^2}{6} f''(n) + \dots \text{ where } n \rightarrow \infty$$

$\rightarrow$  any integer or infinity.

E.G. 1.  $1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}$

$$= (1 + \frac{1}{2} + \dots + \frac{1}{n}) - (1 + \frac{1}{2+h} + \frac{1}{2+h} + \dots + \frac{1}{n+h}) \text{ when } n \rightarrow \infty$$

$$= C_0 + \log n - (1 + \frac{1}{2+h} + \frac{1}{2+h} + \dots + \frac{1}{n+h}) \text{ when } n \rightarrow \infty$$

where  $C_0$  is the constant of  $\log n$

2.  $\lfloor h = \frac{x^h}{(1 + \frac{1}{2}) (1 + \frac{1}{2})} - (1 + \frac{1}{2})$  when  $n = \infty$ .

Sol.  $\lfloor h = \frac{1 \times 2 + h}{1 \times 2} \cdot \frac{1 \times 2 + h}{1 \times 2} = \frac{n^h (1 + \frac{1}{2}) (1 + \frac{1}{2}) - (1 + \frac{1}{2})}{(1 + \frac{1}{2}) (1 + \frac{1}{2})} \cdot \frac{1 \times 2 + h}{1 \times 2}$

$$\therefore \lfloor h = \frac{1 \times 2 + h}{(1 + \frac{1}{2}) (1 + \frac{1}{2})} - (1 + \frac{1}{2}) = \frac{1 \times 2 + h}{(1 + \frac{1}{2}) (1 + \frac{1}{2})} - (1 + \frac{1}{2})$$

$$\text{iii. } \phi(h) = x f(0) - x^{1+h} f(0+h) + x^2 f(2) - x^{2+h} f(2+h) + \dots$$

5. Def. A series is said to be corrected when its constant is subtracted from it.

The differential Coeff<sup>t</sup> of a series is a corrected series.

$$\text{i.e. } \frac{d \{ \phi(0) + \phi(1) + \dots + \phi(x) \}}{dx} = \phi'(0) + \phi'(1) + \dots + \phi'(x) - c'$$

where  $c'$  is the constant of  $\phi(0) + \phi(1) + \dots + \phi(x)$ .

Sol. In the diff<sup>t</sup> coeff<sup>t</sup> of  $\phi(0) + \phi(1) + \dots + \phi(x)$  there can't be any constant. Therefore it should be corrected.

N.B. If  $f(0) + f(1) + \dots + f(x)$  be a convergent series then its constant is the sum of the series.

$$\text{E.G. 1. } \frac{d \left( 1 + \frac{1}{x} + \dots + \frac{1}{x} \right)}{dx} = \frac{1}{(x+1)^2} + \frac{1}{(x+2)^2} + \frac{1}{(x+3)^2} + \dots$$

$$\text{Sol. } \frac{d \sum \frac{1}{x}}{dx} = -\frac{1}{x^2} - \frac{1}{x^2} - \dots - \frac{1}{x^2} - c$$

$$= \frac{1}{(x+1)^2} + \frac{1}{(x+2)^2} + \dots$$

2. If  $c_0$  be the constant of  $\sum \frac{1}{x}$ , then

$$\frac{d \sum \frac{1}{x}}{dx} = \sum \left( \frac{1}{x} - c_0 \right)$$

$$\text{Sol. } \frac{d \sum \frac{1}{x}}{dx} = \sum \frac{d \log \frac{1}{x}}{dx} = \sum \left( \frac{1}{x} - c_0 \right)$$

$$\int_1^x \frac{1}{x} dx = \log Lx + x C_0$$

$$\int_0^x (1^{13} + 2^{13} + \dots + x^{13}) dx = \frac{1}{14} (1^{14} + 2^{14} + \dots + x^{14}) - \frac{x}{14}$$

$$\int_0^x (1^{10} + 2^{10} + \dots + x^{10}) dx = \frac{1}{11} (1^{11} + 2^{11} + \dots + x^{11}) - \frac{x}{11}$$

$$\int_1^x (\sqrt{1} + \sqrt{2} + \dots + \sqrt{x}) dx = \frac{2}{3} (1\sqrt{1} + 2\sqrt{2} + \dots + x\sqrt{x}) - \frac{x}{4\pi} (\frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{x}})$$

6. If  $f^n(x)$  stands for the  $n$ th derivative of  $f(x)$  and  $C_n$  be the constant of  $\{f'(x) + f''(x) + \dots + f^n(x)\}$

$$\text{then } \phi(x) = -C_1 x - C_2 \frac{x^2}{2} - C_3 \frac{x^3}{3} - C_4 \frac{x^4}{4} - \dots$$

$$\text{Sol. } \phi(x) = \phi(0) + \frac{x}{1} \phi'(0) + \frac{x^2}{2} \phi''(0) + \dots$$

But from VI 5 we have  $\phi(0) = 0$ ,  $\phi'(0) = -C_1$ ,  $\phi''(0) = -C_2 \times 2$

E.g. 1.  $\log Lx = -S_1 x + \frac{S_2}{2} x^2 - \frac{S_3}{3} x^3 + \dots$  where  $S_n$  is the constant of  $(\frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots)$

2.  $\frac{1}{x} = S_1 x - S_3 x^3 + S_5 x^5 - \dots$  where  $S_n = \frac{1}{1^n} + \frac{1}{2^n} + \dots$

N.B. This is very useful in finding  $\phi(x)$  for fractional values of  $x$ .

7. If  $C_n'$  be the constant of

$$f(\frac{1}{n}) + f(\frac{2}{n}) + f(\frac{3}{n}) + \dots + f(\frac{x}{n}), \text{ then}$$

$$\phi\left(\frac{x}{n}\right) + \phi\left(\frac{x-1}{n}\right) + \phi\left(\frac{x-2}{n}\right) + \dots + \phi\left(\frac{x-n+1}{n}\right) - nc$$

$$= f\left(\frac{x}{n}\right) + f\left(\frac{x-1}{n}\right) + \dots + f\left(\frac{x}{n}\right) - c'_n$$

Sol. Let  $\psi(x) = \phi\left(\frac{x}{n}\right) + \phi\left(\frac{x-1}{n}\right) + \dots + \phi\left(\frac{x-n+1}{n}\right)$ , then

$$\psi(x) - \psi(x-1) = \phi\left(\frac{x}{n}\right) - \phi\left(\frac{x-1}{n}\right) = f\left(\frac{x}{n}\right)$$

$\therefore \psi(x)$  &  $f\left(\frac{x}{n}\right) + f\left(\frac{x-1}{n}\right) + \dots + f\left(\frac{x}{n}\right)$  differ only by some constant; hence if these be corrected they must be equal.  $\psi(x)$  contains  $n$  terms each each of which is of the form  $\phi(y)$  whose constant is  $c$ .  $\therefore$  The constant of  $\psi(x)$  is  $nc$  & the constant of  $f\left(\frac{x}{n}\right) + f\left(\frac{x-1}{n}\right) + \dots + f\left(\frac{x}{n}\right)$  is  $c'_n$  by our supposition.

Coroll.  $\phi\left(\frac{1}{n}\right) + \phi\left(\frac{2}{n}\right) + \dots + \phi\left(\frac{n-1}{n}\right) = nc - c'_n$

Sol. Put  $x = 0$  in the above theorem.

i.  $\phi\left(\frac{1}{2}\right) = 2c - c'_2$ .

ii.  $c = c_0 = c'_1$ .

iii.  $\phi\left(\frac{1}{3}\right) + \phi\left(\frac{2}{3}\right) = 3c - c'_3$

iv.  $\phi\left(\frac{1}{4}\right) + \phi\left(\frac{3}{4}\right) = 2c + c'_2 - c'_4$ .

v.  $\phi\left(\frac{1}{8}\right) + \phi\left(\frac{5}{8}\right) = c + c'_2 + c'_3 - c'_4$ .

8.  $\phi\left(x - \frac{1}{2}\right) = c + \int f(x) dx - (1 - \frac{1}{2}) \frac{B_{1/2}}{1/2} f'(x) + (1 - \frac{1}{2}) \frac{B_{3/2}}{3/2} f''(x)$   
 $- 2c = \sum_{n=0}^{\infty} \left\{ \left(1 - \frac{1}{2^{n+1}}\right) \frac{B_n}{n!} f^{(n)}(x) \cos \frac{\pi n}{2} \right\}$

Sol. Put  $n=2$ , change  $x$  to  $2x$  and apply VI 1.

- i.  $S(a_1 + a_2 + a_3 + \dots)$  means that the series is a convergent series and its sum to infinity is required
- ii.  $C(a_1 + a_2 + a_3 + \dots)$  means that the series is a divergent series and its constant is reqd.
- iii.  $G(a_1 + a_2 + a_3 + \dots)$  means that the series is oscillating or divergent and the value of its generating function is required.

N.B. Hereafter the series will only be given omitting  $S, C$  or  $G$  and from the nature of the series we should infer whether  $C, S$  or  $G$  is reqd; more over if a series appear to be equal to a finite quantity we must select  $S, C$  or  $G$  from the nature of the series.

- i. The value of an oscillating series is only true when the series is deduced from a regular series. For example the series  $1 - 1 + 1 - 1 + \dots = \frac{1}{2}$  only when it is deduced from a regular series of the form  $\phi(1) - \phi(2) + \phi(3) - \dots$ . Again if we take an irregular series  $a^x - b^x + c^x - d^x + \dots$  we get the same series  $1 - 1 + 1 - 1 + \dots$  when  $a$  becomes 0; yet its value is not  $\frac{1}{2}$  in this case.

ii.  $a_1 - a_2 + a_3 - a_4 + \dots$  is not equal to the series  $(a_1 - a_2) + (a_3 - a_4) + (a_5 - a_6) + \dots$  or to the series



$a_1 - (a_2 - a_3) - (a_4 - a_5) - (a_6 - a_7) - \dots$ ; but to the series  $a_1 - (a_2 - a_3 + a_4 - \dots)$

e.g.  $1 - 2 + 3 - 4 + \dots$  is not equal to  $(1 - 2) + (3 - 4) + (5 - 6) + \dots$  or to  $1 - (2 - 3) - (4 - 5) - \dots$

$$\text{iii. } (a_1 - a_2 + a_3 - \dots) \pm (b_1 - b_2 + b_3 - \dots)$$

$$= (a_1 \pm b_1) - (a_2 \pm b_2) + (a_3 \pm b_3) - \dots$$

Ex. 1. Show that  $(a_1 - a_2 + a_3 - \dots) + (b_1 - b_2 + \dots)$

$$= a_1 + (b_1 - a_2) - (b_2 - a_3) + (b_3 - a_4) - \dots$$

$$\text{Sol. L.S.} = a_1 + (b_1 - b_2 + b_3 - \dots) - (a_2 - a_3 + \dots)$$

$$= a_1 + (b_1 - a_2) - (b_2 - a_3) + \dots$$

$$2. a_1 - a_2 + a_3 - a_4 + \dots = \frac{a_1}{2} + \frac{1}{2} \{ (a_1 - a_2) - (a_2 - a_3) + \dots \}$$

$$3. = \frac{3a_1 - a_2}{4} + \frac{1}{4} \{ (a_1 - 2a_2 + a_3) - (a_2 - 2a_3 + a_4) + \dots \}$$

$$\frac{1}{4} = \frac{7a_1 - 4a_2 + a_3}{8} + \frac{1}{8} \{ (a_1 - 3a_2 + 3a_3 - a_4) - (a_2 - 3a_3 + 3a_4 - a_5) + (a_3 - 3a_4 + 3a_5 - a_6) - \dots \}$$

$$\text{ii. } a_1 - a_2 + a_3 - a_4 + \dots$$

$$= \frac{a_1}{2} + \frac{a_1 - a_2}{4} + \frac{a_1 - 2a_2 + a_3}{8} + \dots$$

$$= x a_1 - x^2 a_2 + x^3 a_3 - x^4 a_4 + \dots$$

$$= x \cdot \frac{a_1}{2} + x^2 \cdot \frac{a_1 - a_2}{4} + x^3 \cdot \frac{a_1 - 2a_2 + a_3}{8} + \dots$$

when  $x$  approaches unity.

11. If  $\frac{2}{3}$  lies between  $\frac{a_1}{a_2}$  &  $\frac{a_3}{a_4}$ , then

$a_2 + a_3 - a_4 + \dots$  lies between  $\frac{a_1^2}{a_1 + a_2}$  &  $a_1 - \frac{a_2^2}{a_2 + a_3}$

12.  $1 - 2 + 3 - 4 + \dots$  lies between  $\frac{1}{3}$  &  $\frac{1}{5}$  and its value is  $\frac{1}{4}$ .  
 $10 - 11 + 12 - 13 + \dots$  lies between  $\frac{1}{2}$  &  $\frac{1}{3}$ ; its value is  $\frac{3}{5}$  very nearly.

But  $2 - 2\frac{1}{2} + 3\frac{1}{3} - 4\frac{1}{4} + 5\frac{1}{5} - \dots$  cannot lie between  $\frac{2^2}{2+2\frac{1}{2}}$  &  $2 - \frac{(2\frac{1}{2})^2}{2\frac{1}{2}+3\frac{1}{3}}$  as  $2\frac{1}{2}$  is not lying between  $\frac{2}{2\frac{1}{2}}$  &  $\frac{3\frac{1}{3}}{2\frac{1}{2}}$ . i.e. it cannot lie between .889 & .929 as its value is .17.

13.  $\phi_1(x) + \phi_2(x) + \phi_3(x) + \dots$  can be expanded in ascending powers of  $x$ , say  $A_0 + A_1x + A_2x^2 + \dots$  where each of  $\phi_1, \phi_2, \dots$  is a series.

Case I when  $A_n$  is a convergent series

(1) If  $A_0 + A_1x + A_2x^2 + \dots$  be a rapidly convergent series what is required is got.

(2) But if it is a slowly convergent or an oscillating series, convergent, or divergent (at least for some values of  $x$ )

(3) Change  $x$  into a suitable function of  $y$  so that the new series in ascending powers

of  $y$  may be a rapidly convergent series;

e.g. let  $\frac{x}{1+x^2} = y$ , then  $x - \frac{x^3}{2} + \frac{x^5}{3} - \frac{x^7}{4} + \dots$   
 $= y + \frac{y^3}{12} + \frac{y^5}{80} + \frac{y^7}{448} + \dots$

(b) or convert it into a continued fraction

e.g.  $x + \frac{x^2}{3} + \frac{2}{15}x^3 - \frac{17}{315}x^5 + \dots = \frac{x}{1 + \frac{x}{3 + \frac{x}{5 + \dots}}}$   
 $\frac{1}{x} - \frac{11}{x^2} + \frac{11}{x^3} - \frac{13}{x^4} + \dots = \frac{1}{x+1} - \frac{12}{x+3} - \frac{2}{x+5} - \dots$

(c) or transform it into another series by applying III 8; e.g.  $\frac{1}{x} - \frac{2}{x^2} + \frac{5}{x^3} - \frac{16}{x^4} + \dots$

$$= \frac{1}{x+1} - \frac{1}{(x+1)(x+2)} + \frac{1}{(x+1)(x+2)(x+3)} - \dots$$

(d) or take the reciprocal of the series and try to make it a rapidly convergent series in any way

Case II When  $A_n$  is an oscillating (convergent or divergent) or a pure divergent series

(1) Let  $C_n$  be the constant or the value of the generating function. Then the given series  $= \psi(x) + C_0 + C_1x + C_2x^2 + C_3x^3 + \dots$  where  $\psi(x)$  can be found in special cases.

(2) But if  $C_0 + C_1x + C_2x^2 + \dots$  be a divergent series find some function of  $x$  (say  $P_n$ ) such that the value of  $P_0 + P_1x + P_2x^2 + \dots$  may be easily

found and  $c_n - p_n$  may rapidly diminish as  $n$  increases. Then the given series =

$$F(x) + (c_0 - p_0) + (c_1 - p_1)x + (c_2 - p_2)x^2 + \dots$$

e.g. 1.  $\frac{1}{x+1} - \frac{1}{x+2} + \frac{1}{x+3} - \dots = \frac{1}{x}(1-1+1-\dots)$

$$- \frac{1}{x^2}(1-2+3-\dots) = \frac{1}{2x} - \frac{1}{4x^2} + \dots$$

2.  $\frac{1}{1-x^2} + \frac{1}{2-x^2} + \frac{1}{3-x^2} + \dots = -\frac{1}{x^2}(1+1+1+\dots)$

$$- \frac{1}{x^6}(1^6+2^6+3^6+\dots) - \frac{1}{x^6}(1^4+2^4+3^4+\dots) = \psi(x)$$

$$+ \frac{1}{2x^2} = \frac{1}{2x^2} - \frac{\pi \cot \pi x}{2x}$$

3.  $\frac{1}{1^x} + \frac{1}{2^x} + \frac{1}{3^x} + \dots = (1+1+1+\dots)$

$$- x(\log 1 + \log 2 + \dots) + \dots = -\frac{1}{2} - x \log \sqrt{2\pi} - \dots$$

$$= \frac{1}{x-1} + 1+x+x^2+\dots - \frac{1}{2} - x \log \sqrt{2\pi} - \dots$$

$$= \frac{1}{x-1} + \frac{1}{2} + (1-.91894)x - \dots$$

$$= \frac{1}{x-1} + \frac{1}{2} + .08106x - \dots$$

4.  $\frac{x}{e^x+1} + \frac{x}{e^{2x}+1} + \frac{x}{e^{3x}+1} + \dots$

$$= \log 2 - \frac{x}{4} + (\beta_2)^2 \frac{x^2(2^2-1)}{2 \cdot 2^2} + (\beta_4)^2 \frac{x^4(2^4-1)}{4 \cdot 2^4} +$$

$$(\beta_6)^2 \frac{x^6(2^6-1)}{6 \cdot 2^6} + \dots$$

Sol.  $\frac{x}{e^x+1} + \frac{x}{e^{2x}+1} + \frac{x}{e^{3x}+1} + \frac{x}{e^{4x}+1} + \dots$

$$= \frac{x}{2}(1+1+1+\dots) - \beta_2 \frac{x^2(2^2-1)}{2^2}(1+2+3+\dots)$$

$$+ \beta_4 \frac{x^4(2^4-1)}{2^4}(1^4+2^4+3^4+\dots) - \dots$$

$$= \psi(x) - \frac{x}{4} + (B_2)^2 \frac{x^2(2^2-1)}{2!2^2} + (B_4)^2 \frac{x^4(2^4-1)}{4!2^4} + \dots$$

Now it is req<sup>d</sup> to find  $\psi(x)$ .

The given series =  $\frac{x}{e^x-1} - \frac{x}{e^{2x}-1} + \frac{x}{e^{3x}-1} - \dots$   
 $= \log 2 +$  terms involving  $x$  & higher powers of  $x$ .  
 $\therefore \psi(x) = \log 2$ .

ii.  $\frac{x}{e^x-1} + \frac{x}{e^{2x}-1} + \frac{x}{e^{3x}-1} + \frac{x}{e^{4x}-1} + \dots$   
 $= C - \log x + \frac{x}{4} - (B_2)^2 \frac{x^2}{2!2^2} - B_4^2 \frac{x^4}{4!2^4} - B_6^2 \frac{x^6}{6!2^6} - \dots$

Sol. Proceeding as in the previous theorem we have the series =  $\psi(x) + C + \frac{x}{4} - B_2^2 \frac{x^2}{2!2^2} - B_4^2 \frac{x^4}{4!2^4} - \dots$

But we know  $\frac{x}{e^x+1} + \frac{x}{e^{2x}+1} + \frac{x}{e^{3x}+1} + \dots$

$$= \left( \frac{x}{e^x-1} + \frac{x}{e^{2x}-1} + \dots \right) - \left( \frac{2x}{e^{2x}-1} + \frac{2x}{e^{4x}-1} + \dots \right)$$

$$\therefore \psi(x) - \psi(2x) = \log 2; \text{ hence } \psi(x) = -\log_2 x.$$

Ex. 1: Show that the constant in the series

$$\sqrt[100]{1} + \sqrt[100]{2} + \sqrt[100]{3} + \sqrt[100]{4} + \dots + \sqrt[100]{x}$$

$$\text{is } -1.4909100$$

2.  $\frac{1}{2+1} + \frac{1}{2^2+1} + \frac{1}{2^3+1} + \dots = \frac{3}{4} + \frac{\log 2}{48}$  nearly

3.  $\frac{1}{1+\frac{10}{9}} + \frac{1}{1+(\frac{10}{9})^2} + \frac{1}{1+(\frac{10}{9})^3} + \dots = 6.331009$ .

$$4. \frac{1}{\frac{10}{9}-1} + \frac{1}{\left(\frac{10}{9}\right)^2-1} + \frac{1}{\left(\frac{10}{9}\right)^3-1} + \dots = 27 \text{ nearly. } 71$$

$$15. \text{ i. } \frac{1}{x-1} + \frac{1}{x^2-1} + \frac{1}{x^3-1} + \dots$$

$$= \frac{1}{2} \cdot \frac{x+1}{x-1} + \frac{1}{x^2} \cdot \frac{x^2+1}{x^2-1} + \frac{1}{x^3} \cdot \frac{x^3+1}{x^3-1} + \dots$$

$$\text{ii. } \frac{1}{x-1} - \frac{1}{x^2-1} + \frac{1}{x^3-1} - \frac{1}{x^4-1} + \dots$$

$$= \frac{1}{x} \cdot \frac{x^2+1}{x^2-1} - \frac{1}{x^2} \cdot \frac{x^4+1}{x^4-1} + \frac{1}{x^3} \cdot \frac{x^6+1}{x^6-1} - \dots$$

$$\text{Sol. } \frac{1}{x-1} = \frac{1}{x-1}$$

$$\pm \frac{1}{x^2-1} = \pm \left\{ \frac{1}{x^2} + \frac{1}{x^2(x^2-1)} \right\}$$

$$\frac{1}{x^3-1} = \frac{1}{x^3} + \frac{1}{x^3} + \frac{1}{x^3(x^3-1)}$$

$$\pm \frac{1}{x^4-1} = \pm \left\{ \frac{1}{x^4} + \frac{1}{x^4} + \frac{1}{x^4(x^4-1)} \right\}$$

$\dots \dots \dots$

Adding up all the terms we can get the results.

$$16. \frac{a}{1-ax} + \frac{a^2}{1-ax^2} + \frac{a^3}{1-ax^3} + \dots \text{ to } n \text{ terms}$$

$$= \frac{axx}{1-ax} + \frac{(axx^2)^2}{1-ax^2} + \frac{(axx^3)^3}{1-ax^3} + \dots \text{ to } n \text{ terms}$$

$$+ \frac{a-a^{n+1}}{1-a} + a \frac{(ax)^2 - (ax)^{n+1}}{1-ax} + a^2 \frac{(ax^2)^3 - (ax^2)^{n+1}}{1-ax^2} + \dots$$

to  $n$  terms.

$$\text{Sol. } \frac{a}{1-ax} = \frac{axx}{1-ax} + a$$

$$\frac{a^2}{1-ax^2} = \frac{(axx^2)^2}{1-ax^2} + a^2 + a^2 x^2$$

$$\frac{a^3}{1-ax^3} = \frac{(arx^3)^3}{1-ax^3} + r^3 + ar^3x^3 + a^2r^3x^6$$

&c &c &c

Adding up all the terms in the n rows we can get the result.

$$\text{Coef. } \frac{a}{1-ax} + \frac{a^2}{1-ax^2} + \frac{a^3}{1-ax^3} + \dots$$

$$= \frac{arx}{1-ax} + \frac{(arx^2)^2}{1-ax^2} + \frac{(arx^3)^3}{1-ax^3} + \dots$$

$$+ \frac{a}{1-a} + \frac{a(arx)^2}{1-ax} + \frac{a^2(arx^2)^3}{1-ax^2} + \frac{a^3(arx^3)^4}{1-ax^3} + \dots$$

$$17. \frac{a}{1-m} + \frac{(a+b)n}{1-mx} + \frac{(a+2b)n^2}{1-mx^2} + \frac{(a+3b)n^3}{1-mx^3} + \dots$$

$$= a \cdot \frac{1-mn}{(1-m)(1-n)} + (a+b) \frac{1-mnx^2}{(1-mx)(1-nx)} (mnx)$$

$$+ (a+2b) \frac{1-mnx^4}{(1-mx^2)(1-nx^2)} (mnx^2)^2 + (a+3b) \frac{1-mnx^6}{(1-mx^3)(1-nx^3)}$$

$$+ \dots + \frac{b}{m} \left\{ \frac{mn}{(1-n)^2} + \frac{(mnx)^2}{(1-nx)^2} + \frac{(mnx^4)^3}{(1-nx^2)^2} + \dots \right\}$$

$$\text{Coef. } \frac{a}{1-m} + \frac{(a+b)n}{1-mx} + \frac{(a+2b)n^2}{1-mx^2} + \dots$$

$$= a \cdot \frac{1+m}{1-m} + (a+b) \frac{1+nx}{1-nx} \cdot (n^2x) + (a+2b) \frac{1+nx^2}{1-nx^2} (n^2x^2)^2$$

$$+ b \left\{ \frac{n}{(1-n)^2} + \frac{n^3x^2}{(1-nx)^2} + \frac{n^5x^6}{(1-nx^2)^2} + \frac{n^7x^{12}}{(1-nx^3)^2} + \dots \right\}$$

2. If  $A_n$  denotes the no. of factors in  $x$  including 1 &  $x$  then  $\frac{A_1}{x} + \frac{A_2}{x^2} + \frac{A_3}{x^3} + \dots = \frac{1}{2-1} + \frac{1}{2^2-1} + \dots$   
and hence deduce  $\sqrt{15}$

$$1. \quad 1^2 + 3^2 + 4^2 + 5^2 + \dots + x^2 = \phi_n(x)$$

$$\phi_n(x) = \frac{B_{n+1} \cos \frac{\pi(n+1)}{2}}{n+1} + \frac{x^{n+1}}{n+1} + \frac{x^n}{2} + B_2 \frac{n}{2} x^{n-1} - B_4 \frac{n(n-1)(n-2)}{24}$$

$$\times x^{n-3} + B_6 \frac{n(n-1)(n-2)(n-3)}{720} x^{n-5} - \dots$$

Sol. The corrected series is found by applying VI b.

$$\text{The coeff. of } x^{n-n} = - \frac{1/2}{n^n |n+1} \cdot B_{n+1} \cos \frac{\pi(n+1)}{2}$$

\(\therefore\) The values of the corrected series when  $x=0$

$$= - \frac{B_{n+1} \cos \frac{\pi(n+1)}{2}}{n+1} \text{ by IV 10 Cor. But } \phi_n(0) = 0$$

$$\therefore \text{The constant} = \frac{B_{n+1} \cos \frac{\pi(n+1)}{2}}{n+1}$$

A.B. If  $C_n$  be the constant, then  $C_{-n} = S_n$  and consequently  $S_{-n}$  is invariably written for this constant.

$$2. \quad 1^2 - 2^2 + 3^2 - 4^2 + \dots = (2^{n+1} - 1) \frac{B_{n+1} \sin \frac{\pi n}{2}}{n+1}$$

$$\text{Sol. } (1 - 2^{n+1}) C_n = (1^2 - 2^2 + \dots) - 2^{n+1} (1^2 - 2^2 + \dots)$$

$$= 1^2 - 2^2 + 3^2 - \dots$$

$$\text{Cor. } \phi_n\left(\frac{1}{2}\right) = 2\left(1 - \frac{1}{2^n}\right) \frac{B_{n+1} \cos \frac{\pi n}{2}}{n+1}$$

$$\text{Sol. } \phi_n\left(\frac{1}{2}\right) = 1^2 - \left(\frac{1}{2}\right)^2 + 2^2 - \left(\frac{1}{2}\right)^2 + \dots$$

$$= - \frac{1}{2^n} (1^2 - 2^2 + 3^2 - \dots) = \left(2 - \frac{1}{2^n}\right) \frac{B_{n+1} \cos \frac{\pi(n+1)}{2}}{n+1}$$

$$3. \quad (a+6)^2 + (a+16)^2 + (a+26)^2 + \dots + \{ \phi_n(a+6) \}^2$$

$$= 6^n \left\{ \phi_n\left(\frac{a}{6} + 1\right) - \phi_n\left(\frac{a}{6}\right) \right\}$$

$$\text{Sol. L.S.} = 6^n \left\{ \left(1 + \frac{a}{6}\right)^2 + \left(2 + \frac{a}{6}\right)^2 + \dots + \left(2 + \frac{a}{6}\right)^2 \right\} = \text{R.S.}$$



$$4. \frac{B_{1-n}}{1-n} \sin \frac{\pi n}{2} = S_n = \frac{(2\pi)^n}{2 \Gamma} B_n$$

From this we can find  $B_n$  for negative values of  $n$

Sol.  $\frac{B_{1+n}}{1+n} \cos \frac{\pi(1+n)}{2}$  is the constant of  $1^m + 2^m + 3^m + \dots$

$\therefore \frac{B_{1-n}}{1-n} \cos \frac{\pi(1-n)}{2}$  is that of  $\frac{1}{1^n} + \frac{1}{2^n} + \dots = S_n$

$$\therefore \frac{B_{1-n}}{1-n} \sin \frac{\pi n}{2} = \frac{(2\pi)^n}{2 \Gamma} B_n$$

$$\text{Ex. 1. } B_{-2} = 2S_3; B_{-4} = -4S_5; B_{-6} = 6S_7; B_{-8} = -8S_9 \text{ etc}$$

$$2. \Gamma_{\frac{1}{2}} = \sqrt{\pi}; \text{ sol. } -\frac{B_{1\frac{1}{2}}}{1\frac{1}{2}} \sin \frac{\pi}{4} = \frac{(2\pi)^{-\frac{1}{2}}}{2 \Gamma_{\frac{1}{2}}} B_{-\frac{1}{2}}$$

Again  $\frac{B_{-\frac{1}{2}}}{-\frac{1}{2}} \sin \frac{\pi}{4} = \frac{(2\pi)^{\frac{1}{2}}}{2 \Gamma_{\frac{1}{2}}} B_{\frac{1}{2}}$  multiplying the

results we have  $\frac{2}{3} = \frac{2}{3} \cdot \frac{\pi}{(\Gamma_{\frac{1}{2}})^2} \therefore \Gamma_{\frac{1}{2}} = \sqrt{\pi}$ .

3. In a similar manner we can prove that

$$\Gamma_{n-1} \Gamma_n = \pi \operatorname{Cosec} \pi n.$$

$$4. \pi \left( \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}+\sqrt{4}} + \frac{1}{\sqrt{4}+\sqrt{6}} - \frac{1}{\sqrt{6}+\sqrt{8}} + \dots \right)$$

$$= \frac{1}{\sqrt{1}} + \frac{1}{3\sqrt{3}} + \frac{1}{5\sqrt{5}} + \frac{1}{7\sqrt{7}} + \dots$$

$$\text{Sol. } L.S = \frac{\pi}{\sqrt{2}} \{ 1 - (\sqrt{2}-1) + (\sqrt{3}-\sqrt{2}) - (\sqrt{4}-\sqrt{3}) + \dots \}$$

$$= \pi \sqrt{2} (\sqrt{2}-\sqrt{2} + \sqrt{3}-\sqrt{4} + \dots) = 2(2\sqrt{2}-1) \frac{B_{1\frac{1}{2}}}{\Gamma_{\frac{1}{2}}} \cdot \frac{\pi}{2}$$

$$= (1 - \frac{1}{2\sqrt{2}}) \left( \frac{1}{\sqrt{1}} + \frac{1}{2\sqrt{2}} + \frac{1}{3\sqrt{3}} + \dots \right)$$

$$= \frac{1}{\sqrt{1}} + \frac{1}{3\sqrt{3}} + \frac{1}{5\sqrt{5}} + \dots$$

$$5. \frac{2\pi (\frac{1}{\sqrt{1}} - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} - \frac{1}{\sqrt{4}} + \dots)}{(2\sqrt{2} + 4\sqrt{2}) (\frac{1}{\sqrt{1}} - \frac{1}{\sqrt{4}} + \frac{1}{\sqrt{9}} - \dots)} = \frac{1}{3}$$

$$6. \sqrt{2+4x} - (\frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{x}})$$

= (2+1) ( \frac{1}{\sqrt{1}} - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} - \dots ) when x is great

$$7. \frac{2}{3} \sqrt{(x+\frac{1}{4})(x+\frac{1}{2})(x+\frac{3}{4})} - (\sqrt{1} + \sqrt{2} + \dots + \sqrt{x})$$

= \frac{1}{4\sqrt{1}} ( \frac{1}{\sqrt{1}} + \frac{1}{2\sqrt{2}} + \frac{1}{3\sqrt{3}} + \frac{1}{4\sqrt{4}} + \dots )

$$8. \frac{2}{5} \sqrt{x(x+\frac{1}{2})(x+\frac{1}{2})(x+\frac{3}{4})(x+1) + \frac{5}{168}(x+\frac{1}{2})}$$

- (1\sqrt{1} + 2\sqrt{2} + 3\sqrt{3} + \dots + x\sqrt{x})

= \frac{2}{167\pi} ( \frac{1}{\sqrt{1}} + \frac{1}{2\sqrt{2}} + \frac{1}{9\sqrt{3}} + \frac{1}{16\sqrt{4}} + \dots )

$$9. (a+6)^2 - (a+26)^2 + (a+36)^2 - \dots = 6^2 \{ \phi_{12}(\frac{a}{26}) - \phi_{12}(\frac{a-6}{26}) \}$$

$$i. \frac{(x^n+x)^n}{2} = \frac{\pi}{11} \frac{\phi_{2n-1}(x)}{2n-1} + \frac{\pi(n-1)(n-2)}{13} \phi_{2n-3}(x) + \frac{\pi(n-1)(n-2)(n-3)(n-4)}{15} \phi_{2n-5}(x) + \dots$$

$$ii. \frac{(x+\frac{1}{2})(x^n+x)^n}{2} = \frac{(n+\frac{1}{2})}{11} \phi_{2n}(x) + \frac{\pi(n-1)(n-\frac{1}{2})}{13} \phi_{2n-2}(x) + \frac{\pi(n-1)(n-2)(n-3)(n-\frac{3}{2})}{15} \phi_{2n-4}(x) + \dots$$

$$Sol. \frac{(x^2+x)^n - (x^2-x)^n}{2} = \frac{\pi}{11} x^{2n-1} + \frac{\pi(n-1)(n-2)}{13} x^{2n-3} + \dots$$

change x to x-1, x-2 &c up to 1 & add up all the terms

$$\frac{(x+\frac{1}{2})(x^2+x)^n - (x-\frac{1}{2})(x^2-x)^n}{2} = \frac{\pi}{2} \{ (x^2+x)^n - (x^2-x)^n \} + \frac{\pi}{2} \{ (x^2+x)^n + (x^2-x)^n \}$$

& proceed as in i.

Cor. If  $x^2 + x = y$  &  $x + \frac{1}{2} = a$ , then

$$1. \phi_1(x) = \frac{y}{2}; \phi_2(x) = a \frac{y}{3}; \phi_3(x) = \frac{y^2}{4}; \phi_4(x) = \frac{a}{5} y(y - \frac{1}{3})$$

$$\phi_5(x) = \frac{y^2}{6}(y - \frac{1}{2}); \phi_6(x) = \frac{a}{7} y(y - y + \frac{1}{3}); \phi_7(x) = \frac{y^2}{8}(y - \frac{2}{3}y + \frac{1}{3})$$

$$\phi_8(x) = \frac{a}{9} y(y^3 - xy^2 + \frac{y}{3} - \frac{3}{5}); \phi_9(x) = \frac{y^2}{10}(y-1)(y^2 - \frac{2}{3}y + \frac{1}{3})$$

$$\phi_{10}(x) = \frac{a}{11} y(y-1)(y^3 - \frac{7}{3}y^2 + \frac{10}{3}y - \frac{5}{3})$$

$$\phi_{11}(x) = \frac{y^2}{12}(y^4 - 4y^3 + 8\frac{1}{2}y^2 - 10y + 5)$$

$$2. i. (\frac{1+\sqrt{5}}{2})^9 + (\frac{3+\sqrt{5}}{2})^9 + \dots + (\frac{2n-1+\sqrt{5}}{2})^9 = \phi_9(\frac{2n-1+\sqrt{5}}{2})$$

$$ii. (\frac{2+\sqrt{5}}{2})^{10} + (\frac{3+\sqrt{5}}{2})^{10} + \dots + (\frac{2n-1+\sqrt{5}}{2})^{10} = \phi_{10}(\frac{2n-1+\sqrt{5}}{2})$$

iii. If  $n$  be even then

$$1^n + 3^n + 5^n + 7^n + \dots + (2p-1)^n = 2^n \phi_n(p - \frac{1}{2})$$

7. If  $n$  is a positive integer excluding zero

$$\phi_n(x-1) + (-1)^n \phi_n(x) = 0,$$

Sol. Let  $L.S. = \psi(x)$ ; then  $\psi(x+1) - \psi(x) = 0$

Cor. If  $n > 1$ , then  $\phi_n(x)$  is divisible by  $\frac{x^2(x+1)}{4}$  or  $\frac{x(x+\frac{1}{2})(x+1)}{3}$  according as  $n$  is odd or even

$$8. \phi_n(x) = -n x S_{1-n} - \frac{n(n-1)x^2 S_{2-n}}{2} - \frac{n(n-1)(n-2)x^3 S_{3-n}}{6}$$

$$- \dots = -B_n x \cos \frac{\pi n}{2} - \frac{n}{2} B_{n-1} x^2 \sin \frac{\pi n}{2} +$$

$$\frac{n(n-1)}{6} B_{n-2} x^3 \cos \frac{\pi n}{2} + \frac{n(n-1)(n-2)}{24} B_{n-3} x^4 \sin \frac{\pi n}{2}$$

- etc; Sol. Apply VI 6.

9.  $\phi_n(x) = \{ - (1+x)^n + x^2 - (2+x)^n + 2x \dots + \phi_n(\frac{x-n+1}{n}) \}$   
 10.  $\phi_n(x) = x^n \{ \phi_n(\frac{x}{n}) + \phi_n(\frac{x-1}{n}) + \phi_n(\frac{x-2}{n}) + \dots + \phi_n(\frac{x-n+1}{n}) \}$   
 $= (x^{n+1} - 1) \frac{\beta_{n+1}}{n+1} \frac{\sin \frac{\pi x}{2}}{2} \text{ etc } (1 - x^{n+1}) S_{-n}$

Sol. Apply VI 7.

Ans.  $\phi_n(\frac{x}{n}) + \phi_n(\frac{x-1}{n}) + \phi_n(\frac{x-2}{n}) + \dots + \phi_n(\frac{x-n+1}{n})$   
 $= (x - x^{-n}) S_{-n}$

11. If  $n$  is a negative integer, then  
 $\phi_n(x-1) + (-1)^n \phi_n(x) = \{ 1 + (-1)^n \} S_{-n} + \frac{(-1)^n}{[-n]} d_{-(n+1)x} \pi \cot \pi x$

Sol.  $\phi_n(x-1) - \phi_{-n}(x) = -\pi \cot \pi x$  by II 10.

Differentiate both sides  $n$  times.

Note. The above theorem is true even for positive integral values of  $n$  and hence VII 7 can be deduced from VII 11.

N.B. The following method is very useful in finding the derivatives of  $\pi \cot \pi x$ . Let  $\pi \cot \pi x = y$ ; then the coeff<sup>ts</sup> in the coeff<sup>ts</sup> of  $\pi^n$  are the same as those in the expansion of  $(\tan^{-1} \frac{1}{y})^{-n}$ .

Each derivative is divisible by  $y^2 + 1$  so that the last term can be exactly found.

Write under each term the quotient obtained

$\pi y$  by dividing the sum of the products of the  
 $\pi^2(y^2+1)$  coeff<sup>s</sup>. and the index of that term and of  
 $\pi^3(y^3+y)$  the preceding term by the index of  $\pi$ .

$$\pi^4(y^4 + \frac{4}{3}y^2 + \frac{1}{3})$$

$$\pi^5(y^5 + \frac{5}{3}y^3 + \frac{2}{3}y)$$

$$\pi^6(y^6 + 2y^4 + \frac{17}{15}y^2 + \frac{2}{15})$$

$$\pi^7(y^7 + \frac{7}{3}y^5 + \frac{77}{45}y^3 + \frac{17}{45}y)$$

$$\pi^8(y^8 + \frac{8}{3}y^6 + \frac{12}{5}y^4 + \frac{248}{315}y^2 + \frac{17}{315})$$

$$\pi^9(y^9 + 3y^7 + \frac{16}{5}y^5 + \frac{88}{63}y^3 + \frac{62}{315}y)$$

$$\pi^{10}(y^{10} + \frac{10}{3}y^8 + \frac{37}{9}y^6 + \frac{424}{189}y^4 + \frac{1382}{2835}y^2 + \frac{62}{2835})$$

Cor. For all values of  $a$

$$i. \phi_n(x) - 2^n \left\{ \phi_n\left(\frac{x}{2}\right) + \phi_n\left(\frac{x-1}{2}\right) \right\} = (1 - 2^{n+1}) S_{-n}$$

$$ii. \phi_n\left(-\frac{1}{2}\right) = (2 - \frac{1}{2^n}) S_{-n}$$

$$iii. \phi_n\left(-\frac{1}{3}\right) + \phi_n\left(-\frac{2}{3}\right) = (3 - \frac{1}{3^n}) S_{-n}$$

$$iv. \phi_n\left(-\frac{1}{4}\right) + \phi_n\left(-\frac{3}{4}\right) = (2 + \frac{1}{2^n} - \frac{1}{4^n}) S_{-n}$$

$$v. \phi_n\left(-\frac{1}{6}\right) + \phi_n\left(-\frac{5}{6}\right) = (1 + \frac{1}{2^n} + \frac{1}{3^n} - \frac{1}{6^n}) S_{-n}$$

Ex. If  $n$  is a positive odd integer show that

$$i. \phi_n\left(-\frac{1}{3}\right) = (3 - \frac{1}{3^n}) \frac{S_{-n}}{2}$$

$$ii. \phi_n\left(-\frac{1}{4}\right) = (1 + \frac{1}{2^{n+1}} - \frac{1}{2^{2n+1}}) S_{-n}$$

$$iii. \phi_n\left(-\frac{1}{6}\right) = (1 + \frac{1}{2^n} + \frac{1}{3^n} - \frac{1}{6^n}) \frac{S_{-n}}{2}$$

$$iv. \phi_n\left(-\frac{1}{5}\right) + \phi_n\left(-\frac{2}{5}\right) = (5 - \frac{1}{5^n}) \frac{S_{-n}}{2}$$

$$v. \phi_n\left(-\frac{1}{6}\right) + \phi_n\left(-\frac{5}{6}\right) = (2 + \frac{1}{2^{n+1}} - \frac{1}{2^{2n+1}}) S_{-n}$$

vii.  $\phi_n(\frac{1}{10}) + \phi_n(\frac{3}{10}) = (5 + \frac{1}{5}n - \frac{1}{10}n^2) \frac{S_{-n}}{2}$

viii.  $\phi_n(\frac{1}{12}) + \phi_n(\frac{5}{12}) = (6 + \frac{1}{6}n - \frac{1}{12}n^2) \frac{S_{-n}}{2}$

12.  $2^n \{ \phi_n(\frac{1}{8}) - \phi_n(\frac{5}{8}) \} = (2^n + 1) \{ \phi_n(\frac{1}{3}) - \phi_n(\frac{2}{3}) \}$

Sol.  $\phi_n(\frac{1}{3}) - 2^n \{ \phi_n(\frac{1}{8}) + \phi_n(\frac{5}{8}) \} = (2^{2n+1} - 1) S_{-n}$   
 $\phi_n(\frac{2}{3}) - 2^n \{ \phi_n(\frac{1}{3}) + \phi_n(\frac{5}{3}) \} = (2^{2n+1} - 1) S_{-n}$  } by (viii)

$\therefore 2^n \{ \phi_n(\frac{1}{8}) - \phi_n(\frac{5}{8}) \} = (2^n + 1) \{ \phi_n(\frac{1}{3}) - \phi_n(\frac{2}{3}) \}$

13. Since all these theorems and the following theorems are true for all values of n, the properties of  $\pi \approx \frac{1}{2}, \sqrt{x}, \frac{1}{n} + \frac{1}{n^2} + \dots + \frac{1}{n^2}$  &c &c are only their particular cases.

Ex-1.  $\frac{1}{13} + \frac{1}{33} + \frac{1}{53} + \dots = \frac{7}{8} S_3$

2.  $\frac{1}{12} + \frac{1}{42} + \frac{1}{72} + \dots = \frac{2}{81\sqrt{3}} \pi^3 + \frac{13}{27} S_3$

3.  $\frac{1}{13} + \frac{1}{53} + \frac{1}{93} + \dots = \frac{\pi^2}{64} + \frac{7}{16} S_3$

4.  $\frac{1}{12} + \frac{1}{72} + \frac{1}{132} + \dots = \frac{\pi^3}{36\sqrt{3}} + \frac{91}{216} S_3$

13. If  $C_n$  be the constant of  $\frac{(\log 1)^2}{1} + \frac{(\log 2)^2}{2} + \dots$

then  $S_{n+1} = \frac{1}{n} + C_0 - \frac{\pi^2}{6} C_1 + \frac{\pi^4}{72} C_2 - \frac{\pi^6}{120} C_3 + \dots$   
 $= \frac{1}{n} + .5772156649 + .0728158455\pi$   
 $- (.00485\pi^2 + .00034\pi^3) + E$

where E, the error is less than  $(\frac{\pi}{10})^4$

20.  
 Sol. It is proved in  $\nabla$  26 Cor. 1. that  $S_{m, -n}$  is finite when  $m=0$ ; the remaining part is obtained from VI. 13. N.B. The theorem is true for all values of  $n$ .

Ex. 1.  $S_{1+n} + S_{1-n} = \frac{2C_0}{1 + .00839n^2 + .0001n^4, \dots}$

2.  $\frac{1}{1\sqrt{1}} + \frac{1}{2\sqrt{2}} + \frac{1}{3\sqrt{3}} + \dots = 10.58444842$

3.  $\frac{1}{1\sqrt{1}} + \frac{1}{2\sqrt{2}} + \frac{1}{3\sqrt{3}} + \dots = 2.6123752$  correct

4.  $\frac{1}{1^2\sqrt{1}} + \frac{1}{2^2\sqrt{2}} + \frac{1}{3^2\sqrt{3}} + \dots = 1.341490$

5.  $B_{1/2} = .4409932$ ;  $B_{-1/2} = -1.032627$

6.  $B_{1/3} = -.9420745$ ;  $B_{-1/3} = -1.3841347$

7.  $B_{-1/2} = -1.847228$ .

14.  $\frac{1}{2(2^n-1)} + \frac{1}{3(3^n-1)} + \frac{1}{4(4^n-1)} + \dots$   
 $= \frac{.7946786 - \log n}{n} + .2115922$   
 $- .0060680n - .0000028n^3 + \dots$

Sol. We can easily prove that  $L.S. = \frac{e^{-\log n}}{n} + \dots$   
 where  $C$  is the constant in  $\frac{1}{2\log 2} + \frac{1}{3\log 3} + \frac{1}{4\log 4} + \dots$   
 If  $n=1$  then  $L.S. = \frac{1}{1.2} + \frac{1}{2.3} + \dots = 1$ ; hence  $C = 1$ .

Cor. 1.  $\frac{1}{2\log 2} + \frac{1}{3\log 3} + \dots + \frac{1}{n\log n}$   
 $= .7946786 + \log \log (n+1)$  nearly.

2.  $\frac{1}{2^{m+1}\log 2} + \frac{1}{3^{m+1}\log 3} + \frac{1}{4^{m+1}\log 4} + \frac{1}{5^{m+1}\log 5} + \dots$

$$= -\log x + .2174630 + .4227843x$$

$$- .0364079x^2 + .001617x^3 + .000095x^4$$

$$- .00002x^5 - \dots$$

Sol. Integrate III 13.

$$15. \frac{\phi_{2n}(x-1) - \phi_n(x)}{1 \sqrt{2}} = -\cos \frac{\pi x}{2} \left\{ \frac{\sin 2\pi x}{(2\pi)^{2n+1}} + \frac{\sin 4\pi x}{(4\pi)^{2n+1}} \right.$$

$$\left. + \frac{\sin 6\pi x}{(6\pi)^{2n+1}} + \dots \right\}$$

Sol.  $\phi_{2n}(x-1) - \phi_n(x) = (1-x)^{2n} - x^{2n} + (2-x)^{2n} - (1+x)^{2n} + (3-x)^{2n}$   
 $- (2+x)^{2n} + \dots$ ; then arrange the terms in ascending powers of  $x$  and substitute  $\frac{B_n}{n} \cos \frac{\pi x}{2}$  for  $S_{1-n}$ . Similarly

$$16. \frac{\phi_{2n}(x-1) + \phi_n(x) - 2S_{-n}}{1 \sqrt{2}} = \sin \frac{\pi x}{2} \left\{ \frac{\cos 2\pi x}{(2\pi)^{2n+1}} + \frac{\cos 4\pi x}{(4\pi)^{2n+1}} + \frac{\cos 6\pi x}{(6\pi)^{2n+1}} + \dots \right\}$$

N.B. The above two theorems are true for all values of  $x$  when  $n$  is an integer but when  $n$  is fractional they are true only when  $x$  lies between 0 and 1  
 Cor. If  $\frac{p}{q}$  lies between 0 and 1 and  $p, q$  are integers

$$i. \frac{(2\pi p)^n}{1 \sqrt{2}} \left\{ \phi_{n-1}\left(\frac{p}{q}-1\right) - \phi_{n-1}\left(\frac{p}{q}\right) \right\} = -\sin \frac{\pi p}{2} \left[ \left\{ S_n - \phi_{n-1}\left(\frac{p}{q}-1\right) \right\} \times \right.$$

$$\left. \sin \frac{2\pi p}{q} + \left\{ S_n - \phi_{n-1}\left(\frac{p}{q}\right) \right\} \sin \frac{4\pi p}{q} + \left\{ S_n - \phi_{n-1}\left(\frac{p}{q}\right) \right\} \right]$$



$$\times \sin \frac{6\pi p}{q} + \dots + \left\{ S_n - \phi_n \left( \frac{q-1}{q} \right) \right\} \sin \frac{(2q-2)\pi p}{q}$$

$$\text{ii. } \frac{(2\pi q)^n}{4^{n-1}} \left\{ \phi_{n-1} \left( \frac{p}{q} \right) + \phi_n \left( -\frac{p}{q} \right) - 2S_{1-n} \left( 1 - \frac{1}{q} \right) \right\}$$

$$= -\cos \frac{\pi a}{2} \left[ \left\{ S_n - \phi_n \left( \frac{1}{q} \right) \right\} \cos \frac{2\pi p}{q} + \left\{ S_n - \phi_n \left( \frac{q-1}{q} \right) \right\} \cos \frac{4\pi p}{q} \right.$$

$$\left. + \left\{ S_n - \phi_n \left( \frac{2}{q} \right) \right\} \cos \frac{6\pi p}{q} + \dots + \left\{ S_n - \phi_n \left( \frac{q-1}{q} \right) \right\} \cos \frac{(2q-2)\pi p}{q} \right]$$

$$17. \phi_n \left( \frac{1}{2} \right) - \phi_n \left( -\frac{1}{2} \right) = 2 \cdot \frac{E_{n+1}}{4^{n+1}} \cos \frac{\pi a}{2}$$

Sol. Put  $x = \frac{1}{2}$  in VII. 15.

$$\text{Cor. } 1^n - 3^n + 5^n - 7^n + \dots = \frac{1}{2} E_{n+1} \cos \frac{\pi a}{2}$$

$$18. E_{1-n} \cos \frac{\pi a}{2} = \left( \frac{\pi}{2} \right)^n \frac{E_n}{\Gamma(n)}$$

Sol. Change  $n$  to  $-n$  in VII 17 Cor.

$$\text{Cor. } \pi \left\{ \frac{1}{2} - \frac{1}{\sqrt{1+\sqrt{3}}} + \frac{1}{\sqrt{3+\sqrt{5}}} - \frac{1}{\sqrt{5+\sqrt{7}}} + \dots \right\}$$

$$= \frac{1}{\sqrt{1}} - \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{5}} - \frac{1}{\sqrt{7}} + \dots$$

19. If  $\frac{p}{q}$  lies between 0 & 1,  $p$  being any integer &  $q$  an odd integer, then

$$\text{i. } \frac{(2\pi q)^n}{4^{n-1}} \left\{ \phi_{n-1} \left( \frac{p}{q} \right) - \phi_n \left( -\frac{p}{q} \right) \right\} = \sin \frac{\pi a}{2} \left[ \left\{ \phi_n \left( \frac{1}{q} \right) - \phi_n \left( -\frac{1}{q} \right) \right\} \right.$$

$$\left. \times \sin \frac{2\pi p}{q} + \left\{ \phi_n \left( \frac{2}{q} \right) - \phi_n \left( -\frac{2}{q} \right) \right\} \sin \frac{4\pi p}{q} + \dots \text{to } \frac{q-1}{2} \text{ terms} \right]$$

$$\text{ii. } \frac{(2\pi q)^n}{4^{n-1}} \left\{ \phi_{n-1} \left( \frac{p}{q} \right) + \phi_n \left( -\frac{p}{q} \right) - 2S_{1-n} \left( 1 - \frac{1}{q} \right) \right\}$$

$$= \cos \frac{\pi a}{2} \left[ \left\{ \phi_n \left( \frac{1}{q} \right) - \phi_n \left( -\frac{1}{q} \right) \right\} \cos \frac{2\pi p}{q} + \left\{ \phi_n \left( \frac{2}{q} \right) - \phi_n \left( -\frac{2}{q} \right) \right\} \cos \frac{4\pi p}{q} + \dots \right]$$

$x \cos \frac{2\pi x}{l} + \&c$  to  $\frac{l-1}{2}$  terms]

$$\text{Coroll. } \frac{l^{n-1}}{l} \phi_n(x) = \frac{\sin \pi x}{\pi^{n+1}} \cos(\pi x + \frac{\pi a}{2}) + \frac{\sin 2\pi x}{(2\pi)^{n+1}} \cos(2\pi x + \frac{\pi a}{2}) \\ + \frac{\sin 3\pi x}{(3\pi)^{n+1}} \cos(3\pi x + \frac{\pi a}{2}) + \&c$$

Sol. Combine the results of VII 15 & 16.

$$2. \frac{1}{\sqrt{x}} - \frac{1}{\sqrt{1-x}} + \frac{1}{\sqrt{1+x}} - \frac{1}{\sqrt{2-x}} + \frac{1}{\sqrt{2+x}} - \&c \\ = 2 \left( \frac{\sin 2\pi x}{\sqrt{1}} + \frac{\sin 4\pi x}{\sqrt{2}} + \frac{\sin 6\pi x}{\sqrt{3}} + \&c \right)$$

$$2c. \frac{(6\pi)^2}{2!2! \sqrt{3}} \left\{ \phi_{n-1}(\frac{1}{3}) - \phi_n(\frac{1}{3}) \right\} = \left\{ \phi_{n-1}(\frac{1}{3}) - \phi_n(\frac{1}{3}) \right\} \sin \frac{\pi n}{2}$$

Sol. Put  $p=1$  &  $q=3$  in VII 19. i.

$$2l. \phi(0) + \frac{n}{l} \phi(1)x + \frac{n(n-1)}{l^2} \phi(2)x^2 + \frac{n(n-1)(n-2)}{l^3} \phi(3)x^3 \\ + \&c = (1+x)^n \phi_{\infty}(\frac{nx}{1+x}), \text{ where}$$

$$\phi_n(x) = \phi_{n-1}(x) + \frac{n P_{n-1}}{l L_n} \phi_{n-1}^2(x) + \frac{(n P_{n-1})^2}{l^2 (L_n)^2} \phi_{n-1}^{2n}(x) \\ + \frac{(n P_{n-1})^3}{l^3 (L_n)^3} \phi_{n-1}^{3n}(x) + \&c \text{ and } \phi_1(x) = \phi(x).$$

$$\text{and } P_n = 1^2 x - 2^2 x^2 + 3^2 x^3 - 4^2 x^4 + \&c.$$

Sol. Prove the theorem by substituting  $e^{ax}$  for  $\phi(x)$  or proceed as in III 10.

$$\text{Cor. } \left\{ \phi(0) + \frac{n}{l} x \phi(1) + \frac{n(n-1)}{l^2} x^2 \phi(2) + \&c \right\} (1+x)^{-n} \\ = \phi(\frac{nx}{1+x}) + \frac{nx}{(1+x)^2} \phi''(\frac{nx}{1+x}) + \&c.$$

22. If  $A_n = (1^n + 2^n + 3^n + \dots)(1 + \cos \pi n)$ , then

$$2^n + 6^n + 12^n + 20^n + \dots = A_n + \frac{\pi}{2L} A_{n+\pi} + \frac{\pi(n-1)}{L^2} A_{n+2}$$

Ex.  $10 = \pi^2 + \frac{1}{2^3} + \frac{1}{6^3} + \frac{1}{12^3} + \frac{1}{20^3} + \dots$

23.  $\log_e \Gamma(x) = (x + \frac{1}{2}) \log x - x + \frac{1}{2} \log 2\pi + \frac{B_2}{1 \cdot 2x} - \frac{B_4}{3 \cdot 4x^3} + \frac{B_6}{5 \cdot 6x^5} - \dots$

Sol. Equate the coeff<sup>s</sup> of  $x$  in VIII 1; the coeff<sup>t</sup> of  $x$  in  $S_n$

$$= \text{that in } -\frac{1x}{\pi(2\pi)^x} S_{n+1} \sin \frac{\pi x}{2} = \text{that of } x \text{ in}$$

$$-\frac{B_2}{2} (1 - x \log_e 2\pi + \dots) (\frac{1}{x} + C_0 - \dots) (1 - x C_0 + \dots)$$

$$= \frac{1}{2} \log_e 2\pi. \text{ or as follows}$$

Let  $c$  be the constant in  $\log_e \Gamma(x)$  &  $f(x) = \log_e \frac{\Gamma(x)}{\Gamma(x-1)}$

then we see that  $f(x) - f(x-1) = \log_e 2$ .

$\therefore \log_e \frac{\Gamma(x)}{\Gamma(x-1)} - x \log_e 2 = \text{some constant}$ ; by put-

ting  $x=0$  we find this constant is  $-\frac{1}{2} \log_e \pi$ .

But the constant in  $\log_e \frac{\Gamma(x)}{\Gamma(x-1)} = \frac{1}{2} \log_e 2 - c$ .

$$\therefore c = \frac{1}{2} \log_e 2\pi = .918938533204673.$$

Cor. When  $x$  is great  $\frac{e^x \Gamma(x)}{x^x} = \sqrt{2\pi x + \frac{\pi}{3}}$  nearly.

24.  $\Gamma(x-1) \Gamma(x) = \pi \operatorname{Cosec} \pi x$ ; Cor  $\Gamma(\frac{1}{2}) = \sqrt{\pi}$ .

25.  $\Gamma(\frac{x}{n}) \Gamma(\frac{x-1}{n}) \Gamma(\frac{x-2}{n}) \Gamma(\frac{x-3}{n}) \dots \Gamma(\frac{x-n+1}{n}) = \frac{(2\pi)^{\frac{n-1}{2}}}{n^{x+\frac{1}{2}}} \Gamma(x)$

Cor. 1.  $\Gamma(\frac{1}{2}) \Gamma(\frac{3}{2}) \Gamma(\frac{5}{2}) \dots \Gamma(\frac{2n-1}{2}) = \frac{(2\pi)^{\frac{n}{2}}}{\sqrt{2\pi n}}$

2.  $\Gamma(\frac{1}{3}) = \sqrt{\Gamma(\frac{2}{3})} \sqrt[3]{\Gamma(\frac{4}{3})} \sqrt[4]{\Gamma(\frac{5}{3})}$

$$3. \frac{\sqrt{x}}{\sqrt[3]{x}} = \frac{x^{\frac{1}{2}}}{x^{\frac{1}{3}}} = x^{\frac{1}{6}}$$

$$\ln \log \sqrt{x-\frac{1}{2}} = x \log x - x + \frac{1}{2} \log 2\pi + (1-\frac{1}{2}) \frac{B_2}{1 \cdot 2 x} -$$

$$(1-\frac{1}{2^2}) \frac{B_4}{3 \cdot 4 x^3} + (1-\frac{1}{2^4}) \frac{B_6}{5 \cdot 6 x^5} - 2x e$$

$$26. \log \sqrt{x} = -C_0 x + \frac{S_2}{2} x^2 - \frac{S_3}{3} x^3 + \frac{S_4}{4} x^4 - \dots$$

$$\text{i.e. } \log_e \frac{\sqrt{x+1}}{2} = .9227948351x + .1974670334x^2$$

$$- .0256856344x^3 + .0049558084x^4$$

$$- .0011355510x^5 + .0002863487x^6$$

$$- .0000766825x^7 + .0000213883x^8$$

$$- .0000061409x^9 + .0000054047x^{10}$$

$$\text{Ex. 1 } \log_e \sqrt{-\frac{1}{3}} = .5341990853$$

$$2. \log_e \sqrt{-\frac{1}{6}} = .1211436313$$

$$3. \log_e \sqrt{-\frac{1}{10}} = .0663762397.$$

$$27. i 2\pi x \left\{ 1 + \left(\frac{x}{n+1}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+2}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+3}\right)^2 \right\} \dots \text{ ad inf}$$

$$= \left(\frac{ix}{x^n}\right)^2 (e^{\pi x} - e^{-\pi x}) e^{-\frac{S_2}{2x^2} + \frac{S_4}{2x^4} - \frac{S_6}{3x^6} + \dots}$$

$$\text{where } S_p = 1^p + 2^p + 3^p + \dots + n^p.$$

Sol. Let  $L.S = f(x)$ ; then  $\frac{f(n+1)}{f(n)} = 1 + \left(\frac{x}{n}\right)^2$ ; find  $f(x)$  by applying  $\Delta$  or in anyway.

H.P.  $\theta = \cos 2\pi x$  exactly or very nearly according as  $2\pi x$  is an integer or not.

Sol. For even values of  $2n$ ,  $e^{\pi x} - e^{-\pi x}$  appears in  
 And is but for odd values  $e^{\pi x} + e^{-\pi x}$

$$\begin{aligned} \text{ii. } & 2\pi (x^2 + n^2)^{n+\frac{1}{2}} \left\{ 1 + \left(\frac{x}{n+1}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+2}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+3}\right)^2 \right\} \&c \\ & = (2n)^2 (e^{\pi x} - e^{-\pi x}) e^{2\pi x} - 2x \tan^{-1} \frac{x}{n} - \frac{13x S_2}{x} - \frac{B_4 S_4}{2x^2} \\ & - \frac{B_6 S_6}{3x^3} - \&c \text{ where } S_p = \frac{\pi}{x} - \frac{p(p+1)}{15} \left(\frac{x}{n}\right)^3 - \\ & \frac{p(p+1)(p+2)(p+3)}{15} \left(\frac{x}{n}\right)^5 - \&c. \end{aligned}$$

Sol. Find  $S_2, S_4, S_6$  etc in the previous theorem  
 by VII 1. and then simplify.

$$\begin{aligned} \text{iii. } & 2\pi (n^2 + x^2)^{n-\frac{1}{2}} \left\{ 1 + \left(\frac{x}{n}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+1}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+2}\right)^2 \right\} \&c \\ & = (2n-1)^2 e^{2n + 2x\beta} - 2 \frac{B_2 \cos \beta}{1.2n} + \frac{2 B_4 \cos 3\beta}{8.4n^2} - \&c \\ & \times (1 - e^{-2\pi x}) \text{ where } n^2 = n^2 + x^2 \& \tan \beta = \frac{x}{n}. \end{aligned}$$

$$\begin{aligned} \text{Sol. } & \frac{(n+x)}{(n-x)} = \frac{(n+x)}{(n-x)} \cdot \frac{(1^2+x^2)(2^2+x^2)(3^2+x^2)}{(1^2+x^2)(2^2+x^2)(3^2+x^2)} \\ & \dots (n^2+x^2) = \frac{(2n)^2}{\left\{ 1 + \left(\frac{x}{n+1}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+2}\right)^2 \right\} \&c} \text{ and int} \\ & \text{then find } \frac{(n+x)}{(n-x)} \text{ by VII 23.} \end{aligned}$$

- N.B. i. is useful only when  $x$  is great &  $n$  small  
 ii. when  $x$  is great when compared to  $n$   
 iii. in all cases.

1.  $\frac{B_2}{n} \cos \frac{\pi n}{2} + \frac{1}{n}$ , when  $n$  varies, is a finite quantity which is invariably denoted by  $C_0$ ; it is the constant of  $S_1$ , and its value is found from VIII 2 to be  $\cdot 577215664901533$  and  $\log_e C_0 = + \cdot 56145948356$ .

Sol. L.S. in VII 1 is finite when  $n=1$ .

$$\therefore \frac{B_2}{n} \cos \frac{\pi n}{2} + \frac{x^2}{n} \text{ is finite when } n=0$$

$$\text{i.e. } \frac{B_2}{n} \cos \frac{\pi n}{2} + \frac{1}{n} + \frac{x^3-1}{n} \text{ is finite when } n=0$$

$$\text{But } \frac{x^3-1}{n} = \log_e x \text{ when } n=0.$$

2.  $1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{x} = \sum \frac{1}{x}$  or  $\phi(x)$ . (Suppose).

$$\sum \frac{1}{x} = C_0 + \log_e x + \frac{1}{2x} - \frac{B_2}{2x^2} + \frac{B_4}{4x^4} - \frac{B_6}{6x^6} + \dots$$

$$3. \sum \frac{1}{x} = 1 - \frac{1}{x+1} + \frac{1}{2} - \frac{1}{x+2} + \frac{1}{3} - \frac{1}{x+3} + \dots$$

$$= \frac{x}{1(1+x)} + \frac{x}{2(2+x)} + \frac{x}{3(3+x)} + \dots$$

$$4. \sum \frac{1}{x} = xS_2 - x^2S_3 + x^3S_4 - x^4S_5 + \dots$$

$$5. \sum \frac{1}{x-1} = \sum \frac{1}{-x} = -\pi \cot \pi x.$$

$$6. n \sum \frac{1}{x} = \left\{ \sum \frac{1}{x/n} + \sum \frac{1}{x/n} + \dots + \sum \frac{1}{x/n} \right\}$$

$$= n \log n.$$

$$\text{Cor. 1. } \sum \frac{1}{x-\frac{1}{2}} = C_0 + \log x + (1-\frac{1}{2}) \frac{B_2}{2x^2} - (1-\frac{1}{2}) \frac{B_4}{4x^4} + \dots$$

2.  $\sum_{n=1}^{\infty} \frac{1}{n} + \sum_{n=1}^{\infty} \frac{1}{2n} + \dots + \sum_{n=1}^{\infty} \frac{1}{mn} = -n \log n$

3. i.  $\phi(\frac{1}{2}) = -2 \log 2$ ; ii.  $\phi(\frac{1}{3}) = -\frac{3}{2} \log 3 - \frac{\pi}{2\sqrt{3}}$

iii.  $\phi(\frac{1}{4}) = -\frac{\pi}{2} - 3 \log 2$ ; iv.  $\phi(\frac{1}{8}) = -\frac{\pi}{2}\sqrt{3} - 2 \log 2 - \frac{3}{2} \log 3$

v.  $3\phi(\frac{1}{2}) - 2\phi(\frac{1}{3}) = \pi$ .

4.  $\phi(\frac{1}{2n}) + \phi(\frac{1}{2^2n}) + \dots + \phi(\frac{1}{2^m n}) = -n \log \frac{1}{2^m}$

7.  $\frac{1}{a+b} + \frac{1}{a+2b} + \frac{1}{a+3b} + \dots + \frac{1}{a+xb} = \frac{1}{b} \{ \phi(\frac{a}{b}) - \phi(\frac{a}{xb}) \}$

8.  $\frac{1}{a+b} - \frac{1}{a+2b} + \frac{1}{a+3b} - \dots = \frac{1}{2b} \{ \phi(\frac{a}{2b}) - \phi(\frac{a}{4b}) \}$

9.  $\phi(\frac{x}{1+x}) = \phi(\frac{x}{2}) - \log 2 + x \int_0^1 \frac{x^x}{1+x^x} dx$

10.  $\phi(\frac{1}{2}) = -x \int_0^1 \frac{(1-x)^2}{x(x^2-1)} dx$

11.  $\phi(\frac{1}{2}-1) + \phi(\frac{1}{2}) = -x \{ 1 + \frac{2}{x^2-x} + \frac{2}{(2x)^2-2x} + \dots \}$

12.  $\frac{2}{x^2-x} + \frac{2}{(2x)^2-2x} + \frac{2}{(3x)^2-3x} + \dots = \int_0^1 \frac{x^{2-2(1-x)^n}}{1-x^2} dx$

13.  $1 + \frac{2}{(2x)^2-2x} + \frac{2}{(4x)^2-4x} + \frac{2}{(6x)^2-6x} + \dots$

$= \frac{1}{2} \{ 1 + \frac{2}{x^2-x} + \frac{2}{(2x)^2-2x} + \dots \} + \frac{\log 2}{x}$

+ Log part of  $(1 - \frac{1}{1+x} + \frac{1}{1+2x} - \frac{1}{1+3x} + \dots)$

N.B. i.  $x - \frac{x^{1+n}}{1+n} + \frac{x^{1+2n}}{1+2n} - \dots = \int \frac{x dx}{1+x^n}$

ii.  $x + \frac{x^{1+n}}{1+n} + \frac{x^{1+2n}}{1+2n} + \dots = \int_0^x \frac{dx}{1-x^n}$

iii. If n is odd  $\int_0^x \frac{dx}{1-x^n} = \int_0^x \frac{1}{1+(-x)^n} dx$

iv. If n is even  $\int_0^x \frac{dx}{1-x^n} = \frac{1}{2} \int_0^x \frac{dx}{1+x^2} + \frac{1}{2} \int_0^x \frac{dx}{1-x^2}$

v.  $\int \frac{1}{x^l} < n+1$

(a)  $\int \frac{x^{l-1}}{x^n-1} dx = \frac{1}{n} \log(x-1) +$

$$\frac{(-1)^l}{n} \log(x+1) + \frac{1}{n} \sum \cos \frac{r l \pi}{n} \log(x^2 - 2x \cos \frac{r \pi}{n} + 1)$$

$$- \frac{2}{n} \sum \sin \frac{r l \pi}{n} \tan^{-1} \frac{x - \cos \frac{r \pi}{n}}{\sin \frac{r \pi}{n}}$$

$r = 2, 4, 6, \dots$  up to  $n-2$ .

(b)  $\int \frac{x^{l-1}}{x^n+1} = \frac{(-1)^{l-1}}{n} \log(x+1)$   $n$  being odd.

$$- \frac{1}{n} \sum \cos \frac{r l \pi}{n} \log(x^2 - 2x \cos \frac{r \pi}{n} + 1)$$

$$+ \frac{2}{n} \sum \sin \frac{r l \pi}{n} \tan^{-1} \frac{x - \cos \frac{r \pi}{n}}{\sin \frac{r \pi}{n}}$$

$r = 1, 3, 5, \dots$  up to  $n-2$ .

vi.  $\int \frac{1}{x^{n+1}} < n$  even

(a)  $\int \frac{x^{l-1}}{x^n-1} dx = \frac{1}{n} \log(x-1) + \frac{1}{n} \sum \cos \frac{r l \pi}{n} x$

$$\log(x^2 - 2x \cos \frac{r \pi}{n} + 1) - \frac{2}{n} \sum \sin \frac{r l \pi}{n} x$$

$$\tan^{-1} \frac{x - \cos \frac{r \pi}{n}}{\sin \frac{r \pi}{n}}, \quad r = 2, 4, 6, \dots, (n-1).$$

(b)  $\int \frac{x^{l-1}}{x^n+1} dx = -\frac{1}{n} \sum \cos \frac{r l \pi}{n} \log(x^2 - 2x \cos \frac{r \pi}{n} + 1)$

$$+ \frac{2}{n} \sum \sin \frac{r l \pi}{n} \tan^{-1} \frac{x - \cos \frac{r \pi}{n}}{\sin \frac{r \pi}{n}} \quad n \text{ being even}$$

$r = 1, 3, 5, \dots, (n-1).$



14. If  $A_n = \int_0^x \frac{dx}{1+x^n}$ , then

i.  $A_1 = \log(1+x)$ ; ii.  $A_2 = \log^{-1} x$ .

iii.  $A_3 = \frac{1}{6} \log \frac{(1+x)^3}{1+x^3} + \frac{1}{\sqrt{3}} \tan^{-1} \frac{x\sqrt{3}}{2-x}$ .

iv.  $A_4 = \frac{1}{2\sqrt{2}} \log \frac{1+x\sqrt{2}+x^2}{1-x\sqrt{2}+x^2} + \frac{1}{2\sqrt{2}} \tan^{-1} \frac{x\sqrt{2}}{1-x^2}$ .

v.  $A_5 = \frac{1}{20} \log \frac{(1+x)^5}{1+x^5} + \frac{1}{4\sqrt{5}} \log \frac{1+x \cdot \frac{\sqrt{5}-1}{2} + x^2}{1-x \cdot \frac{\sqrt{5}-1}{2} + x^2}$

+  $\frac{1}{10} \sqrt{10-2\sqrt{5}} \tan^{-1} \frac{x \sqrt{10-2\sqrt{5}}}{4-x(\sqrt{5}+1)} + \frac{\sqrt{10+2\sqrt{5}}}{10} \tan^{-1} \frac{x \sqrt{10+2\sqrt{5}}}{4+x(\sqrt{5}-1)}$

vi.  $A_6 = \frac{1}{2} \tan^{-1} x + \frac{1}{2} \tan^{-1} x^3 + \frac{1}{4\sqrt{3}} \log \frac{1+x\sqrt{3}+x^2}{1-x\sqrt{3}+x^2}$

vii.  $A_8 = \frac{\sqrt{2+\sqrt{2}}}{16} \left\{ \log \frac{1+x\sqrt{2+\sqrt{2}}+x^2}{1-x\sqrt{2+\sqrt{2}}+x^2} + 2 \tan^{-1} \frac{x\sqrt{2+\sqrt{2}}}{1-x^2} \right\}$

+  $\frac{\sqrt{2-\sqrt{2}}}{16} \left\{ \log \frac{1+x\sqrt{2-\sqrt{2}}+x^2}{1-x\sqrt{2-\sqrt{2}}+x^2} + 2 \tan^{-1} \frac{x\sqrt{2-\sqrt{2}}}{1-x^2} \right\}$

viii.  $A_{10} = \frac{1}{4} \tan^{-1} x - \frac{1}{20} \tan^{-1} x^5 + \frac{1}{4\sqrt{5}} \tan^{-1} \frac{(x-1)\sqrt{5}}{1-3x^2+x^4}$

+  $\frac{1}{40} \sqrt{10-2\sqrt{5}} \log \frac{1+\frac{x}{2} \sqrt{10-2\sqrt{5}}+x^2}{1-\frac{x}{2} \sqrt{10-2\sqrt{5}}+x^2}$

+  $\frac{1}{40} \sqrt{10+2\sqrt{5}} \log \frac{1+\frac{x}{2} \sqrt{10+2\sqrt{5}}+x^2}{1-\frac{x}{2} \sqrt{10+2\sqrt{5}}+x^2}$ .

Ex. 1. i.  $\frac{1}{1.2} - \frac{1}{2.2^4} + \frac{1}{7.2^7} - \dots = \frac{\pi}{6\sqrt{3}} + \frac{1}{4} \log 3$ .

ii.  $\frac{\sqrt{3}-1}{1} - \frac{(\sqrt{3}-1)^2}{2} + \frac{(\sqrt{3}-1)^3}{7} - \dots = \frac{\pi}{4\sqrt{3}} + \frac{1}{3} \log \frac{1+\sqrt{3}}{\sqrt{2}}$ .

iii.  $\frac{2-\sqrt{3}}{1} - \frac{(2-\sqrt{3})^2}{5} + \frac{(2-\sqrt{3})^3}{9} - \dots = \frac{\pi}{16} (\sqrt{3}-1)$

-  $\frac{\sqrt{3}-1}{20} \log(\sqrt{3}-1)$ .

2. If  $A_n = 1 + \frac{2}{2^2 - n} + \frac{2}{(2n)^2 - 2n} + \frac{2}{(3n)^2 - 3n} + \dots$ , then

$A_2 = 2 \log_e 2$ ;  $A_3 = \log_e 3$ ;  $A_4 = \frac{3}{2} \log_e 2$ ;  $A_6 = \frac{1}{2} \log_e 3 + \frac{1}{3} \log_e 4$

$A_5 = \frac{1}{2} \log_e 5 + \frac{1}{\sqrt{5}} \log_e \frac{\sqrt{5}+1}{2}$ ;  $A_8 = \log_e 2 + \frac{1}{2\sqrt{2}} \log_e (1+\sqrt{2})$

$A_{10} = \frac{2}{5} \log_e 2 + \frac{1}{4} \log_e 5 + \frac{3}{2\sqrt{5}} \log_e \frac{1+\sqrt{5}}{2}$ ;  $A_{12} = \frac{1}{2} \log_e 2 + \frac{1}{4} \log_e 3$

$- \frac{1}{\sqrt{3}} \log_e (\sqrt{3}-1)$ ;  $A_{16} = \frac{5}{8} \log_e 2 + \frac{1}{4\sqrt{2}} \log_e (1+\sqrt{2})$

$+ \frac{\sqrt{2}+\sqrt{2}}{16} \log_e \frac{2+\sqrt{2}+\sqrt{2}}{2-\sqrt{2}+\sqrt{2}}$   $+ \frac{\sqrt{2}-\sqrt{2}}{16} \log_e \frac{2+\sqrt{2}-\sqrt{2}}{2-\sqrt{2}-\sqrt{2}}$

$A_{20} = \frac{1}{8} \log_e 5 + \frac{3}{10} \log_e 2 + \frac{3}{4\sqrt{5}} \log_e \frac{\sqrt{5}+1}{2}$

$+ \frac{\sqrt{10}-2\sqrt{5}}{40} \log_e \frac{4+\sqrt{10}-2\sqrt{5}}{2-\sqrt{10}-2\sqrt{5}}$   $+ \frac{\sqrt{10}+2\sqrt{5}}{40} \log_e \frac{4+\sqrt{10}+2\sqrt{5}}{4-\sqrt{10}+2\sqrt{5}}$

15. If  $\epsilon \frac{1}{x} = C_0 + \log_e a$ , then

$$\left(\frac{x+\frac{1}{2}}{a}\right)^{4x} = 1 - \frac{x}{11} \cdot \frac{1}{6a^2} + \frac{x(x+\frac{1}{10})}{12} \cdot \frac{1}{(6a^2)^2} -$$

$$\frac{x(x^2 + 3\frac{3}{10}x + 12\frac{5}{70})}{18(6a^2)^3} + \dots$$

Cor.  $Lx$  is minimum when  $x = \frac{6}{13}$  very nearly.

Sol.  $Lx$  is minimum when  $\epsilon \frac{1}{x} = C_0$  i.e.  $a=1$

$\therefore x = \frac{1}{2} - \frac{1}{24} + \dots$  or  $x = \frac{1}{2}$  very nearly.

16.  $C_0 = \log_e 2 - 1\left(\frac{2}{3^2-3}\right) - 2\left(\frac{2}{6^2-6} + \frac{2}{9^2-9} + \frac{1}{12^2-12}\right) - \dots$

the last term in the  $n$ th group =  $\frac{2}{\left(\frac{3^n+3}{2}\right)^2 - \frac{3^n+3}{2}}$

$$17. i. \frac{\log 1}{1} + \frac{\log 2}{2} + \frac{\log 3}{3} + \dots + \frac{\log x}{x} = \phi(x)$$

$$\phi(x) = (\epsilon \frac{1}{x} - c_0) \log x - \frac{1}{2} (\log x)^2 + c_1 + \frac{B_2}{2x^2} \cdot 1$$

$$- \frac{B_4}{4x^4} (1 + \frac{1}{2} + \frac{1}{3}) + \frac{B_6}{6x^6} (1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5}) - \dots$$

where  $c_1 = -0.72815845483680$

Sol. Write  $n-1$  for  $x$  in VIII 1, then divide both sides by  $x^2$  and find the coeff<sup>ts</sup> of  $n$  from both sides and equate them

Cor. When  $x = \infty$ ,  $\phi(x) - \frac{1}{2} (\epsilon \frac{1}{x} - c_0)^2 = c_1$

ii.  $\phi(x) = \frac{\log 1}{1} - \frac{\log(1+x)}{1+x} + \frac{\log 2}{2} - \frac{\log(2+x)}{2+x} + \dots$

Cor.  $\frac{\log 1}{1} - \frac{\log 3}{3} + \frac{\log 5}{5} - \dots = \frac{\pi}{2} \log 2 + \frac{1}{2} \{ \phi(\frac{1}{2}) - \phi(\frac{3}{2}) \}$

iii.  $n \phi(x) - \{ \phi(\frac{x}{n}) + \phi(\frac{x-1}{n}) + \dots + \phi(\frac{x-n+1}{n}) \}$

$$= n \log n (\epsilon \frac{1}{x} - c_0) - \frac{n}{2} (\log n)^2$$

Cor.  $\phi(\frac{1}{n}) + \phi(\frac{2}{n}) + \dots + \phi(\frac{n-1}{n})$

$$= n c_0 \log n + \frac{n}{2} (\log n)^2$$

Ex. 1.  $\frac{\sqrt{1}}{\sqrt{2}} \cdot \frac{\sqrt{3}}{\sqrt{4}} \cdot \frac{\sqrt{5}}{\sqrt{6}} \cdot \frac{\sqrt{7}}{\sqrt{8}} \cdot \frac{\sqrt{9}}{\sqrt{10}} \dots$  ad inf =  $2^{\frac{1}{2} \log 2 - c_0}$

2.  $\phi(\frac{1}{2}) = (\log 2)^2 + 2 c_0 \log 2$

3.  $\phi(\frac{1}{3}) + \phi(\frac{2}{3}) = \frac{3}{2} (\log 3)^2 + 3 c_0 \log 3$

4.  $\phi(\frac{1}{4}) + \phi(\frac{3}{4}) = 7 (\log 2)^2 + 6 c_0 \log 2$

5.  $\phi(\frac{1}{8}) + \phi(\frac{7}{8}) = c_0 (3 \log 3 + 4 \log 2) + \frac{3}{2} (\log 12)^2 - (\log 4)^2$

iv. When  $x$  lies between 0 & 1

$$\frac{\pi}{2} \left\{ \log \frac{1-x}{1-x} + (C_0 + \log 2\pi)(1-x) \right\}$$

$$= \frac{\log 1}{1} \sin 2\pi x + \frac{\log 2}{2} \sin 4\pi x + \frac{\log 3}{3} \sin 6\pi x + \dots$$

N.B.  $\frac{\pi}{2} - \pi x = \sin 2\pi x + \frac{1}{2} \sin 4\pi x + \frac{1}{3} \sin 6\pi x + \dots$

v.  $\phi(x-1) - \phi(-x) = (C_0 + \log 2\pi) \pi \cot \pi x$  (for the same limits)  $+ 2\pi \left\{ \sin 2\pi x \log 1 + \sin 4\pi x \log 2 + \dots \right\}$

N.B.  $\sin 2\pi x + \sin 4\pi x + \sin 6\pi x + \dots = \frac{1}{2} \cot \pi x$ .

Ex. 1. Find  $\phi\left(\frac{1}{2}\right)$ ,  $\phi\left(\frac{2}{3}\right)$ ,  $\phi\left(\frac{3}{4}\right)$  and  $\phi\left(\frac{5}{8}\right)$ .

2.  $\frac{\log 1}{1} - \frac{\log 3}{3} + \frac{\log 5}{5} - \dots = \frac{\pi}{2} \log \pi - \pi \log \sqrt{\frac{1}{2}} - \frac{\pi}{4} C_0$

3.  $\frac{\left(\frac{\sqrt{1}}{\sqrt{2}} \cdot \frac{\sqrt{3}}{\sqrt{4}} \cdot \frac{\sqrt{5}}{\sqrt{6}} \cdot \frac{\sqrt{7}}{\sqrt{8}} \dots\right)^{\log 2}}{\left(\frac{\sqrt{1}}{\sqrt{3}} \cdot \frac{\sqrt{5}}{\sqrt{7}} \cdot \frac{\sqrt{9}}{\sqrt{11}} \cdot \frac{\sqrt{13}}{\sqrt{15}} \dots\right)^{\frac{4}{\pi}}} = \frac{\sqrt{2}}{\pi} \left(1 - \frac{1}{2}\right)^4$

18.  $(\log 1)^2 + (\log 2)^2 + (\log 3)^2 + \dots + (\log x)^2 = \phi(x)$

i.  $\phi(x) = 2 \log x \log \frac{1-x}{\sqrt{2\pi}} - (x + \frac{1}{2})(\log x)^2 + 2x + \frac{1}{2} C_1^2$   
 $+ C_1 - \frac{\pi^2}{24} - \frac{1}{2} (\log 2\pi)^2 + 2 \left\{ \frac{B_4}{3 \cdot 4} \cdot \frac{1 + \frac{1}{2}}{x^3} - \frac{B_6}{5 \cdot 6} \cdot \frac{1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}}{x^5} + \dots \right\}$

Sol. Equate the Coeff<sup>s</sup> of  $x^2$  in VIII 1.

ii.  $\phi(x) - \left\{ \phi\left(\frac{x}{n}\right) + \phi\left(\frac{x-1}{n}\right) + \dots + \phi\left(\frac{x-n+1}{n}\right) \right\} =$

$$2 \log n \log \frac{x}{\sqrt{2\pi}} - x(\log n)^2 - (n-1) \left\{ \frac{1}{2} C_0^2 + C_1 - \frac{\pi^2}{24} - \frac{1}{2} (\log 2\pi)^2 \right\} \\ - \frac{1}{2} (\log n)^2. \text{ If } C \text{ be the constant in this series then}$$

$$\text{Cor. } \phi\left(-\frac{1}{2n}\right) + \phi\left(-\frac{2}{2n}\right) + \phi\left(-\frac{3}{2n}\right) + \dots + \phi\left(-\frac{n-1}{2n}\right) \\ = \log n \log 2\pi + (n-1)C + \frac{1}{2} (\log n)^2.$$

Ex. 1. If  $x$  becomes infinite then

$$\frac{1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{x}}{\frac{1}{\log 1} + \frac{1}{\log 2} + \frac{1}{\log 3} + \dots + \frac{1}{\log x}} \cdot x^{x \log x - 2x} \\ \times e^{2x + \frac{1}{2} (\frac{1}{2} - \log x)^2} = e^{\frac{\pi^2}{24}} (2\pi)^{\frac{1}{2} \log 2\pi}$$

2. Find  $\phi\left(\frac{1}{2}\right)$ ,  $\phi\left(\frac{1}{3}\right) + \phi\left(\frac{2}{3}\right)$ ,  $\phi\left(\frac{1}{4}\right) + \phi\left(\frac{3}{4}\right)$  and  $\phi\left(\frac{1}{8}\right) + \phi\left(\frac{7}{8}\right)$ .

$$\text{iii } \frac{\phi(x-1) + \phi(-x)}{2} = C_1 - \frac{\pi^2}{24} + \frac{1}{2} (C_0 + \log 2\pi) (C_0 - \log \frac{\pi}{2 \sin \pi x}) \\ - \left\{ \frac{\log 1}{1} \cos 2\pi x + \frac{\log 2}{2} \cos 4\pi x + \dots + \frac{\log x}{x} \cos 2\pi x \right\}$$

19. If  $C_n$  be the constant in  $(\log 1)^n + (\log 2)^n + \dots + (\log x)^n$  and if  $\phi_n(x) = (\log 1)^n + (\log 2)^n + \dots + (\log x)^n - C_n$  then

$$\text{i The logarithmic part of } \phi_n(x) = n \log x \phi_{n-1}(x) \\ - \frac{n(n-1)}{2} (\log x)^2 \phi_{n-2}(x) + \frac{n(n-1)(n-2)}{6} (\log x)^3 \phi_{n-3}(x) - \dots$$

and the non-logarithmic part can be found from VII 1.

$$\text{ii } \phi_0(x) (\log x)^n - \frac{n}{1} \phi_1(x) (\log x)^{n-1} + \frac{n(n-1)}{2} \phi_2(x) (\log x)^{n-2} - \dots$$

$$\begin{aligned}
 &= x^{\frac{1}{2}} - \frac{1}{x^{\frac{3}{2}}} \frac{B_{n+1}}{n+1} \sin \frac{\pi x}{2} - \frac{\pi}{2} \cdot \frac{1}{x^{\frac{5}{2}}} \frac{B_{n+2}}{n+2} \cos \frac{\pi x}{2} \\
 &+ \frac{n(n+\frac{5}{3})}{2 \cdot 4} \cdot \frac{1}{x^{\frac{7}{2}}} \frac{B_{n+3}}{n+3} \sin \frac{\pi x}{2} + \frac{n(n+2)(n+3)}{2 \cdot 4 \cdot 6} \cdot \frac{1}{x^{\frac{9}{2}}} \\
 &\times \frac{B_{n+4}}{n+4} \cos \frac{\pi x}{2} - \frac{\pi(n+2)(n+4)^2 + \frac{n(n+2)}{3} + \frac{4n}{5}}{2 \cdot 4 \cdot 6 \cdot 8} \cdot \frac{1}{x^{\frac{11}{2}}} \\
 &\times \frac{B_{n+5}}{n+5} \sin \frac{\pi x}{2} - \frac{\pi(n+4)(n+5) \{ (n+2)(n+4) + \frac{2}{3}(n+1) \}}{2 \cdot 4 \cdot 6 \cdot 8 \cdot 10} \\
 &\times \frac{1}{x^{\frac{13}{2}}} \frac{B_{n+6}}{n+6} \cos \frac{\pi x}{2} + \dots
 \end{aligned}$$

iii.  $\phi_n(\frac{x}{n}) + \phi_n(\frac{x-1}{n}) + \dots + \phi_n(\frac{x-n+1}{n})$   
 $= \phi_n(x) - n \log n \phi_{n-1}(x) + \frac{n(n-1)}{1^2} (\log n)^2 \phi_{n-2}(x) - \dots$

Case 1.  $\phi_n(\frac{1}{n}) + \phi_n(\frac{2}{n}) + \dots + \phi_n(\frac{n-1}{n})$   
 $= - \{ c_n - n \log n c_{n-1} + \frac{n(n-1)}{1^2} (\log n)^2 c_{n-2} - \dots \}$

Case 2. There will be no logarithmic functions in  $\phi_n(\frac{x}{n}) + \phi_n(\frac{x-1}{n}) + \dots + \phi_n(\frac{x-n+1}{n})$ .

20. Let  $1^x + 2^x k + 3^x k^2 + \dots + x^x k^{x-1} = k^x \phi(x) = F_k(x)$

i.  $\phi(x) = C_n(k) + x^n \frac{\psi_0(k)}{k-1} - \frac{n}{1} \cdot x^{n-1} \frac{\psi_1(k)}{(k-1)^2} + \frac{n(n-1)}{1^2} \cdot \frac{\psi_2(k)}{(k-1)^3} - \dots$  where  $\psi$  is the same  $\psi$  in

ii.  $C_n(k) = \frac{\psi_n(k)}{(1-k)^{n+1}}$  and  $k \psi_n(k) = k^2 \psi(\frac{1}{k})$

iii.  $F_k(\frac{x}{n}) + F_k(\frac{x-1}{n}) + F_k(\frac{x-2}{n}) + \dots + F_k(\frac{x-n+1}{n}) - n C_n(k)$   
 $= \frac{\sqrt[n]{k}}{k^{n^2}} \{ F_{\sqrt[n]{k}}(x) - C_n(\sqrt[n]{k}) \}$

$$\text{Cor. } F_n\left(\frac{1}{n}\right) + F_n\left(\frac{2}{n}\right) + \dots + F_n\left(\frac{n-1}{n}\right) = n C_n'(k) - \frac{\gamma_k C_n'(k)}{k+2}$$

21. Let  $\frac{\log 1}{1^n} + \frac{\log 2}{2^n} + \frac{\log 3}{3^n} + \dots + \frac{\log x}{x^n} = \phi_n(x)$  and let  $C_n'$  be the constant. Then,

$$\begin{aligned} \text{i. } \phi_n(x) &= C_n' - \left\{ \frac{1}{(x+1)^n} + \frac{1}{(x+2)^n} + \frac{1}{(x+3)^n} + \dots \right\} \log x - \frac{1}{(x-1)^n x^n} \\ &+ B_2 \frac{n}{1!} \cdot \frac{1}{n x^{n+1}} - B_4 \frac{n(n+1)(n+2)}{4!} \left( \frac{1}{x} + \frac{1}{x+1} + \frac{1}{x+2} \right) \frac{1}{x^{n+3}} \\ &+ B_6 \frac{n(n+1)(n+2)(n+3)(n+4)}{6!} \left( \frac{1}{x} + \frac{1}{x+1} + \frac{1}{x+2} + \frac{1}{x+3} + \frac{1}{x+4} \right) \\ &\times \frac{1}{x^{n+5}} - \dots \end{aligned}$$

$$\begin{aligned} \text{ii. } \phi_n(x) &= n x C_{n+1}' - \frac{n(n+1)}{1!} x^2 C_{n+2}' + \frac{n(n+1)(n+2)}{2!} x^3 C_{n+3}' \\ &- \dots - n \cdot \frac{1}{n} x S_{n+1} + \frac{n(n+1)}{2!} \left( \frac{1}{n} + \frac{1}{n+1} \right) x^2 S_{n+2} - \dots \end{aligned}$$

$$\begin{aligned} \text{iii. } x^n \phi_n(x) &= \left\{ \phi_n\left(\frac{x}{n}\right) + \phi_n\left(\frac{x-1}{n}\right) + \dots + \phi_n\left(\frac{x-n+1}{n}\right) \right\} \\ &= C_n' (n^n - n) - n^n \log n \left\{ \frac{1}{(x+1)^n} + \frac{1}{(x+2)^n} + \frac{1}{(x+3)^n} + \dots \right\} \end{aligned}$$

$$\begin{aligned} \text{Cor. } \phi_n\left(\frac{1}{n}\right) + \phi_n\left(\frac{2}{n}\right) + \phi_n\left(\frac{3}{n}\right) + \dots + \phi_n\left(\frac{n-1}{n}\right) \\ = n^n \log n S_n - (n^n - n) C_n'. \end{aligned}$$

22. Let  $(\log 1)^e + \frac{1}{2}(\log 2)^e + \frac{1}{3}(\log 3)^e + \dots$  to  $x$  terms  $= \psi_n(x)$  and let  $C_n$  be its constant; then

$$\text{i. } \psi_n(x) - \frac{1}{n+1} (\log x)^{n+1} = C_n \text{ when } x \rightarrow \infty$$

$$\begin{aligned} \text{ii. } n \psi_n(x) &= \left\{ \psi_n\left(\frac{x}{n}\right) + \psi_n\left(\frac{x-1}{n}\right) + \psi_n\left(\frac{x-2}{n}\right) + \dots + \psi_n\left(\frac{x-n+1}{n}\right) \right\} \\ &= \frac{\pi}{n+1} (\log x)^{n+1} \cos \pi n + n \log n \left\{ \psi_n(x) - C_{n-1} \right\} \end{aligned}$$

$$- \frac{n(n-1)}{12} m(\log n)^2 \{ \phi_{n-1}(x) - C_{n-2} \} + \&c \text{ the last term being}$$

$$(-1)^{n-1} m(\log m)^2 \{ \phi_0(m) - C_0 \}$$

$$23. \frac{(\log 1)^2}{1^{n+1}} + \frac{(\log 2)^2}{2^{n+1}} + \frac{(\log 3)^2}{3^{n+1}} + \&c$$

$$= \frac{1/2}{n^{n+1}} + C_n - \frac{n}{1} C_{n+1} + \frac{n^2}{2} C_{n+2} - \frac{n^3}{6} C_{n+3} + \&c$$

Sol. Differentiate both sides n times in

$$\text{Ex. 1. } \frac{(\log 1)^3}{1\sqrt{1}} + \frac{(\log 2)^3}{2\sqrt{2}} + \frac{(\log 3)^3}{3\sqrt{3}} + \&c = 96.001 \text{ nearly}$$

$$2. \frac{\log 1}{1^2} + \frac{\log 2}{2^2} + \frac{\log 3}{3^2} + \&c = .9382 \text{ nearly}$$

$$3. \frac{(\log 1)^4}{1^2} + \frac{(\log 2)^4}{2^2} + \frac{(\log 3)^4}{3^2} + \&c = 24 \text{ nearly.}$$

$$4. \frac{(\log 1)^5}{1\sqrt{1}} + \frac{(\log 2)^5}{2\sqrt{2}} + \frac{(\log 3)^5}{3\sqrt{3}} + \&c = 7680 \text{ nearly.}$$

$$5. \frac{(\log 1)^5}{1^2} \sqrt{\log 1} + \frac{(\log 2)^5}{2^2} \sqrt{\log 2} + \&c = 288 \text{ nearly.}$$

$$24. \frac{\log 1}{\sqrt{1}} + \frac{\log 2}{\sqrt{2}} + \frac{\log 3}{\sqrt{3}} + \dots + \frac{\log x}{\sqrt{x}} = \phi(x)$$

$$i. \phi(x) = \frac{\log 1}{\sqrt{1}} - \frac{\log(1+x)}{\sqrt{1+x}} + \frac{\log 2}{\sqrt{2}} - \frac{\log(2+x)}{\sqrt{2+x}} + \&c$$

$$ii. \phi(x) = \left( \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{x}} \right) \log x$$

$$+ (\sqrt{2+1}) \left( \frac{1}{\sqrt{1}} - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} - \&c \right) \left( \log x + \frac{1}{2} C_0 + \frac{\pi}{4} + \frac{1}{2} \log \pi \right)$$

$$- 4\sqrt{x} + \frac{1}{2} \cdot \frac{B_2}{x\sqrt{x}} - \frac{1.3.5}{2.4.6} \left( 1 + \frac{1}{3} + \frac{1}{5} \right) \frac{B_4}{2x^2\sqrt{x}}$$

$$+ \frac{1.3.5.7.9}{2.4.6.8.10} \left( 1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{9} \right) \frac{B_6}{3^2 x^3 \sqrt{x}} - \&c.$$



$$\text{iii. } \phi(x) = \frac{1}{\sqrt{x}} \left\{ \phi\left(\frac{x}{2}\right) + \phi\left(\frac{x-1}{2}\right) + \dots + \phi\left(\frac{x-n+1}{2}\right) \right\}$$

$$= \left( \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{x}} \right) \log x$$

$$- (1 + \sqrt{2}) \left( \frac{1}{\sqrt{1}} - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} - 2c \right) \left\{ (\sqrt{n}-1) \left( \frac{1}{2} C_0 + \frac{\pi}{2} + \log \sqrt{8\pi} \right) - \log n \right\}$$

$$\text{iv. } \text{If } \psi(x) = \frac{1}{\sqrt{x}} + \frac{1}{\sqrt{2}} + \dots + \frac{1}{\sqrt{x}}, \text{ then}$$

$$\left\{ \phi(x-1) + \phi(x) - 2c \right\} + (C_0 + \frac{\pi}{2} + \log 8\pi) \left\{ \psi(x-1) + \psi(x) - 2c \right\}$$

$$= 2 \left\{ \frac{\log 1}{\sqrt{1}} \cos 2\pi x + \frac{\log 2}{\sqrt{2}} \cos 4\pi x + 2c \right\}$$

$$\text{v. } \left\{ \phi(x-i) - \phi(x) \right\} + (C_0 - \frac{\pi}{2} + \log 8\pi) \left\{ \psi(x-1) - \psi(x) \right\}$$

$$= 2 \left\{ \frac{\log 1}{\sqrt{1}} \sin 2\pi x + \frac{\log 2}{\sqrt{2}} \sin 4\pi x + 2c \right\}$$

In both cases  $c$  &  $c'$  are the constants of  $\phi(x)$  and  $\psi(x)$  respectively.

Ex. 1. Find the values of  $\phi\left(\frac{1}{2}\right)$ ,  $\phi\left(\frac{2}{3}\right)$ , &  $\phi\left(\frac{3}{4}\right)$ .

2. Show that the constant in  $\phi(x)$

$$= -\frac{1}{2} \delta_{\frac{1}{2}} (C_0 + \frac{\pi}{2} + \log 8\pi) = 3.92265$$

$$= 2 \left\{ 2 - \frac{1}{2} \cdot \frac{B_{\frac{1}{2}}}{2} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \left( 1 + \frac{1}{3} + \frac{1}{5} \right) \frac{B_{\frac{1}{2}}}{2} - 2c \right\}$$

Sol. Write  $\frac{1+h}{2}$  for  $n$  in VIII 4, and equate the coeffs. of  $h$ . Put  $x=1$  in VIII 24. ii; then the second result is at once obtained.

1. If  $S_n = \frac{1}{(1-a)^n} - \frac{1}{(1+a)^n} + \frac{1}{(3-a)^n} - \frac{1}{(3+a)^n} + \dots$  then

i. If  $n$  is odd,

$$\frac{\cos(1-a)x}{(1-a)^n} - \frac{\cos(1+a)x}{(1+a)^n} + \frac{\cos(3-a)x}{(3-a)^n} - \frac{\cos(3+a)x}{(3+a)^n} + \dots$$

$$= S_n - \frac{x^2}{1!} S_{n-2} + \frac{x^4}{2!} S_{n-4} - \dots$$

as far as the term containing  $S_1$ ,

ii. If  $n$  is even

$$\frac{\sin(1-a)x}{(1-a)^n} - \frac{\sin(1+a)x}{(1+a)^n} + \frac{\sin(3-a)x}{(3-a)^n} - \frac{\sin(3+a)x}{(3+a)^n} + \dots$$

$$= \frac{x}{1!} S_{n-1} - \frac{x^3}{3!} S_{n-3} + \frac{x^5}{5!} S_{n-5} - \dots$$

as far as the term containing  $S_1$ ,

2. If  $S_n = \frac{1}{(1-a)^n} + \frac{1}{(1+a)^n} + \frac{1}{(3-a)^n} + \frac{1}{(3+a)^n} + \dots$  then

i. If  $n$  is even

$$\frac{\cos(1-a)x}{(1-a)^n} + \frac{\cos(1+a)x}{(1+a)^n} + \frac{\cos(3-a)x}{(3-a)^n} + \frac{\cos(3+a)x}{(3+a)^n} + \dots$$

$$= S_n - \frac{x^2}{1!} S_{n-2} + \frac{x^4}{2!} S_{n-4} - \dots$$

as far as the term containing  $S_2$ ,

ii. If  $n$  is odd

$$\frac{\sin(1-a)x}{(1-a)^n} + \frac{\sin(1+a)x}{(1+a)^n} + \frac{\sin(3-a)x}{(3-a)^n} + \frac{\sin(3+a)x}{(3+a)^n} + \dots$$

$$= \frac{x}{1!} S_{n-1} - \frac{x^3}{3!} S_{n-3} + \frac{x^5}{5!} S_{n-5} - \dots$$

as far as the

term containing  $S_2$

Sol. In both 1 & 2 expand the series in ascending powers of  $x$  and apply

$$3. \text{ If } \phi(x) = \frac{\cos x}{1^n} - (1+\frac{1}{2}) \frac{\cos 3x}{3^n} + (1+\frac{1}{2}+\frac{1}{3}) \frac{\cos 5x}{5^n} - \dots$$

then if  $n$  is odd  $\phi(n-2) - \phi(n) =$

$$x \left\{ \left( \frac{\sin x}{1^{n-2}} - \frac{\sin 3x}{3^{n-2}} + \frac{\sin 5x}{5^{n-2}} - \dots \right) \right.$$

$$\left. - \left( \frac{\sin x}{1^n} - \frac{\sin 3x}{3^n} + \frac{\sin 5x}{5^n} - \dots \right) \right\}$$

$$+ n \left\{ \left( \frac{\cos 3x}{1^{n-1}} - \frac{\cos 3x}{3^{n-1}} + \frac{\cos 5x}{5^{n-1}} - \dots \right) \right.$$

$$\left. - \left( \frac{\cos x}{1^{n+1}} - \frac{\cos 3x}{3^{n+1}} + \frac{\cos 5x}{5^{n+1}} - \dots \right) \right\}$$

$$4. \text{ Let } \psi(n) = \left\{ \frac{\sin x}{1^n} - \frac{1}{2} \cdot \frac{\sin 3x}{3^n} + \frac{1.3}{2.4} \cdot \frac{\sin 5x}{5^n} - \dots \right\}$$

$$- \cos \pi n \left\{ \left( \frac{\sin 2x}{2^n} - \frac{1}{2} \cdot \frac{\sin 4x}{4^n} + \frac{1.3}{2.4} \cdot \frac{\sin 6x}{6^n} - \dots \right) \right.$$

$$\left. - \left( \frac{\sin 2x}{2^{n+1}} - \frac{1}{2} \cdot \frac{\sin 4x}{4^{n+1}} + \frac{1.3}{2.4} \cdot \frac{\sin 6x}{6^{n+1}} - \dots \right) \right\} \text{ and}$$

$$\psi(n) = \left\{ \frac{\cos x}{1^n} - \frac{1}{2} \cdot \frac{\cos 3x}{3^n} + \frac{1.3}{2.4} \cdot \frac{\cos 5x}{5^n} - \dots \right\}$$

$$+ \cos \pi n \left\{ \left( \frac{\cos 2x}{2^n} - \frac{1}{2} \cdot \frac{\cos 4x}{4^n} + \frac{1.3}{2.4} \cdot \frac{\cos 6x}{6^n} - \dots \right) \right.$$

$$\left. - \left( \frac{\cos 2x}{2^{n+1}} - \frac{1}{2} \cdot \frac{\cos 4x}{4^{n+1}} + \frac{1.3}{2.4} \cdot \frac{\cos 6x}{6^{n+1}} - \dots \right) \right\} \text{ then}$$

If  $n$  is odd,

$$i. \frac{F(n)}{2} \sin \frac{\pi n}{2} = \frac{x^n}{1^n} S_0 \phi(0) - \frac{x^{n-2}}{1^{n-2}} \left\{ S_0 \phi(2) + \frac{S_2}{2^2} \phi(0) \right\}$$

$$+ \frac{x^{n-4}}{1^{n-4}} \left\{ S_0 \phi(4) + \frac{S_2}{2^2} \phi(2) + \frac{S_4}{2^4} \phi(0) \right\} - \dots$$

$$= \frac{A_{n-1}}{1^{n-1}} \phi(0) - \frac{A_{n-3}}{1^{n-3}} \phi(2) + \frac{A_{n-5}}{1^{n-5}} \phi(4) - \dots \quad I.$$

$$ii. \frac{F(x+1)}{3} \sin \frac{\pi x}{2} = \frac{x^n}{1^n} S_0 \phi(1) - \frac{x^{n-1}}{1^{n-1}} \left\{ S_0 \phi(2) + \frac{S_2}{2!} \phi(1) \right\}$$

$$+ \frac{x^{n-2}}{1^{n-2}} \left\{ S_0 \phi(3) + \frac{S_2}{2!} \phi(2) + \frac{S_4}{4!} \phi(1) \right\} - \dots$$

$$= \frac{A_{n-1}}{1^{n-1}} \phi(1) - \frac{A_{n-3}}{1^{n-3}} \phi(3) + \frac{A_{n-5}}{1^{n-5}} \phi(5) - \dots \quad II.$$

where  $S_n = \frac{1}{1^n} - \frac{1}{2^n} + \frac{1}{3^n} - \frac{1}{4^n} + \dots$

$$\frac{\pi}{2} \phi(x) = \frac{1}{1^{n+1}} + \frac{1}{2} \cdot \frac{1}{3^{n+1}} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{5^{n+1}} + \dots$$

and  $\frac{2}{\pi} A_n = \left(\frac{\pi}{2}\right)^2 + \left(\frac{3\pi}{2}\right)^2 + \left(\frac{5\pi}{2}\right)^2 + \dots + \left(x - \frac{\pi}{2}\right)^2$

If  $x$  is even  $\psi\left(\frac{n+1}{2}\right) \cos \frac{\pi x}{2} = -1$  or  $\psi\left(\frac{n+1}{2}\right) \cos \frac{\pi x}{2} = 1$

Sub. From the following identities the I part of the theorem is obtained.

i.  $\sin x - \frac{1}{2} \sin 3x + \frac{1 \cdot 3}{2 \cdot 4} \sin 5x - \dots = \frac{1}{2} \sin 2x - \frac{1 \cdot 3}{2 \cdot 4} \sin 4x$   
 $+ \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \sin 6x - \dots = \frac{\sin \frac{x}{2}}{\sqrt{2} \cos x}$

ii.  $\cos x - \frac{1}{2} \cos 3x + \frac{1 \cdot 3}{2 \cdot 4} \cos 5x - \dots = 1 - \frac{1}{2} \cos 2x +$   
 $\frac{1 \cdot 3}{2 \cdot 4} \cos 4x - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cos 6x + \dots = \frac{\cos \frac{x}{2}}{\sqrt{2} \cos x}$

iii.  $\sin 2x - \frac{1}{2} \sin 4x + \frac{1 \cdot 3}{2 \cdot 4} \sin 6x - \dots = \frac{\sin \frac{3x}{2}}{\sqrt{2} \cos x}$

iv.  $\cos 2x - \frac{1}{2} \cos 4x + \frac{1 \cdot 3}{2 \cdot 4} \cos 6x - \dots = \frac{\cos \frac{3x}{2}}{\sqrt{2} \cos x}$

v.  $\frac{\sin 2x}{2} - \frac{1}{2} \cdot \frac{\sin 4x}{4} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\sin 6x}{6} - \dots = \sin \frac{x}{2} \sqrt{2} \cos x$

vi.  $\frac{\cos 2x}{2} - \frac{1}{2} \cdot \frac{\cos 4x}{4} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\cos 6x}{6} - \dots = \cos \frac{x}{2} \sqrt{2} \cos x - 1$

vii.  $\frac{\sin x}{1} - \frac{1}{2} \cdot \frac{\sin 3x}{3} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\sin 5x}{5} - \dots = \sin^{-1}(\sqrt{2} \sin \frac{x}{2})$

viii.  $\frac{\cos x}{1} - \frac{1}{2} \cdot \frac{\cos 3x}{3} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\cos 5x}{5} - \dots = \log(\sqrt{\cos x} + \sqrt{2} \cos \frac{x}{2})$

$$ix. \frac{\sin 2x}{2^2} = \frac{1}{2} \cdot \frac{\sin 4x}{4^2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{\sin 6x}{6^2} + \dots = \sin \frac{x}{2} \sqrt{2 \cos x} + \sin^{-1}(\sqrt{2} \sin \frac{x}{2}) - x.$$

$$x. \frac{\cos 2x}{2^2} = \frac{1}{2} \cdot \frac{\cos 4x}{4^2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{\cos 6x}{6^2} + \dots = \cos \frac{x}{2} \sqrt{2 \cos x} - \log(\sqrt{\cos x} + \sqrt{2} \cos \frac{x}{2}) - 1 + \log 2.$$

$$5. i. \sin a \theta + \frac{x^n}{4} \sin(a+2)\theta + \frac{x^{n(n-1)}}{16} \sin(a+4)\theta + \dots = 2^n \cos^n \theta \sin(a+n)\theta.$$

$$ii. \cos a \theta + \frac{x^n}{4} \cos(a+2)\theta + \frac{x^{n(n-1)}}{16} \cos(a+4)\theta + \dots = 2^n \cos^n \theta \cos(a+n)\theta.$$

6. If  $\phi(x) = \frac{x}{12} + \frac{x^2}{24} + \frac{x^3}{32} + \frac{x^4}{48} + \dots$  then

$$i. \phi(1-x) + \phi(1-\frac{1}{x}) = -\frac{1}{2}(\log x)^2.$$

$$ii. \phi(x) + \phi(-\frac{1}{x}) = -\frac{\pi^2}{6} - \frac{1}{2}(\log x)^2.$$

$$iii. \phi(x) + \phi(1-x) = \frac{\pi^2}{6} - \log x \log(1-x)$$

$$iv. \phi(x) + \phi(-x) = \frac{1}{2} \phi(x^2).$$

v. If  $\phi(x) - \phi(-x) = 2\psi(x) = 2(\frac{x}{12} + \frac{x^3}{48} + \frac{x^5}{56} + \dots)$  then

$$\psi(x) + \psi(\frac{1}{1+x}) = \frac{\pi^2}{8} + \frac{1}{2} \log x \log \frac{1+x}{1-x}$$

$$vi. \phi(\frac{x}{1-y}) + \phi(\frac{y}{1-x}) = \phi(x) + \phi(y) + \frac{xy}{(1-x)(1-y)} + \log(1-x) \log(1-y)$$

$$vii. \phi(e^{-x}) = \frac{\pi^2}{6} + x \log x - x - \frac{x^2}{4} + \frac{B_2}{24} x^4 - \frac{B_4}{48} x^6 + \dots$$

$$viii. \phi(0 - e^{-x}) = x - \frac{x^2}{4} + B_2 \frac{x^3}{12} - B_4 \frac{x^5}{48} + B_6 \frac{x^7}{12} - \dots$$

$$E.g. i. \phi(\frac{1}{2}) = \frac{\pi^2}{12} - \frac{1}{2}(\log 2)^2.$$

$$ii. \phi(\frac{\sqrt{5}-1}{2}) = \frac{\pi^2}{10} - (\log \frac{\sqrt{5}-1}{2})^2$$

$$iii. \phi(\frac{3-\sqrt{5}}{2}) = \frac{\pi^2}{15} - (\log \frac{\sqrt{5}-1}{2})^2$$

$$iv. \psi(\sqrt{2}-1) = \frac{\pi^2}{16} - \frac{1}{2}(\log \sqrt{2}-1)^2$$

$$v. \psi(\frac{\sqrt{5}-1}{2}) = \frac{\pi^2}{12} - \frac{3}{4}(\log \frac{\sqrt{5}-1}{2})^2$$

$$vi. \psi(\sqrt{5}-2) = \frac{\pi^2}{24} - \frac{3}{4}(\log \frac{\sqrt{5}-1}{2})^2$$

7.  $f/\phi(x) = \frac{x^2}{1^2} + \frac{x^4}{2^2} + \frac{x^6}{3^2} + \dots$  then  
 i.  $\phi(x) + \phi(1-x) + \phi(x) = S_3 + \frac{\pi^2}{6} \log x - \frac{1}{2} (\log x)^2 \log(1-x) + \frac{1}{2} (\log x)^3$

ii.  $\phi(x) - \phi(1-x) = -\frac{1}{6} (\log x)^3 - \frac{\pi^2}{6} \log x$

iii.  $\phi(x) + \phi(-x) = \frac{1}{2} \phi(x^2)$

∴ g. i.  $\phi(\frac{1}{2}) = \frac{1}{6} (\log 2)^3 - \frac{\pi^2}{12} \log 2 + (\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots)$

ii.  $\phi(\frac{1-\sqrt{5}}{2}) = \frac{1}{3} (\log \frac{1+\sqrt{5}}{2})^3 - \frac{2\pi^2}{15} \log \frac{1+\sqrt{5}}{2} + S_3$

8.  $f/\phi(x) = x + (1+\frac{1}{2})\frac{x^2}{3} + (1+\frac{1}{2}+\frac{1}{3})\frac{x^3}{5} + \dots$  then  
 $\phi(\frac{x}{1-x}) = \frac{1}{8} (\log |1-x|)^2 + \frac{1}{2} (\frac{\pi}{10} + \frac{\pi^2}{15} + \frac{x^3}{3^2} + \dots)$

∴ g. i.  $\phi(\frac{1}{3}) = \frac{\pi^2}{24} - \frac{1}{8} (\log 2)^2$

ii.  $\phi(\frac{1}{5}) = \frac{\pi^2}{20} - \frac{3}{8} (\log \frac{1+\sqrt{5}}{2})^2$

iii.  $\phi(\frac{1}{10}) = \frac{\pi^2}{30} - \frac{3}{4} (\log \frac{1+\sqrt{5}}{2})^2$

9.  $f/\phi(x) = \frac{x^2}{1^2} + (1+\frac{1}{2})\frac{x^3}{3^2} + (1+\frac{1}{2}+\frac{1}{3})\frac{x^4}{5^2} + \dots$  then

i.  $\phi(1-x) = \frac{1}{2} \log(1-x) (\log x)^2 + \log x (\frac{\pi}{12} + \frac{x^2}{2^2} + \frac{x^3}{3^2} + \dots) - (\frac{\pi}{12} + \frac{x^2}{2^2} + \frac{x^3}{3^2} + \dots) + S_3$

ii.  $\phi(1-x) - \phi(1-\frac{1}{2}) = \frac{1}{8} (\log x)^3$

iii.  $\phi(1-x) = \frac{1}{2} \log(1-x) (\log x)^2 - \frac{1}{3} (\log x)^3 - \log x (\frac{1}{1^2} x + \frac{1}{2^2} x^2 + \dots) - (\frac{1}{1^2} x + \frac{1}{2^2} x^2 + \frac{1}{3^2} x^3 + \dots) + S_3$

iv.  $\phi(-x) + \phi(-\frac{1}{2}) = -\frac{1}{6} (\log x)^3 + \log x (\frac{\pi}{12} - \frac{x^2}{12} + \frac{x^3}{3^2} - \dots) - (\frac{\pi}{12} - \frac{x^2}{12} + \frac{x^3}{3^2} - \dots) + S_2$

10.  $f/\phi(x) = \frac{x^2}{2^2} + (1+\frac{1}{2})\frac{x^3}{3^2} + (1+\frac{1}{2}+\frac{1}{3})\frac{x^4}{5^2} + \dots$  then

i.  $\phi(1-x) - \phi(1-\frac{1}{2}) = \frac{1}{24} (\log x)^4 - \frac{1}{6} (\log x)^3 \log(1-x) - S_3 \log x$

$$10h. + 2 \left( \frac{x}{14} + \frac{x^2}{28} + \frac{x^3}{42} + \dots \right) - \log x \left( \frac{x}{12} + \frac{x^2}{24} + \frac{x^3}{36} + \dots \right) - \frac{\pi^2}{48}$$

$$ii. \phi(x) - \phi\left(\frac{x}{2}\right) = \frac{1}{24} (\log x)^2 - \log x \left( \frac{x}{12} - \frac{x^2}{24} + \frac{x^3}{36} - \dots \right) + 2 \left( \frac{x}{14} - \frac{x^2}{28} + \frac{x^3}{42} - \dots \right) - S_3 \log x - \frac{7\pi^2}{360}$$

$$11. \text{ If } \phi(x) = \frac{x^2}{2} + (1+\frac{1}{2}) \frac{x^3}{3} + (1+\frac{1}{2}+\frac{1}{3}) \frac{x^4}{4} + \dots + \dots$$

$$\psi(x) = \frac{x^2}{2} + (1+\frac{1}{2}) \frac{x^3}{3} + (1+\frac{1}{2}+\frac{1}{3}) \frac{x^4}{4} + \dots + \dots$$

$$i. \phi\left(\frac{1-x}{1+x}\right) = \frac{1}{8} (\log x)^2 \log \frac{1-x}{1+x} + \frac{1}{2} \log x \left( \frac{x}{12} + \frac{x^2}{24} + \frac{x^3}{36} + \dots \right) + \frac{1}{2} \left( \frac{1-x}{12} + \frac{1-x^2}{36} + \frac{1-x^3}{54} + \dots \right)$$

$$ii. \psi(x) + \psi\left(\frac{1-x}{1+x}\right) = \phi(x) \log x + \phi\left(\frac{1-x}{1+x}\right) \log \frac{1-x}{1+x}$$

$$- \frac{1}{16} (\log x)^2 (\log \frac{1-x}{1+x})^2 + \frac{\pi^2}{24} \left( \frac{1}{12} - \frac{1}{36} + \frac{1}{54} - \dots \right) - \frac{\pi^2}{3\sqrt{3}} \left( \frac{1}{12} + \frac{1}{36} + \frac{1}{54} + \dots \right)$$

$$12. \text{ If } \phi(x) = x + (1+\frac{1}{2}) \frac{x^2}{3} + (1+\frac{1}{2}+\frac{1}{3}) \frac{x^3}{5} + \dots + \dots \text{ then}$$

$$\phi\left(\frac{1-x}{1+x}\right) = -(1-\log 2) \log x + \frac{1+x}{1-x} \log \frac{1-x}{(1+x)^2} + \frac{1}{2} (\log x)^2 + \frac{\pi^2}{12}$$

$$= \left( \frac{x}{12} - \frac{x^2}{24} + \frac{x^3}{36} - \dots \right)$$

$$\text{E.g. } i. \frac{1}{2} + \frac{1+\frac{1}{2}}{2^2} \cdot \frac{1}{2} + \frac{1+\frac{1}{2}+\frac{1}{3}}{3^2} \cdot \frac{1}{2} + \dots = \frac{1}{3} - \frac{\pi^2}{12} \log 2$$

$$ii. \frac{1}{12} + \frac{1+\frac{1}{2}+\frac{1}{3}}{3^2} + \frac{1+\frac{1}{2}+\frac{1}{3}+\frac{1}{4}}{4^2} + \dots = \frac{3}{2} \left( \frac{1}{12} + \frac{1}{36} + \frac{1}{54} + \dots \right)$$

$$iii. \frac{1}{12} + \frac{1+\frac{1}{2}}{2^2} + \frac{1+\frac{1}{2}+\frac{1}{3}}{3^2} + \dots = 2 \left( \frac{1}{12} + \frac{1}{36} + \frac{1}{54} + \dots \right)$$

$$iv. (\sqrt{5}-2) + \frac{1+\frac{1}{2}}{3} (\sqrt{5}-2)^2 + \frac{1+\frac{1}{2}+\frac{1}{3}}{5} (\sqrt{5}-2)^3 + \dots$$

$$= \frac{\pi^2}{60} + \frac{3}{4} (\log \frac{\sqrt{5}-1}{2})^2 + (\sqrt{5}+2) \log 4 + (3\sqrt{5}+5+\log 2) \log \frac{\sqrt{5}-1}{2}$$

$$13. S_{n+1} \cos \frac{\pi n}{2} \int_0^{\frac{\pi}{2}} \frac{x^n}{2} \cot \frac{x}{2} dx + x^n \left( \frac{\cos x}{1} + \frac{\cos 2x}{2} + \dots \right)$$

$$= n x^{n-1} \left( \frac{\sin x}{12} + \frac{\sin 2x}{24} + \frac{\sin 3x}{36} + \dots \right)$$

$$= n(n-1) x^{n-2} \left( \frac{\cos 2x}{12} + \frac{\cos 4x}{24} + \frac{\cos 6x}{36} + \dots \right)$$

$$+ n(n-1)(n-2) x^{n-3} \left( \frac{\sin x}{12} + \frac{\sin 2x}{24} + \frac{\sin 3x}{36} + \dots \right) + \dots$$

where  $S_n = \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots$  and  $S_1 = -\log 2$ .

Sol.  $\sin x + \sin 2x + \sin 3x + \dots = \frac{1}{2} \cot \frac{x}{2}$ .

$\therefore \int x^n (\sin x + \sin 2x + \dots) dx = \int \frac{x^n}{2} \cot \frac{x}{2} dx$ .

4.  $S_{n+1} \cos \frac{\pi}{2} \lfloor n = \int \frac{x^n}{2 \sin x} dx$   
 $+ x^n \left( \frac{\cos x}{1} + \frac{\cos 3x}{3} + \frac{\cos 5x}{5} + \dots \right)$   
 $- n x^{n-1} \left( \frac{\sin x}{1^2} + \frac{\sin 3x}{3^2} + \frac{\sin 5x}{5^2} + \dots \right)$   
 $- n(n-1) x^{n-2} \left( \frac{\cos x}{1^3} + \frac{\cos 3x}{3^3} + \frac{\cos 5x}{5^3} + \dots \right) + \dots$

Sol.  $\sin x + \sin 2x + \sin 3x + \dots = \frac{1}{2} \operatorname{cosec} x$ .

15. If  $\int x^n \cot x dx = f_n(x)$  then

$\int_0^{\frac{\pi}{2}} f_n\left(\frac{\pi}{2}-x\right) dx = \pi^n \left\{ f_0(\pi) - f_0(0) \right\} - \frac{n}{2} \pi^{n-1} \left\{ f_1(\pi) - 2f_1(0) \right\}$   
 $+ \frac{n(n-1)}{2} \pi^{n-2} \left\{ f_2(\pi) - 2^2 f_2(0) \right\} - \frac{n(n-1)(n-2)}{6} \pi^{n-3} \left\{ f_3(\pi) - 2^3 f_3(0) \right\} + \dots$

Sol.  $\tan x = \cot x - 2 \cot 2x$  and

$\int_0^{\frac{\pi}{2}} (\frac{\pi}{2}-x) dx = - \int (\frac{\pi}{2}-x) \cot (\frac{\pi}{2}-x) dx = - \int (\frac{\pi}{2}-x) \tan x dx$ .

N.B. Let  $\sin x = y$  and  $\tan x = z$ , then

$\int x^n \cot x dx = \int \frac{x^n}{\sin x} \cos x dx = \int \left( \frac{\sin^{-1} y}{y} \right)^n dy$ , and

$\int \frac{z x^n}{\sin^2 x} dx = \int \frac{x^n}{\cos x \sin x} dx = \int \frac{x^n}{\tan x} \sec^2 x dx$   
 $= \int \frac{(\tan^{-1} z)^n}{z} dz$ .

i.  $\frac{1}{2} (\tan^{-1} x)^2 = \frac{x^2}{2} - (1 + \frac{1}{3}) \frac{x^4}{4} + (1 + \frac{1}{3} + \frac{1}{5}) \frac{x^6}{6} - \dots$



ii.  $(2(\sin^{-1}x))^2 = \frac{x^2}{2} + \frac{1}{3} \cdot \frac{x^4}{4} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{x^6}{6} + \dots$

iii.  $\frac{1}{13} (\sin^{-1}x)^3 = \frac{1}{2} \cdot \frac{x^3}{3} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{x^5}{5} (1 + \frac{1}{3 \cdot 2}) + \frac{1 \cdot 2 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{x^7}{7} (1 + \frac{1}{3 \cdot 2} + \frac{1}{5 \cdot 4}) + \dots$

iv.  $\frac{1}{13} (\sin^{-1}x)^4 = \frac{1}{3} \cdot \frac{x^4}{4} \cdot \frac{1}{2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{x^6}{6} (\frac{1}{2} + \frac{1}{4}) + \frac{1 \cdot 2 \cdot 6}{2 \cdot 4 \cdot 7} \cdot \frac{x^8}{8} (\frac{1}{2} + \frac{1}{4} + \frac{1}{6}) + \dots$

16.  $\frac{\sin x}{1^2} + \frac{1}{2} \cdot \frac{\sin^3 x}{3^2} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{\sin^5 x}{5^2} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{\sin^7 x}{7^2} + \dots$   
 $= x \log 2 \sin x + \frac{1}{2} \left( \frac{\sin^3 x}{3^2} + \frac{\sin^5 x}{5^2} + \frac{\sin^7 x}{7^2} + \dots \right)$

c.g.  $\frac{1}{1^2} + \frac{1}{2} \cdot \frac{1}{3^2} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{5^2} + \dots = \frac{\pi}{2} \log 2.$

ii.  $\frac{1}{1^2} + \frac{1}{2} \cdot \frac{1}{3^2} \cdot \frac{1}{2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{1}{5^2} \cdot \frac{1}{2} + \frac{1 \cdot 2 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{1}{7^2} \cdot \frac{1}{2} + \dots$   
 $= \frac{\pi}{4\sqrt{2}} (\log 2 + \frac{1}{\sqrt{2}} (\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots))$

iii.  $\frac{1}{1^2} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{3^2} \cdot \frac{1}{2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{1}{5^2} \cdot \frac{1}{2} + \dots$   
 $= \frac{3\sqrt{3}}{4} (\frac{1}{1^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots) - \frac{\pi^2}{6\sqrt{3}}$

iv.  $\frac{1}{1^2} + \frac{1}{2} \cdot \frac{1}{3^2} \cdot \frac{3}{2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{1}{5^2} \cdot (\frac{3}{2})^2 + \dots = \frac{\pi}{3\sqrt{3}} \log 3 - \frac{2\pi^2}{27}$   
 $+ (\frac{1}{1^2} + \frac{1}{4^2} + \frac{1}{7^2} + \dots)$

17.  $\frac{\tan x}{1^2} - \frac{\tan^3 x}{3^2} + \frac{\tan^5 x}{5^2} - \dots$   
 $= x \log \tan x + \frac{\sin^2 x}{1^2} + \frac{\sin^6 x}{5^2} + \frac{\sin^{10} x}{5^2} + \dots$

c.g. i.  $\int_0^{\sqrt{3}} \frac{\tan^{-1} x}{x} dx = -\frac{\pi}{12} \log 3 - \frac{5\pi^2}{18\sqrt{3}} + \frac{5\sqrt{3}}{4} (\frac{1}{1^2} + \frac{1}{4^2} + \frac{1}{7^2} + \dots)$

ii.  $\int_0^{\sqrt{2}-1} \frac{\tan^{-1} x}{x} dx = \frac{\pi}{8} \log(\sqrt{2}-1) - \frac{\pi^2}{16} + \sqrt{2} (\frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots)$

iii.  $\int_0^{2-\sqrt{3}} \frac{\tan^{-1} x}{x} dx = \frac{\pi}{12} \log(2-\sqrt{3}) + \frac{2}{3} \int_0^1 \frac{\tan^{-1} x}{x} dx$

N. 13.  $\int_0^1 \frac{\tan^{-1} x}{x} dx = \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \frac{1}{7^2} + \dots$   
 $= .915965594177$

18 When x lies between 0 and  $\frac{\pi}{4}$ .

$$\frac{\cos^2 x - \sin^2 x}{2} + \frac{1}{2} \cdot \frac{\cos^4 x - \sin^4 x}{3^2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{\cos^6 x - \sin^6 x}{5^2} + \dots$$

$$= \frac{\pi}{2} \log 2 \cos x - \frac{1}{2} \left\{ \frac{\sin^2 2x}{1^2} + \frac{1}{2} \cdot \frac{\sin^4 2x}{3^2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{\sin^6 2x}{5^2} + \dots \right\}$$

e.g. If  $\psi(x) = \int_0^x \frac{\sin t}{t} dt$  then

$$\psi\left(\frac{2}{3}\right) - \frac{1}{2} \psi\left(\frac{2\sqrt{2}}{2\sqrt{2}}\right) = \frac{\pi}{2} \log 2 + 2 \psi\left(\frac{1}{\sqrt{2}}\right) - 2 \psi\left(\frac{2}{\sqrt{2}}\right)$$

$$19. \frac{\cos^2 x + \sin^2 x}{1^2} + \frac{1}{2} \cdot \frac{\cos^4 x + \sin^4 x}{3^2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{\cos^6 x + \sin^6 x}{5^2} + \dots$$

$$= \frac{\pi}{2} \log 2 \cos x + \frac{\tan^2 x}{1^2} - \frac{\tan^4 x}{3^2} + \frac{\tan^6 x}{5^2} - \dots$$

e.g.  $1^2 \cdot \frac{1+2}{5} + \frac{1}{2} \cdot \frac{1}{2^2} \cdot \frac{1+2^3}{5^2} + \frac{1 \cdot 2}{2 \cdot 4} \cdot \frac{1}{5^2} \cdot \frac{1+2^5}{5^2} + \dots$

$$= \frac{\pi}{2\sqrt{5}} \log \frac{4}{\sqrt{5}} + \frac{1}{\sqrt{5}} \left( \frac{1}{1^2} \cdot 2 - \frac{1}{3^2} \cdot \frac{1}{23} + \frac{1}{5^2} \cdot \frac{1}{25} - \dots \right)$$

$$20. \frac{\sin^2 x}{2^2} + \frac{2}{3} \cdot \frac{\sin^4 x}{4^2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{\sin^6 x}{6^2} + \dots$$

$$= \frac{x^2}{2} \log 2 \sin x + \frac{\pi}{2} \left( \frac{\sin 2x}{1^2} + \frac{\sin 4x}{2^2} + \frac{\sin 6x}{3^2} + \dots \right)$$

$$+ \frac{1}{2} \left( \frac{\cos 2x}{1^2} + \frac{\cos 4x}{2^2} + \frac{\cos 6x}{3^2} + \dots \right)$$

e.g. i.  $\frac{1}{2^2} + \frac{2}{3} \cdot \frac{4}{4^2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{1}{6^2} + \frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7} \cdot \frac{1}{8^2} + \dots$

$$= \frac{\pi^2}{8} \log 2 - \frac{1}{2} \left( \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right)$$

ii.  $\frac{1}{2^2} \cdot \frac{1}{2} + \frac{2}{3} \cdot \frac{4}{4^2} \cdot \frac{1}{2^2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{1}{6^2} \cdot \frac{1}{2^2} + \dots$

$$= \frac{\pi^2}{64} \log 2 + \frac{\pi}{8} \left( \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots \right) - \frac{5}{16} \left( \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right)$$

$$21. \frac{\tan^2 x}{2^2} - \left(1 + \frac{1}{3}\right) \frac{\tan^4 x}{4^2} + \left(1 + \frac{1}{3} + \frac{1}{5}\right) \frac{\tan^6 x}{6^2} - \dots$$

$$= \frac{x^2}{2} \log \tan x + \pi \left( \frac{\sin 2x}{1^2} + \frac{\sin 6x}{3^2} + \frac{\sin 10x}{5^2} + \dots \right)$$

$$+ \frac{1}{2} \left( \frac{\cos 2x}{1^2} + \frac{\cos 6x}{3^2} + \frac{\cos 10x}{5^2} + \dots \right) - \frac{1}{2} \left( \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right)$$

e.g.  $\frac{1}{1-x} = \frac{1+\frac{1}{2}}{2x} + \frac{1+\frac{1}{2}+\frac{1}{4}}{5x^2} + \dots$

$$= \frac{\pi}{2} \left( \frac{1}{1-x} + \frac{1}{1-x} + \frac{1}{1-x} + \dots \right) - \frac{1}{2} \left( \frac{1}{1-x} + \frac{1}{1-x} + \frac{1}{1-x} + \dots \right)$$

22.  $\frac{\cos^2 x + \sin^2 x}{2} + \frac{2}{3} \cdot \frac{\cos^4 x + \sin^4 x}{4} + \frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7} \cdot \frac{\cos^6 x + \sin^6 x}{8} + \dots$

$$= -\frac{\pi^2}{8} \log 2 \cos x$$

$$+ \frac{\pi}{6} \left\{ \frac{\cos^2 x}{1^2} + \frac{1}{2} \cdot \frac{\cos^4 x}{2^2} + \frac{1 \cdot 3}{6 \cdot 4} \cdot \frac{\cos^6 x}{3^2} + \dots \right\}$$

$$+ \frac{1}{4} \left\{ \frac{\sin^2 2x}{2^2} + \frac{2}{3} \cdot \frac{\sin^4 2x}{4^2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{\sin^6 2x}{6^2} + \dots \right\}$$

$$- \frac{1}{2} \left( \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots \right)$$

23.  $\frac{\tan^2 x}{2} - (1+\frac{1}{3}) \frac{\tan^4 x}{4} + (1+\frac{1}{3}+\frac{1}{5}) \frac{\tan^6 x}{6} - \dots$

$$= 2 \left\{ \frac{\sin^2 x}{2^2} + \frac{2}{3} \cdot \frac{\sin^4 x}{4^2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{\sin^6 x}{6^2} + \dots \right\}$$

$$- \frac{1}{4} \left\{ \frac{\sin^2 2x}{2^2} + \frac{2}{3} \cdot \frac{\sin^4 2x}{4^2} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{\sin^6 2x}{6^2} + \dots \right\}$$

24. If  $x \cos \theta + y \cos \phi = 1$ , and  $x \sin \theta + y \sin \phi = 0$  then

i.  $\frac{x^2}{1^2} \cos \theta + \frac{x^2}{2^2} \cos 2\theta + \frac{x^2}{3^2} \cos 3\theta + \dots$

$$+ \frac{y^2}{1^2} \cos \phi + \frac{y^2}{2^2} \cos 2\phi + \frac{y^2}{3^2} \cos 3\phi + \dots$$

$$= \frac{\pi^2}{6} - \log x \log y + \theta \phi$$

ii.  $\frac{x^2}{1^2} \sin \theta + \frac{x^2}{2^2} \sin 2\theta + \frac{x^2}{3^2} \sin 3\theta + \dots$

$$+ \frac{y^2}{1^2} \sin \phi + \frac{y^2}{2^2} \sin 2\phi + \frac{y^2}{3^2} \sin 3\phi + \dots = -\phi (\log x - \theta \log y)$$

25. If  $x \cos \theta + y \cos \phi = xy \cos(\theta + \phi)$ , &  $x \sin \theta + y \sin \phi = xy \sin(\theta + \phi)$ , then

i.  $\frac{x^2}{1^2} \cos \theta + \frac{x^2}{2^2} \cos 2\theta + \frac{x^2}{3^2} \cos 3\theta + \dots$

$$+ \frac{y^2}{1^2} \cos \phi + \frac{y^2}{2^2} \cos 2\phi + \frac{y^2}{3^2} \cos 3\phi + \dots$$

$$= \frac{1}{2} \log(1-2x \cos \theta + x^2) \log(1-2y \cos \phi + y^2)$$

$$- \frac{1}{2} \tan^{-1} \frac{\sin \theta}{1-x \cos \theta} \tan^{-1} \frac{y \sin \phi}{1-y \cos \phi}$$

$$\begin{aligned}
 11. & \frac{x}{12} \sin \theta + \frac{x^2}{12} \sin 2\theta + \frac{x^3}{12} \sin 3\theta + \dots \\
 & + \frac{y}{12} \sin \phi + \frac{y^2}{12} \sin 2\phi + \frac{y^3}{12} \sin 3\phi + \dots \\
 & = -\frac{1}{4} \log(1 - 2x \cos \theta + x^2) \tan^{-1} \frac{y \sin \phi}{1 - y \cos \phi} \\
 & \quad - \frac{1}{4} \log(1 - 2y \cos \phi + y^2) \tan^{-1} \frac{x \sin \theta}{1 - x \cos \theta}
 \end{aligned}$$

26.  $x \cos \theta + y \cos \phi + xy \cos(\theta + \phi) = 1$  and  
 $x \sin \theta + y \sin \phi + xy \sin(\theta + \phi) = 0$ , then

$$\begin{aligned}
 i. & \frac{x}{12} \cos \theta + \frac{x^3}{32} \cos 3\theta + \frac{x^5}{52} \cos 5\theta + \dots \\
 & + \frac{y}{12} \cos \phi + \frac{y^3}{32} \cos 3\phi + \frac{y^5}{52} \cos 5\phi + \dots \\
 & = \frac{\pi^2}{8} - \frac{1}{2} \log x \log y + \frac{1}{2} \theta \phi
 \end{aligned}$$

$$\begin{aligned}
 ii. & \frac{x}{12} \sin \theta + \frac{x^3}{32} \sin 3\theta + \frac{x^5}{52} \sin 5\theta + \dots \\
 & + \frac{y}{12} \sin \phi + \frac{y^3}{32} \sin 3\phi + \frac{y^5}{52} \sin 5\phi + \dots \\
 & = -\frac{1}{2} \phi \log x - \frac{1}{2} \theta \log y
 \end{aligned}$$

27.  $1^2 \log 1 + 2^2 \log 2 + 3^2 \log 3 + 4^2 \log 4 + \dots + x^2 \log x = \phi_n(x)$

$$\begin{aligned}
 \phi_n(x) &= C_n + (1^2 + 2^2 + 3^2 + \dots + x^2 - S_{-2}) \log x - \frac{x^{n+1}}{(n+1)!} \\
 & + \frac{B_2}{12} \cdot n \cdot \frac{1}{2} \cdot x^{n-1} - \frac{B_4}{120} n(n-1)(n-2) \left( \frac{1}{n} + \frac{1}{n^2} + \frac{1}{n^3} \right) x^{n-3} + \dots
 \end{aligned}$$

$$\begin{aligned}
 \text{and } C_n &= \frac{B_n}{n} \left\{ \cos \frac{\pi n}{2} \left( \frac{1}{n-1} - C_0 - \log 2\pi \right) - \frac{\pi}{2} \sin \frac{\pi n}{2} \right\} \\
 & - 2 \frac{(n-1)}{(2\pi)^n} \cos \frac{\pi n}{2} \left\{ \frac{\log 1}{1^2} + \frac{\log 2}{2^2} + \frac{\log 3}{3^2} + \dots \right\}
 \end{aligned}$$

Cor. If  $n$  is even  $C_n = -\frac{\pi}{2} \cdot \frac{B_{n+1}}{n+1} \cos \frac{\pi n}{2} = -\frac{12}{2(2\pi)^{n+1}} \sum_{r=1}^{\infty} \cos \frac{\pi n r}{2}$

$$C_2 = \frac{1}{2} \log 2\pi, \quad C_4 = \frac{5}{4\pi^2}, \quad C_6 = -\frac{3}{4\pi^4}, \quad C_8 = \frac{45}{8\pi^6} \dots$$

e.g. i.  $\frac{(1^2 \cdot 2^2 \cdot 3^2 \cdot 4^2 \dots x^2) \cdot C_n}{(x^2 - \frac{1}{2})}$  when  $x = \infty$

$$\begin{aligned}
 & \sqrt[3]{\frac{x(x+3)x}{x+\frac{1}{2}}} \Big|_{x=\infty} \\
 & = 1 \cdot \frac{1}{\pi} + 2 \cdot \frac{1}{2\pi} + 3 \cdot \frac{1}{3\pi} + 4 \cdot \frac{1}{4\pi} + \dots
 \end{aligned}$$

ii.  $\left\{ \left(\frac{1}{x}\right)^1 \cdot \left(\frac{2}{x}\right)^2 \cdot \left(\frac{3}{x}\right)^3 \cdot \left(\frac{4}{x}\right)^4 \dots \left(\frac{x}{x}\right)^{x^2} \right\} e^{\frac{x^2}{9} \frac{x}{12}}$  when  $x = \infty$   
 $= e^{\frac{x^2}{9} \left( \frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8} + \dots \right)}$

28.  $\phi_n(x) + n C_{n,1} x + \frac{n(n-1)}{2} C_{n,2} x^2 + \frac{n(n-1)(n-2)}{6} C_{n,3} x^3 + \dots + C_{n,n} x^n$   
 $+ S_1 \frac{x^{n+1}}{n+1} - S_2 \frac{x^{n+2}}{(n+1)(n+2)} + S_3 \frac{x^{n+3}}{(n+1)(n+2)(n+3)} - \dots = f(x, c)$

where  $C_n$  is the constant of  $1^n \log 1 + 2^n \log 2 + 3^n \log 3 + \dots$   
 &  $S_n$  is that of  $\frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots$  and

$f(x, c) = (1^n + 2^n + 3^n + \dots + x^n) S_{-n} = \frac{1}{2} - \frac{1}{2^n} B_n x^{n-1}$   
 $+ \frac{n(n-1)(n-2)}{6} B_3 x^{n-3} (1 + \frac{1}{2} + \frac{1}{3}) - \frac{n(n-1)(n-2)(n-3)(n-4)}{24} B_5 x^{n-5} (1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5}) + \dots$   
 $= \frac{1^n + 2^n + 3^n + \dots + x^n}{n} + n \int_0^x f(x, n-1) dx$

29.  $\phi_n(x) - n^2 \left\{ \phi_n\left(\frac{x}{n}\right) + \phi_n\left(\frac{x-1}{n}\right) + \phi_n\left(\frac{x-2}{n}\right) + \dots + \phi_n\left(\frac{x-n+1}{n}\right) \right\}$   
 $= (1^n + 2^n + 3^n + \dots + x^n - S_{-n}) \log n - (n^2 + 1) C_n$

Cor. 1.  $\phi_n\left(\frac{1}{n}\right) + \phi_n\left(\frac{2}{n}\right) + \dots + \phi_n\left(\frac{n-1}{n}\right) = \frac{\log n}{n^2} S_{-n} + (n - \frac{1}{n}) C_n$

Cor. 2.  $\phi_n\left(\frac{1}{2}\right) = \frac{\log 2}{2^n} S_{-n} + (2 - \frac{1}{2^n}) C_n$

30. i. If  $n$  is even

$\phi_n(x-1) + \phi_n(x) = 2C_n + \frac{1x}{(2\pi)^2} \cos \frac{\pi x}{2} \left\{ \frac{\cos 3\pi x}{1^{2+1}} + \frac{\cos 5\pi x}{2^{2+1}} + \dots \right\}$

ii. If  $n$  is odd

$\phi_n(x-1) - \phi_n(x) = \frac{1x}{(2\pi)^2} \sin \frac{\pi x}{2} \left\{ \frac{\sin 2\pi x}{1^{2+1}} + \frac{\sin 4\pi x}{2^{2+1}} + \dots \right\}$

Sol.  $\frac{1}{x-1} - \frac{1}{x} = -\pi \cot \pi x = -\frac{\pi}{2} (\sin 2\pi x + \sin 4\pi x + \dots)$

Integrate both sides  $n+1$  times

13. More general theorems true for all values of  $n$  can be got by differentiating VIII. 15. and 16. with respect to  $n$ .

31. If  $1 \log 1 + 2 \log 2 + 3 \log 3 + \dots + x \log x = \phi_1(x)$

and  $\pi \{ \phi_1(x-1) - \phi_1(x) \} + \pi x \log 2 \sin \pi x = \psi(x)$  then

$$i. \psi(x) = \sin \pi x + \frac{2}{3} \cdot \frac{\sin^3 \pi x}{3^2} + \frac{2 \cdot 4}{2 \cdot 5} \frac{\sin^5 \pi x}{5^2} + \dots$$

$$= \tan \pi x - (1 + \frac{1}{3}) \frac{\tan^3 \pi x}{3} + (1 + \frac{1}{3} + \frac{1}{5}) \frac{\tan^5 \pi x}{5} - \dots$$

$$ii. \psi(x) + \psi(\frac{1}{2}-x) = \frac{\pi}{2} \log 2 \cos \pi x$$

$$+ \tan \pi x - \frac{\tan^3 \pi x}{3^2} + \frac{\tan^5 \pi x}{5^2} - \dots$$

$$iii. \psi(\frac{1}{2}-x) + \frac{1}{2} \psi(2x) - \psi(x) = \frac{\pi}{2} \log 2 \cos \pi x$$

$$iv. \psi(\frac{1}{2}-x) + \psi(\frac{1}{2}+x) = \pi \log 2 \cos \pi x$$

$$e.g. i. \psi(\frac{1}{2}) = \frac{\pi}{2} \log 2$$

$$ii. \psi(\frac{1}{3}) = (\frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots) + \frac{\pi}{2} \log 2$$

$$iii. \psi(\frac{1}{4}) = \frac{\sqrt{3}}{2} (\frac{1}{1^2} + \frac{1}{4^2} + \frac{1}{7^2} + \dots) - \frac{\pi^2}{9\sqrt{3}} + \frac{\pi}{6} \log 3$$

$$iv. \psi(\frac{1}{6}) = \frac{\sqrt{3}}{4} (\frac{1}{1^2} + \frac{1}{4^2} + \frac{1}{7^2} + \dots) - \frac{\pi^2}{6\sqrt{3}}$$

$$v. 2\psi(x) - \frac{1}{2}\psi(2x) = \tan \pi x - \frac{\tan^3 \pi x}{3^2} + \frac{\tan^5 \pi x}{5^2} - \dots$$

Similarly we can find peculiarities for  $\phi_2(x), \phi_3(x)$  &c.

$$27. \sin 2x + \frac{2}{3^2} \sin^3 2x + \frac{2 \cdot 4}{3 \cdot 5^2} \sin^5 2x + \dots$$

$$= 2(\tan x - \frac{\tan^3 x}{3^2} + \frac{\tan^5 x}{5^2} - \dots)$$

$$\text{Cos. } 1 + \frac{2 \cdot 4x}{3^2(1+x)^2} + \frac{2 \cdot 4}{3 \cdot 5^2} \left\{ \frac{4x}{1+x} \right\}^2 + \dots$$

$$= (1+x) \left( \frac{1}{1^2} - \frac{x}{3^2} + \frac{x^2}{5^2} - \frac{x^3}{7^2} + \dots \right)$$

$$ii. \tan 2x - \frac{2}{3^2} \tan^3 2x + \frac{2 \cdot 4}{3 \cdot 5^2} \tan^5 2x - \dots$$

$$= 2(\tan x + \frac{\tan^3 x}{3^2} + \frac{\tan^5 x}{5^2} + \dots)$$

$$e.g. i. 1 + \frac{2}{3^2} + \frac{2 \cdot 4}{3 \cdot 5^2} + \dots = 2 \left( \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots \right)$$

$$ii. 1 + \frac{2}{3^2} \cdot \frac{3}{2} + \frac{2 \cdot 4}{3 \cdot 5^2} \cdot \left(\frac{3}{2}\right)^2 + \dots = -\frac{\pi}{3\sqrt{3}} \log 3 - \frac{10}{27} \pi^2 + 5 \left( \frac{1}{1^2} + \frac{1}{4^2} + \frac{1}{7^2} + \dots \right)$$

$$iii. \frac{1}{2} + \frac{2}{3^2} \cdot \frac{1}{2^2} + \frac{2 \cdot 4}{3 \cdot 5^2} \cdot \frac{1}{2^3} + \dots = -\frac{\pi}{6} \log(2+\sqrt{3})$$

$$+ \frac{1}{3} \left( \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots \right)$$

iv.  $1 + \frac{2}{3^2} \cdot \frac{1}{2} + \frac{2 \cdot 4}{3 \cdot 5^2} \cdot \frac{1}{2^2} + \frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7^2} \cdot \frac{1}{2^3} + \dots$   
 $= -\frac{\pi}{2\sqrt{2}} \log(1+\sqrt{2}) - \frac{\pi^2}{4\sqrt{2}} + 4\left(\frac{1}{1^2} - \frac{1}{5^2} + \frac{1}{7^2} - \dots\right)$

v.  $(1 - \frac{1}{2}) + \frac{2}{3^2} (1 - \frac{1}{4}) + \frac{2 \cdot 4}{3 \cdot 5^2} (1 - \frac{1}{2^2}) + \dots = \frac{\pi}{2} \log(2 + \sqrt{3})$

vi.  $1 - \frac{2}{3^2} + \frac{2 \cdot 4}{3 \cdot 5^2} - \frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7^2} + \dots = \frac{\pi^2}{8} - \frac{1}{2} \{\log(1 + \sqrt{2})\}^2$

vii.  $2 - \frac{2}{3^2} \cdot \frac{1}{2^2} + \frac{2 \cdot 4}{3 \cdot 5^2} \cdot \frac{1}{2^3} - \dots = \frac{\pi^2}{12} - \frac{3}{2} (\log \frac{\sqrt{5}+1}{2})^2$

33. i.  $\int_0^{\frac{\pi}{2}} x \cos^n x \sin nx \, dx = \frac{\pi}{2n+1} (1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n})$   
 ii.  $\int_0^{\frac{\pi}{2}} \cos^n x \sin nx \, dx = \frac{1}{2n+1} (\frac{1}{1} + \frac{2}{2} + \frac{3}{3} + \dots + \frac{2n}{n})$

The above theorems are true for all values of  $n$ .  
 Cor. 1.  $\frac{x-1}{1} + \frac{x^2-1}{2} + \frac{x^3-1}{3} + \dots + \frac{x^n-1}{n}$  can be expanded in ascending powers of  $x$  in a convergent series the first two terms being  $\frac{5}{2}x + \frac{5}{4}x^2 + \dots$

L. If  $\phi(x) = \frac{x-1}{1} + \frac{x^2-1}{2} + \frac{x^3-1}{3} + \dots + \frac{x^n-1}{n}$  then  
 $\phi(x) + \frac{1}{x-1} + \frac{1}{x-2} + \frac{1}{(x+1)2^{n+1}} + \frac{1}{(x+2)2^{n+2}} + \dots = 0$   
 and hence the values of the series  $\frac{1}{1^2} \cdot \frac{1}{2} + \frac{1}{2^2} \cdot \frac{1}{3} + \dots$

34.  $\frac{x}{1+x} + \frac{1}{3^2} \cdot (\frac{x}{1+x})^2 + \frac{1}{5^2} \cdot (\frac{x}{1+x})^3 + \dots$   
 $= x - \frac{2}{3}(1+\frac{1}{3})x^2 + \frac{2 \cdot 4}{2 \cdot 5} x^3 (\dots + \frac{1}{5}) - \dots$

35. If  $A_n = (1^n + 2^n + 3^n + \dots)(1 + \cos \pi n)$   
 $2^2 + 6^2 + 12^2 + 20^2 + 30^2 + \dots$   
 $= A_n + \frac{1}{2} A_{n+1} + \frac{n(n-1)}{12} A_{n+2} + \frac{n(n-1)(n-2)}{12} A_{n+3} + \dots$

e.g. i.  $\frac{1}{2^2} + \frac{1}{6^2} + \frac{1}{12^2} + \dots = \frac{\pi^2}{8}$   
 ii.  $\frac{1}{2^2} + \frac{1}{6^2} + \frac{1}{12^2} + \dots = 10 - \pi^2$   
 iii.  $\frac{1}{2^2} + \frac{1}{6^2} + \frac{1}{12^2} + \dots = \frac{\pi^4}{6^2} + \frac{10\pi^2}{3} - 35$   
 iv.  $\frac{1}{2^2} + \frac{1}{6^2} + \frac{1}{12^2} + \dots = 126 - \frac{3\pi^2}{2} - \frac{\pi^4}{9}$

1. If any one of  $x, y, z$  be composite integers,

$$\frac{(x+n)(y+n)(z+n)(x+y+z+n)(y+z+u+n)(x+u+v+n)(x+y+u+n)}{x(x+n)(y+n)(z+n)(x+u+n)(y+v+n)(x+y+z+u+n)} \\ = n - (n-2) \frac{n}{u} \cdot \frac{x}{x+n+1} \cdot \frac{y}{y+n+1} \cdot \frac{z}{z+n+1} \cdot \frac{u}{u+n+1} \cdot \frac{x+y+z+u+n+1}{x+y+z+u+n} \\ + (n+4) \frac{n(n+1)}{u} \cdot \frac{x(x-1)}{(x+n+1)(x+n+2)} \cdot \frac{y(y-1)}{(y+n+1)(y+n+2)} \cdot \frac{z(z-1)}{(z+n+1)(z+n+2)} \\ \times \frac{u(u-1)}{(u+n+1)(u+n+2)} \cdot \frac{(x+y+z+u+2n+1)(x+y+z+u+2n+2)}{(x+y+z+u+n)(x+y+z+u+n-1)} + \&c$$

2. If any one of  $x, y, z$  be positive integers,

$$\frac{\frac{1}{x} \frac{x+y+n}{x+n} \frac{y+z+n}{y+n} \frac{z+x+n}{z+n}}{\frac{1}{x+n} \frac{y+n}{y+n} \frac{z+n}{z+n} \frac{x+y+z+n}{x+y+z+n}} = 1 + \frac{xyz}{u(n+1)(x+y+z+n)} \\ + \frac{x(x-1)y(y-1)z(z-1)}{u(n+1)(n+2)(x+y+z+n)(x+y+z+n-1)} + \&c$$

3. If any one of  $x, y, z$  be positive integers,

$$\frac{(x+n)(y+n)(z+n)(x+y+z+n)}{(x+y+n)(y+z+n)(z+x+n)} = n + (n+2) \frac{x}{x+n+1} \cdot \frac{y}{y+n+1} \\ \times \frac{z}{z+n+1} \cdot \frac{x+y+z+n}{x+y+z+n-1} + (n+4) \frac{x(x-1)}{(x+n+1)(x+n+2)} \cdot \frac{y(y-1)}{(y+n+1)(y+n+2)} \\ \times \frac{z(z-1)}{(z+n+1)(z+n+2)} \cdot \frac{(x+y+z+n-1)(x+y+z+n-2)}{(x+y+z+n)(x+y+z+n-1)} + \&c$$

4. If any one of  $x, y, z$  be a positive integer,

$$\geq \frac{1}{x+n} + \geq \frac{1}{y+n} + \geq \frac{1}{z+n} - \geq \frac{1}{x+y+n} - \geq \frac{1}{y+z+n} \\ - \geq \frac{1}{x+z+n} + \geq \frac{1}{x+y+z+n} - \geq \frac{1}{n} \\ = (1 + \frac{1}{n+1}) \cdot \frac{x}{x+n+1} \cdot \frac{y}{y+n+1} \cdot \frac{z}{z+n+1} \cdot \frac{x+y+z+2n+1}{x+y+z+n} \\ + (\frac{1}{2} + \frac{1}{n+2}) \frac{x(x-1)}{(x+n+1)(x+n+2)} \cdot \frac{y(y-1)}{(y+n+1)(y+n+2)} \cdot \frac{z(z-1)}{(z+n+1)(z+n+2)} \\ \times \frac{(x+y+z+2n+1)(x+y+z+2n+2)}{(x+y+z+n)(x+y+z+n-1)} + \&c$$

e.g. If  $x$  is a positive integer

$$1 - 3 \left( \frac{x-1}{x+1} \right)^4 \frac{x-1}{2x-3} + 5 \left( \frac{x-1}{x+1} \cdot \frac{x-2}{x+1} \right)^4 \frac{x-1}{4x-5} \cdot \frac{x}{4x-4} - \&c$$



114. 
$$= \frac{(x(x-2))^2}{(12x-1)^6 |4x-3}$$

ii. 
$$1 - \frac{(x-1)^3}{(x+1)} \frac{3x-1}{3x-3} + \frac{1}{2} \left( \frac{x-1}{x+1} \cdot \frac{x-1}{x+3} \right)^3 \frac{(3x-1)(3x+1)}{(3x-3)(3x-5)} + \dots$$

$$= \frac{1}{2} \leq \frac{1}{x-1} - \frac{3}{2} \leq \frac{1}{x-1} + \frac{1}{2} \leq \frac{1}{2x-1}$$

iii. 
$$1 + 3 \cdot \left( \frac{x-1}{x+1} \right)^3 \frac{3x-1}{3x-3} + 5 \cdot \left( \frac{x-1}{x+1} \cdot \frac{x-2}{x+2} \right)^3 \frac{(3x-1)3x}{(3x-3)(3x-4)} + \dots$$

$$= \left( \frac{x}{2x-1} \right)^3 (3x-2)$$

iv. 
$$1 + \left( \frac{x}{11} \right)^2 \frac{x}{3x} + \left\{ \frac{x(x-1)}{12} \right\}^2 \frac{x(x-1)}{3x(3x-2)} + \dots = \left( \frac{12x}{11} \right)^3 |3x$$

v. 
$$1 + \frac{x}{11} \cdot \frac{x-1}{x+1} \cdot \frac{x}{4x-1} + \frac{x(x-1)}{12} \cdot \frac{(x-1)(x-2)}{(x+1)(x+2)} \cdot \frac{x(x-1)}{(4x-1)(4x-2)} + \dots$$

$$= \frac{8}{9} \left( \frac{3x}{2x} \right)^3 \frac{|x}{|4x}$$

5. 
$$n \frac{x+n}{1x} \frac{y+n}{x+y+n} \frac{z+n}{y+z+n} \frac{x+y+z+n}{z+x+n} = x - (n+2) \frac{x}{11} \frac{y}{(x+n+1)(y+n+1)}$$

$$\times \frac{z}{z+n+1} + (n+4) \frac{n(n+1)}{12} \frac{x(x-1)}{(x+n+1)(x+n+2)} \cdot \frac{y(y-1)}{(y+n+1)(y+n+2)} \times$$

$$\frac{z}{z(n-1)} - \dots$$

$$\frac{(x+n+1)(z+n+2)}{(x+n+1)(z+n+2)}$$

6. If  $a + \beta + \gamma + 1 = n$ , then

$$(n+1) \frac{1}{10} \frac{|a| |b| |c|}{|n-a| |n-b| |n-c|} + (n+3) \frac{1}{11} \frac{|a+1| |b+1| |c+1|}{|n-a+1| |n-b+1| |n-c+1|}$$

$$+ (n+5) \frac{1}{12} \frac{|a+2| |b+2| |c+2|}{|n-a+2| |n-b+2| |n-c+2|} + \dots$$

(when  $k = \infty$ )  $= - \leq \frac{1}{x} \leq \frac{1}{y} = \dots + \dots$

Cor. 
$$\frac{x^2}{4} \left\{ 1 + 5 \left( \frac{1}{2} \right)^4 (1-x) + 9 \left( \frac{1.3}{2.4} \right)^4 (1-x)^2 + 13 \left( \frac{1.3.5}{2.4.6} \right)^4 (1-x)^3 + \dots \right\}$$

$$+ \log x = 3 \log 2, \text{ when } x \text{ vanishes.}$$

7. 
$$1 + \frac{n}{11} \frac{x}{x+n+1} \frac{y}{y+n+1} \frac{z}{z+n} \frac{x(n+1)}{12} \frac{x(x-1)}{(x+n+1)(x+n+2)} \times$$

$$\frac{y(y-1)}{(y+n+1)(y+n+2)} + \dots = \frac{|x+n| |y+n|}{n |x+y+z|} \frac{|z| |x+y+z|}{|2+z|}$$

$$= 1 \leq \frac{1}{x+n} + 2 \frac{1}{y+n} = \frac{1}{x+y+n} = \frac{1}{n} \frac{x(x-1)}{x(x-1)}$$

$$= \left(1 + \frac{1}{n+1}\right) \frac{x}{x+n+1} \cdot \frac{1}{y+n+1} + \left(\frac{1}{2} + \frac{1}{n+2}\right) \frac{x(x-1)}{(x+n+1)(x+n+2)} \times$$

$$\frac{y(y-1)}{(y+n+1)(y+n+2)} + \dots$$

$$2. \quad n \frac{\frac{x+y}{x} \frac{y+n}{y}}{\frac{x}{x} \frac{y}{y}} \cdot \frac{\frac{1}{n} \frac{y+n}{y+n}}{\frac{1}{n} \frac{x+y+n}{x+y+n}} = n + (n+2) \frac{n^2}{(n)^2} \frac{x y}{(x+n+1)(y+n+1)}$$

$$+ (n+4) \frac{n^2(n+1)^2}{(n)^2} \frac{x(x-1)}{(x+n+1)(x+n+2)} \frac{y(y-1)}{(y+n+1)(y+n+2)} + \dots$$

$$3. \quad \frac{(x+n)(y+n)}{x+y+n} = n + (n+2) \frac{x y}{(x+n+1)(y+n+1)} +$$

$$(n+4) \frac{x(x-1)}{(x+n+1)(x+n+2)} \frac{y(y-1)}{(y+n+1)(y+n+2)} + \dots$$

$$4. \quad n \frac{\frac{x+n}{n} \frac{y+n}{y}}{\frac{1}{n} \frac{x+y+n}{x+y+n}} \cdot \frac{\frac{1}{n} \frac{x+y+n}{x+y+n}}{\frac{1}{n} \frac{x+y+n}{x+y+n}} = n +$$

$$(n+2) \frac{n}{n} \frac{x y}{(x+n+1)(y+n+1)} + (n+4) \frac{n(n+1)}{n} \frac{x(x-1)}{(x+n+1)(x+n+2)}$$

$$\frac{y(y-1)}{(y+n+1)(y+n+2)} + \dots$$

$$5. \quad n \frac{\frac{x+n}{n} \frac{y+n}{y}}{\frac{1}{n} \frac{x+y+n}{x+y+n}} = n - (n+2) \frac{n}{n} \frac{x y}{(x+n+1)(y+n+1)} +$$

$$(n+4) \frac{n(n+1)}{n} \frac{x(x-1)}{(x+n+1)(x+n+2)} \frac{y(y-1)}{(y+n+1)(y+n+2)} - \dots$$

$$6. \quad \left\{ (n+1)^2 + \frac{1}{(n+2)^2} + \frac{1}{(n+3)^2} + \dots \right\} - \left\{ \frac{1}{(x+n+1)^2} + \frac{1}{(x+n+2)^2} + \dots \right\}$$

$$= \left(1 + \frac{1}{n+1}\right) \frac{x}{x+n+1} \cdot \frac{1}{n+1} - \left(\frac{1}{2} + \frac{1}{n+2}\right) \frac{1}{(n+1)(n+2)} \times$$

$$\frac{x(x-1)}{(x+n+1)(x+n+2)} + \dots$$

$$7. \quad \frac{\frac{x+n}{x} \frac{x-n}{x}}{(x)^2} \cdot \frac{\sin \pi x}{\pi} = n - (n+2) \frac{n^3}{(n)^3} \frac{x}{x+n+1} +$$

$$(n+4) \frac{n^3(n+1)^3}{(n)^3} \frac{x(x-1)}{(x+n+1)(x+n+2)} - \dots$$

$$8. \quad \frac{\frac{x+n}{x} \frac{1}{n}}{\frac{1}{n} \frac{x-n}{x}} = 1 - \frac{n^2}{(n)^2} \frac{x}{x+n+1} +$$

116. 
$$+ \frac{n^2(n+1)^2}{(L)^4} \frac{x(x-1)}{(x+n+1)(x+n+2)} - \&c.$$

9. 
$$n \frac{x - \frac{n+1}{L}}{x - \frac{n+1}{L}} \cdot \frac{x+n}{x + \frac{n-1}{L}} \frac{\frac{n-1}{L}}{L} = n - (n+2) \frac{n^2}{(L)^2} \frac{x}{x+n+1} + (n+4) \frac{n^2(n+1)^2}{(L)^2} \frac{x(x-1)}{(x+n+1)(x+n+2)} + \&c.$$

10. 
$$\frac{n(x+n)}{LxLn} = n + (n+2) \frac{n^2}{(L)^2} \frac{x}{x+n+1} + \&c.$$

11. 
$$\frac{LxLn \frac{(n-1)^2}{L^2}}{nLn \left(x + \frac{n-1}{L}\right)^2} = \frac{1}{n} - \frac{n}{L} \cdot \frac{x}{x+n+1} \cdot \frac{1}{n+2} + \frac{n(n+1)}{L^2} \frac{x(x-1)}{(x+n+1)(x+n+2)} - \&c.$$

12. 
$$\frac{LxLn \frac{x+n}{L}}{LxLn \frac{x+n}{L}} = 1 - \frac{n}{L} \cdot \frac{x}{x+n+1} + \frac{n(n+1)}{L^2} \frac{x(x-1)}{(x+n+1)(x+n+2)} - \&c.$$

13. 
$$\frac{x+n}{Ln} \frac{\frac{n-1}{L}}{x + \frac{n-1}{L}} = 1 + \frac{n}{L} \cdot \frac{x}{x+n+1} + \frac{n(n+1)}{L^2} \frac{x(x-1)}{(x+n+1)(x+n+2)} + \&c.$$

14. 
$$\frac{x+n}{(n-1)L} \frac{\frac{n-1}{L}}{x + \frac{n-1}{L}} = n + (n+2) \frac{n}{L} \cdot \frac{x}{x+n+1} + (n+4) \frac{n(n+1)}{L^2} \frac{x(x-1)}{(x+n+1)(x+n+2)} + \&c.$$

15. 
$$\frac{(L)^2}{Lx} \frac{\sin \pi n \tan \pi n}{\pi^2 n} = n + (n+2) \frac{n^2}{(L)^2} + (n+4) \frac{n^2(n+1)^2}{(L)^4} + \&c.$$

16. 
$$n + (n+2) \frac{x^2}{(L)^2} + (n+4) \frac{n^2(n+1)^2}{(L)^4} + \&c = \frac{\left(\frac{n-1}{L}\right)^2 \left(\frac{n-1}{L}\right)^2 \frac{\sin \pi n}{\pi}}{\left(\frac{n-1}{L}\right)^2}$$

17. 
$$\frac{\sin \pi n}{\pi} = n - (n+2) \frac{x^2}{(L)^2} + (n+4) \frac{n^2(n+1)^2}{(L)^4} - \&c.$$

18. 
$$\frac{\left(\frac{n-1}{L}\right)^2}{(L)^2} \frac{2 \tan \pi n}{\pi n^2} = \frac{1}{n} + \frac{x^2}{(L)^2} \frac{1}{n+2} + \frac{n^2(n+1)^2}{(L)^4} \frac{1}{n+4} + \&c.$$

19. 
$$\frac{\pi \left(\frac{n-1}{L}\right)^2}{2nLn \sin \frac{\pi}{2}} = \frac{1}{n} + \frac{n}{L} \cdot \frac{1}{(n+1)^2} + \frac{n(n+1)}{L^2} \cdot \frac{1}{(n+2)^2} + \&c.$$

20. 
$$\sum \frac{1}{x+n} - \sum \frac{1}{n} = \left(1 + \frac{1}{n+1}\right) \frac{n}{L} \cdot \frac{x}{x+n+1} + \left(\frac{1}{2} + \frac{1}{n+1}\right) \frac{x(x-1)}{(x+n+1)(x+n+2)} - \&c.$$

21. 
$$\sum \frac{1}{x} + \sum \frac{1}{n} - \sum \frac{1}{x+n} = \left(1 + \frac{1}{n+1}\right) \frac{n}{L} \cdot \frac{x}{x+n+1} - \&c.$$

$$\left(\frac{1}{2} + \frac{1}{n+1}\right) \frac{x(n+1)}{L} \cdot \frac{x(x-1)}{(x+n+1)(x+n+2)} + \dots$$

$$21. = \left\{ \frac{1}{n^2} + \frac{1}{(n+1)^2} + \frac{1}{(n+2)^2} + \frac{1}{(n+3)^2} + \dots \right\}$$

$$= \left(1 + \frac{1}{n}\right) + \left(\frac{1}{2} + \frac{1}{n+1}\right) \left(\frac{1}{n+1}\right)^2 + \left(\frac{1}{3} + \frac{1}{n+2}\right) \left(\frac{1}{n+1} \cdot \frac{1}{n+2}\right)^2 + \dots$$

$$22. \left\{ \frac{1}{(1+\frac{x}{2})^2} + \frac{1}{(2+\frac{x}{2})^2} + \frac{1}{(3+\frac{x}{2})^2} + \dots \right\} - \left\{ \frac{1}{(1+x)^2} + \frac{1}{(2+x)^2} + \dots \right\}$$

$$= \left(1 - \frac{1}{n+1}\right) \frac{1}{n+1} + \left(\frac{1}{2} - \frac{1}{n+1}\right) \frac{1}{(n+1)(n+2)} + \dots$$

$$24. = \frac{1}{n} + \dots = \frac{1}{n+1} = \left(1 + \frac{1}{n+1}\right) \frac{x^2}{(L)^2} + \left(\frac{1}{2} - \frac{1}{n+1}\right) \frac{x^2(n+1)^2}{(L)^2} + \dots$$

$$27. 1 - \frac{(x)^3 \sqrt{x-1}}{(Lx-1)^3} = 1 - 3 \cdot \left(\frac{x-1}{x+1}\right)^2 + 5 \cdot \left(\frac{x-1}{x+1} \cdot \frac{x-1}{x+2}\right)^3 - \dots$$

$$2. \frac{x^L}{Lx-1} = 1 + 3 \cdot \left(\frac{x-1}{x+1}\right)^2 + 5 \cdot \left(\frac{x-1}{x+1} \cdot \frac{x-1}{x+2}\right)^2 + \dots$$

$$3. \frac{(x)^4 \sqrt{x}}{(Lx)^2} \cdot \frac{Lx}{4x-1} = 1 + \left(\frac{x-1}{x+1}\right) + \left(\frac{x-1}{x+1} \cdot \frac{x-1}{x+2}\right)^2 + \dots$$

$$4. \frac{(Lx)^L}{Lx-1} = 1 - 3 \cdot \left(\frac{x-1}{x+1}\right)^2 + 5 \cdot \left(\frac{x-1}{x+1} \cdot \frac{x-1}{x+2}\right)^2 - \dots$$

$$5. x = 1 + 3 \cdot \frac{x-1}{x+1} + 5 \cdot \frac{x-1}{x+1} \cdot \frac{x-1}{x+2} + \dots$$

$$6. \frac{\sqrt{x}}{2} \cdot \frac{Lx}{Lx-1} = 1 + \frac{x-1}{x+1} + \frac{(x-1)(x-1)}{(x+1)(x+1)} + \dots$$

$$7. \frac{x}{Lx-1} = 1 - \frac{x-1}{x+1} + \frac{(x-1)(x-1)}{(x+1)(x+2)} + \dots$$

$$8. 1 - 3 \cdot \frac{x-1}{x+1} + 5 \cdot \frac{x-1}{x+1} \cdot \frac{x-1}{x+2} - \dots = 0.$$

$$9. 2 \pm \frac{1}{x+1} + \frac{(x-1)(x-1)}{(Lx-1)^2} = 1 + \frac{1}{x+1} + \frac{1}{3} \cdot \frac{x-1}{x+1} \cdot \frac{x-1}{x+2} + \dots$$

$$10. 2 \pm \frac{1}{2x} - \frac{1}{2} + \frac{1}{2x} = 1 - \frac{1}{2} \cdot \frac{x-1}{x+1} + \frac{1}{3} \cdot \frac{x-1}{x+1} \cdot \frac{x-1}{x+2} - \dots$$

$$11. \frac{2^{4x} (Lx)^4}{4x(Lx)^2} = 1 - \frac{1}{3} \cdot \frac{x-1}{x+1} + \frac{1}{5} \cdot \frac{x-1}{x+1} \cdot \frac{x-1}{x+2} - \dots$$

14.  $\frac{1}{2} (\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{x^2}) + \frac{1}{2} (\frac{1}{x+1} + \frac{1}{x+2} + \dots + \frac{1}{x+x})$

$= 1 - \frac{1}{2} \cdot \frac{x-1}{x+1} + \frac{1}{3^2} \cdot \frac{x-1}{x+1} \cdot \frac{x-2}{x+2} - \dots$

13.  $x(x-3) = 1^3 + 3^3 \cdot \frac{x-1}{x+1} + 5^3 \cdot \frac{x-1}{x+1} \cdot \frac{x-2}{x+2} + \dots$

14.  $\frac{1}{\pi} = 1 - 5 \cdot (\frac{1}{2})^3 + 9 \cdot (\frac{1 \cdot 3}{2 \cdot 4})^3 - 13 \cdot (\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6})^3 + \dots$

15.  $1 + 9 \cdot (\frac{1}{4})^4 + 17 \cdot (\frac{1 \cdot 3}{2 \cdot 4})^4 + \dots = \frac{2\sqrt{3}}{\sqrt{\pi}} \cdot (\frac{1}{2})^2$

16.  $1 + (\frac{1}{2})^2 \cdot \frac{1}{3} + (\frac{1 \cdot 3}{2 \cdot 4})^2 \cdot \frac{1}{9} + \dots = \frac{\pi^2}{4} \cdot (\frac{1}{2})^4$

17.  $1 + \frac{1}{2} \cdot \frac{1}{5^2} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{9^2} + \dots = \frac{\pi^2}{8\sqrt{2}} \cdot \frac{\sqrt{\pi}}{(\frac{1}{2})^2}$

18.  $1 + (\frac{1}{2})^3 + (\frac{1 \cdot 3}{2 \cdot 4})^3 + (\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6})^3 + \dots = (\frac{\pi}{(\frac{1}{2})^4})^2$

19.  $1 - (\frac{1}{2})^2 + (\frac{1 \cdot 3}{2 \cdot 4})^2 - (\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6})^2 + \dots = \frac{\sqrt{\pi/2}}{(\frac{1}{2})^2} \cdot \frac{6 \cdot (\frac{\pi}{2})^3 \sin \pi n \sin \frac{\pi n}{2}}{\pi^2 n^2 (1+2 \cos \pi n) (\frac{1}{2})^2}$

20.  $1 + (\frac{n}{a})^3 + \{ \frac{n(n+1)}{2} \}^3 + \{ \frac{n(n+1)(n+2)}{6} \}^3 + \dots = \frac{\pi^2 n^2 (1+2 \cos \pi n) (\frac{1}{2})^2}{\dots}$

8.  $\frac{12x}{x+n} \frac{1}{y+z} = 1 + \frac{x}{y} \cdot \frac{y}{x+1} + \frac{x(x-1)}{2} \cdot \frac{y(y-1)}{(x+1)(x+2)} + \dots$

Sol. Write  $-n+m$  for  $2$  in  $\S 5$  and make  $x$  infinite or equate the coeffts. of  $u^n$  in  $(1+u)^{n+m} (1+\frac{1}{2}u)^m = \frac{(1+u)^{n+y+z}}{u^x}$

9.  $\frac{1}{\alpha-1} - \frac{1}{\alpha-\beta-1} = \frac{\beta}{\alpha} + \frac{\beta(\beta+1)}{\alpha(\alpha+1)} \cdot \frac{1}{2} + \frac{\beta(\beta+1)(\beta+2)}{\alpha(\alpha+1)(\alpha+2)} \cdot \frac{1}{3} + \dots$

10.  $\frac{12x}{x} \frac{1}{x} = \frac{1}{x} - \frac{x}{2} \cdot \frac{1}{x+1} + \frac{x(x-1)}{2} \cdot \frac{1}{x+2} - \dots$

Ex. 1.  $\frac{1}{(n+1)^2} + \frac{1}{(n+2)^2} + \frac{1}{(n+3)^2} + \frac{1}{(n+4)^2} + \dots$   
 $= \frac{1}{n+1} + \frac{1}{2(n+1)(n+2)} + \frac{1}{3(n+1)(n+2)(n+3)} + \dots$

2.  $\frac{\pi}{\sin \pi n} = \frac{1}{n} + \frac{\pi}{2} \cdot \frac{1}{n+1} + \frac{\pi(n+1)}{2} \cdot \frac{1}{n+2} + \dots$

3.  $\frac{\sqrt{\pi} \pi}{\sqrt{n+\frac{1}{2}}} = \frac{1}{n+1} + \frac{1}{2} \cdot \frac{1}{n+2} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{n+3} + \dots$

4.  $\frac{\sqrt{\pi} \Gamma_n}{2 \Gamma_{n+\frac{1}{2}}} = 1 - \frac{n}{2} \cdot \frac{1}{3} + \frac{n(n-1)}{2!} \cdot \frac{1}{5} - \frac{n(n-1)(n-2)}{3!} + \dots$

5.  $\frac{\sqrt{\pi} \Gamma_{n-1}}{\Gamma_{n+\frac{1}{2}}} (= \frac{1}{x+n} - \frac{1}{n+1}) = \frac{1}{n^2} - \frac{x}{2!} \cdot \frac{1}{(n+1)^2} + \frac{x(x-1)}{2!} \cdot \frac{1}{(n+2)^2} - \dots$

6.  $\frac{\sqrt{\pi} \Gamma_n}{\Gamma_{n+\frac{1}{2}}} (= \frac{1}{n+\frac{1}{2}} - \frac{1}{n}) = \frac{1}{(n+\frac{1}{2})^2} + \frac{1}{2} \cdot \frac{1}{(n+\frac{1}{2})^2} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{(n+\frac{1}{2})^2} + \dots$

7.  $-\frac{\pi}{\sin \pi n} \approx \frac{1}{n-1} = \frac{1}{n^2} + \frac{n}{2!} \cdot \frac{1}{(n+1)^2} + \frac{n(n+1)}{2!} \cdot \frac{1}{(n+2)^2} + \dots$

11.  $\alpha^n = \{ \alpha^n - (\beta+1)^n \} + \{ (\alpha+1)^n - (\beta+2)^n \} \left( \frac{\beta+1}{\alpha+1} \right)^n + \{ (\alpha+2)^n - (\beta+3)^n \} \left( \frac{\beta+1}{\alpha+1} \cdot \frac{\beta+2}{\alpha+2} \right)^n + \dots$

Cor. 1.  $\frac{\beta}{\alpha - \beta - 1} = \frac{\pi}{x} + \frac{\beta(\beta+1)}{\alpha(\alpha+1)} + \frac{\beta(\beta+1)(\beta+2)}{\alpha(\alpha+1)(\alpha+2)} + \dots$

2.  $\frac{\beta^2}{\alpha - \beta - 1} = (\alpha + \beta + 1) \left( \frac{\beta}{\alpha} \right)^2 + (\alpha + \beta + 3) \left( \frac{\beta}{\alpha} \cdot \frac{\beta+1}{\alpha+1} \right)^2 + \dots$

12. If  $e^{A_1 x + A_2 \frac{x^2}{2} + A_3 \frac{x^3}{3} + \dots} = P_0 + P_1 x + P_2 x^2 + \dots$ , then

$P_n = P_{n-1} A_1 + P_{n-2} A_2 + P_{n-3} A_3 + \dots$  to  $n$  terms and  $P_0 = 1$

and consequently if  $S_n = a_1^n + a_2^n + a_3^n + \dots + a_m^n$  and

$P_n$  denote the sum of the products of  $a_1, a_2, a_3, \dots, a_m$

taken  $n$  at a time then  $a P_n = P_{n-1} S_1 - P_{n-2} S_2 + P_{n-3} S_3 -$

$P_{n-4} S_4 + \dots$  and  $P_n = 1$ .

13.  $\frac{1}{n(n+1)} = \frac{x}{2} \cdot \frac{1}{(n+1)^{n+1}} + \frac{x(x-1)}{2!} \cdot \frac{1}{(n+2)^{n+1}} - \dots = \frac{1 \cdot 1 \cdot 1 \cdot x}{1 \cdot 2 \cdot 2} \phi(x)$

where  $\phi(0) = 1$  and  $a \phi(x) = S_1 \phi(x-1) + S_2 \phi(x-2) + S_3 \phi(x-3) + \dots$

to  $n$  terms where  $S_n = \frac{1}{na} - \frac{1}{(x+n)^2} + \frac{1}{(n+1)^2} - \frac{1}{(x+n+2)^2} + \dots$

Cor. 1.  $1 + \frac{1}{2} \cdot \frac{1}{3(n+1)} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{5(n+1)} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{1}{7(n+1)} + \dots = \frac{\pi}{2} \phi(x)$

where  $\phi(0) = 1$  and a  $\phi(n) = S_1 \phi(n-1) + S_2 \phi(n-2) + S_3 \phi(n-3) + \dots$   
 to  $n$  terms where  $S_2 = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$

Cor. 2.  $\frac{1}{2} \log 11 + \frac{1}{2} \cdot \frac{1}{2} \log 11 + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{6} \log 11 + \dots = \phi(n)$  where  $\phi(0) = 1$   
 and a  $\phi(n) = S_1 \phi(n-1) + S_2 \phi(n-2) + \dots$  where  $S_n = \frac{1}{2^n} - \frac{1}{3^n} + \frac{1}{4^n}$   
 $- \frac{1}{5^n} + \dots$  e.g.  $1 + \frac{1}{2} \cdot \frac{1}{3} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{5} + \dots = \frac{11^2}{288} + \frac{\pi}{6} (\log 2)^2$ .

Ex.  $\int_0^{\frac{\pi}{2}} \theta \cot \theta \log \sin \theta \, d\theta = -\frac{\pi^3}{48} - \frac{\pi}{4} (\log 2)^2$ .

14.  $\frac{1}{(x+1)^n} + \frac{1+\frac{1}{2}}{(x+2)^n} + \frac{1+\frac{1}{2}+\frac{1}{3}}{(x+3)^n} + \frac{1+\frac{1}{2}+\frac{1}{3}+\frac{1}{4}}{(x+4)^n} + \dots$   
 $= \frac{\pi}{2} S_{n+1} - (S_1 S_n + S_2 S_{n-1} + S_3 S_{n-2} + \dots)$  the last term  
 being  $S_{\frac{n}{2}} S_{\frac{n+1}{2}}$  or  $\frac{1}{2} S_{\frac{n}{2}} S_{\frac{n+1}{2}}$  according as  $n$  is even or  
 odd) where  $S_n = \frac{1}{2^n} + \frac{1}{(x+1)^n} + \frac{1}{(x+2)^n} + \dots$  and  
 $S_1 = -\sum \frac{1}{x-1}$ .

Sol.  $1(\frac{1}{2} - \frac{1}{n+2}) + (1+\frac{1}{2})(\frac{1}{3} - \frac{1}{n+3}) + (1+\frac{1}{2}+\frac{1}{3})(\frac{1}{4} - \frac{1}{n+4}) + \dots$   
 $= \frac{1}{2} \left\{ (1+\frac{1}{2}+\frac{1}{3}+\dots+\frac{1}{n})^2 + (\frac{1}{2}+\frac{1}{3}+\dots+\frac{1}{n-2}) \right\}$ .

In the above identity write  $n+2=1$  form and equate  
 the coeff<sup>s</sup> of  $n^2$ .

15.  $\frac{1 \cdot 1 \cdot 1}{1 \cdot 1 \cdot 1} + \frac{1 \cdot 1 \cdot 1}{1 \cdot 1 \cdot 1} + \frac{1 \cdot 1 \cdot 1}{1 \cdot 1 \cdot 1} + \dots$  to  $n$  terms  
 $= \log x$  (when  $n = \infty$ )  $= - = \frac{1}{x} = \frac{1}{1} + C_0$ .

Cor.  $\pi \left\{ 1 + \left(\frac{1}{2}\right)^x (1-x) + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x (1-x)^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^x (1-x)^3 + \dots \right\}$   
 $+ \log x = \frac{1}{2} \log 2$  when  $x=0$ .

16. If  $A_0 - n A_1 + \frac{n(n-1)}{2} A_2 - \frac{n(n-1)(n-2)}{6} A_3 + \dots = P_n$ , then  
 $P_0 - n P_1 + \frac{n(n-1)}{2} P_2 - \frac{n(n-1)(n-2)}{6} P_3 + \dots = A_n$ .

$$17. \frac{A_0}{x^2} + \frac{n}{1} \cdot \frac{A_1}{x^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{A_2}{x^{n+2}} + \dots$$

$$= \frac{A_0}{(x+h)^2} + \frac{n}{1} \cdot \frac{A_1 + h A_0}{(x+h)^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{A_2 + 2h A_1 + h^2 A_0}{(x+h)^{n+2}} + \dots$$

$$18. \text{ If } \frac{A_0}{x^n} + \frac{n}{1} \cdot \frac{A_1}{x^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{A_2}{x^{n+2}} + \dots$$

$$= \frac{A_0}{(x-1)^n} + \frac{n}{1} \cdot \frac{A_1}{(x-1)^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{A_2}{(x-1)^{n+2}} - \dots, \text{ then}$$

$$i. e^x = \frac{A_0 + \frac{x}{1} A_1 + \frac{x^2}{1^2} A_2 + \frac{x^3}{1^3} A_3 + \dots}{A_0 - \frac{x}{1} A_1 + \frac{x^2}{1^2} A_2 - \frac{x^3}{1^3} A_3 + \dots}$$

$$ii. \frac{1}{\{\phi(x)\}^n} \left[ A_0 + A_1 \frac{x}{1} \left\{ \frac{\phi(x) - \phi(-x)}{\phi(x)} \right\} + A_2 \frac{x^2}{1^2} \left\{ \frac{\phi(x) - \phi(-x)}{\phi(x)} \right\}^2 + \dots \right]$$

is always an even function of  $x$  whatever be  $\phi(x)$ .

iii. If  $n$  is even, the value of  $A_{n+1}$  depends upon the value of  $A_n$ ; but we may give for  $A_n$  any value we choose.

$$\frac{A_{n-1}}{2} = \frac{n-1}{1^2} (2^2-1) B_2 A_{n-2} - \frac{(n-1)(n-3)(n-5)}{1^4} (2^4-1) B_4 A_{n-4} - \dots$$

$$+ \frac{(n-1)(n-3)(n-5)(n-7)(n-9)}{1^6} (2^6-1) B_6 A_{n-6} - \dots$$

$$19. \frac{1}{x^n} + \frac{n}{1} \cdot \frac{m}{x^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{m(m-1)}{x^{n+2}} + \dots$$

$$= \frac{1}{(x-1)^n} + \frac{n}{1} \cdot \frac{m}{(x-1)^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{(m-n)(m-n-1)}{n(n+1)} \cdot \frac{1}{(x-1)^{n+2}} + \dots$$

$$20. \phi(0) + \frac{m}{n} \cdot \frac{\phi'(0)}{1} + \frac{m(m-1)}{n(n-1)} \cdot \frac{\phi''(0)}{1^2} + \dots$$

$$= \phi(1) + \frac{m-n}{n} \cdot \frac{\phi'(1)}{1} + \frac{(m-n)(m-n-1)}{n(n-1)} \cdot \frac{\phi''(1)}{1^2} + \dots$$

$$21. e^x = \frac{1 + \frac{mx}{n} + \frac{m(m-1)}{n(n-1)} \cdot \frac{x^2}{1^2} + \dots}{1 + \frac{m-m}{n} \cdot \frac{x}{1} + \frac{(m-n)(m-n-1)}{n(n-1)} \cdot \frac{x^2}{1^2} + \dots}$$

$$22. \frac{1}{(2x)^2} + \frac{n}{1} \cdot \frac{2m}{2m} \cdot \frac{1}{(x+1)^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{m(m-1)}{2m(2m-1)} \cdot \frac{1}{(x+1)^{n+2}} + \dots$$

$$= \frac{1}{x^2} - \frac{n}{1} \cdot \frac{m}{2m} \cdot \frac{1}{x^{n+1}} + \frac{n(n-1)}{1^2} \cdot \frac{m(m-1)}{2m(2m-1)} \cdot \frac{1}{x^{n+2}} - \dots$$



$$23. e^x = \frac{1 + \frac{m}{1m} \cdot \frac{x}{1} + \frac{m(m+1)}{2m(1m+1)} \cdot \frac{x^2}{1^2} + \frac{m(m+1)(m+2)}{2m(1m+1)(1m+2)} \cdot \frac{x^3}{1^3} + \dots}{1 - \frac{m}{1m} \cdot \frac{x}{1} + \frac{m(m+1)}{2m(1m+1)} \cdot \frac{x^2}{1^2} - \frac{m(m+1)(m+2)}{2m(1m+1)(1m+2)} \cdot \frac{x^3}{1^3} + \dots}$$

Case 1.  $e^x = \frac{1 + \frac{1}{2} \cdot \frac{x}{1} + \frac{1 \cdot 2}{2 \cdot 2} \cdot \frac{x^2}{1^2} + \frac{1 \cdot 2 \cdot 3}{1 \cdot 2 \cdot 6} \cdot \frac{x^3}{1^3} + \dots}{1 - \frac{1}{2} \cdot \frac{x}{1} + \frac{1 \cdot 2}{2 \cdot 2} \cdot \frac{x^2}{1^2} - \frac{1 \cdot 2 \cdot 3}{1 \cdot 2 \cdot 6} \cdot \frac{x^3}{1^3} + \dots}$

$$2. 1 - \left(\frac{1}{2}\right)^x \frac{x}{1-x} + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)^x \frac{x^2}{(1-x)^2} - \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^x \frac{x^3}{(1-x)^3} + \dots$$

$$= \sqrt{1-x} \left\{ 1 + \left(\frac{1}{2}\right)^x x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)^x x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^x x^3 + \dots \right\}$$

$$24. \frac{1}{n(n-1)} + \frac{m}{1} \cdot \frac{1}{(n+1)(n-1)} + \frac{m(m+1)}{1^2} \cdot \frac{1}{(n+2)(n-1)} + \dots$$

$$= \frac{1}{n(n-1)^n} + \frac{m-1}{1} \cdot \frac{1}{(n+1)(n-1)^{n+1}} + \frac{(m-2)(m-3)}{1^2} \cdot \frac{1}{(n+2)(n-1)^{n+2}}$$

$$25. \frac{1}{2x} + \frac{1}{n(n+1)x^2} + \frac{1}{n(n+1)(n+2)x^3} + \dots$$

$$= \frac{1}{n(n-1)} - \frac{1}{(n+1)(n-1)^2} + \frac{1}{(n+2)(n-1)^3} - \dots$$

$$26. (1-x)^{\alpha+\beta} \left\{ 1 + \frac{\alpha}{1} \cdot \frac{\beta}{x} x + \frac{\alpha(\alpha+1)}{1^2} \cdot \frac{\beta(\beta+1)}{2(x+1)} x^2 + \dots \right\}$$

$$= (1-x)^{\alpha} \left\{ 1 + \frac{(x-\alpha)(x-\beta)}{1 \cdot x} x + \frac{(x-\alpha)(x-\alpha+1)(x-\beta)(x-\beta+1)}{1^2 \cdot x(x+1)} x^2 + \dots \right\}$$

$$27. \frac{1+q+n}{1+x} + \frac{p}{1} \frac{1+q+n+1}{1+x+1} + \frac{p(p-1)q(q-1)}{1^2} x$$

$$\frac{1+q+n+2}{1+x+2} + \dots = \frac{1+p+n}{1+x} + \frac{xy}{1} \frac{1+p+q+n+1}{1+p+1} \frac{1+q+n+1}{1+q+1}$$

$$+ \frac{x(x-1)y(y-1)}{1^2} \frac{1+p+q+n+2}{1+p+2} \frac{1+q+n+2}{1+q+2} + \dots$$

$$28. \frac{1}{p+n} + \frac{x}{1} \cdot \frac{y}{n} \cdot \frac{1}{p+n+1} + \frac{x(x-1)}{1^2} \cdot \frac{y(y-1)}{n(n-1)} \cdot \frac{1}{p+n+2} + \dots$$

$$= \frac{1+n-1}{1+x} \frac{1+q+n}{1+y} - p \frac{1+n-1}{1+x+1} \frac{1+q+n+1}{1+y+1} + p(p-1) x$$

$$\frac{1+n-1}{1+x+2} \frac{1+q+n+2}{1+y+2} - \dots$$

$$\begin{aligned}
29. \frac{\pi}{4} & \left\{ \frac{1}{n+1} + \binom{n}{2}^{-1} \frac{1}{n+2} + \binom{1 \cdot 3 \cdot 5}{2 \cdot 4}^{-1} \frac{1}{n+3} + \dots \right\} \\
&= 1 - \frac{\pi}{4} \cdot \left(\frac{2}{3}\right)^n + \frac{n(n-1)}{2!} \cdot \left(\frac{2 \cdot 4}{3 \cdot 5}\right)^n - \frac{n(n-1)(n-2)}{3!} \cdot \left(\frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7}\right)^n + \dots \\
&= \frac{\pi}{4} \cdot \left(\frac{1}{n+2}\right)^n \left\{ 1 + \binom{n}{2}^{-1} + \binom{1 \cdot 3 \cdot 5}{2 \cdot 4}^{-1} + \dots \text{ to } n+1 \text{ terms} \right\} \\
&= \frac{1}{2n+1} \left\{ \frac{n+\frac{1}{2}}{n+1} \cdot 1 + \frac{n+\frac{1}{2}}{n+1} \cdot \frac{n+\frac{1}{2}}{n+2} \cdot \frac{1}{3} + \frac{n+\frac{1}{2}}{n+1} \cdot \frac{n+\frac{1}{2}}{n+2} \cdot \frac{n+\frac{1}{2}}{n+3} \cdot \frac{1}{5} \right. \\
&\quad \left. + \dots \right\} \\
&= \frac{\sqrt{\pi}}{2} \cdot \frac{1}{n+\frac{1}{2}} \left\{ 1 - \frac{\pi}{4} \cdot \frac{1}{2} \cdot \frac{1}{3} + \frac{n(n-1)}{2!} \cdot \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1}{5} - \dots \right\}
\end{aligned}$$

Case 1.  $\frac{\pi}{4} \left\{ 1 + \binom{n}{2}^{-1} \frac{1}{3} + \binom{1 \cdot 3 \cdot 5}{2 \cdot 4}^{-1} \frac{1}{5} + \binom{1 \cdot 3 \cdot 5 \cdot 7}{2 \cdot 4 \cdot 6}^{-1} \frac{1}{7} + \dots \right\}$

$$= \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \frac{1}{7^2} + \dots$$

2.  $\pi n \left\{ \frac{1}{n} + \binom{n}{2}^{-1} \frac{1}{n+1} + \binom{1 \cdot 3 \cdot 5}{2 \cdot 4}^{-1} \frac{1}{n+2} + \dots \right\} = (1 + \frac{1}{2} + \dots + \frac{1}{n})$

=  $4 \log 2$  when  $n$  becomes infinite.

30.  $\frac{1}{y+n} = \frac{x}{n} \cdot \frac{1}{y+n+1} + \frac{x(x-1)}{n(n+1)} \cdot \frac{1}{y+n+2} - \dots$

$$= \frac{1}{x+n} - \frac{y}{n} \cdot \frac{1}{x+n+1} + \frac{y(y-1)}{n(n+1)} \cdot \frac{1}{x+n+2} - \dots$$

31.  $x = \frac{\pi}{4} \cdot (x+2) \frac{x}{x+n+1} \cdot \frac{y}{y+n+1} \cdot \frac{z}{z+n+1} \cdot \frac{u}{u+n+1}$

$$+ \frac{n(n+1)}{2!} \cdot \frac{(x+2)}{(x+n+1)(x+n+2)} \cdot \frac{x(x-1)}{(y+n+1)(y+n+2)} \cdot \frac{y(y-1)}{(z+n+1)(z+n+2)} \cdot \frac{z}{(u+n+1)(u+n+2)} \cdot \frac{u}{(u+n+1)(u+n+2)} - \dots$$

$$= \pi \cdot \frac{1}{n} \cdot \frac{1}{x+y+n} \left\{ 1 + \frac{\pi y}{4} \cdot \frac{z+u+n+1}{(z+n+1)(u+n+1)} + \frac{x(x-1) y(y-1)}{2} \cdot \frac{(z+u+n+1)(z+u+n+2)}{(z+n+1)(z+n+2)(u+n+1)(u+n+2)} + \dots \right\}$$

$$32. \frac{1}{x} + \frac{x}{4} \cdot \frac{y}{2} \cdot \frac{1}{n+1} + \frac{x(x-1)}{1!} \cdot \frac{y(y-1)}{2!(2+1)} \cdot \frac{1}{n+2} + \dots$$

$$= \frac{1 \cdot |n-1|}{|x+n|} \left\{ 1 + \frac{x}{4} \cdot \frac{y+2}{2} + \frac{x(n+1)}{1!} \cdot \frac{(y+2)(y+2+1)}{2(2+1)} + \dots \right.$$

to  $x+1$  terms  $\left. \right\}$

33. If  $x+y+z=0$ , then

$$\frac{1}{x} + \frac{x}{4} \cdot \frac{y}{2} \cdot \frac{1}{n+1} + \frac{x(x-1)}{1!} \cdot \frac{y(y-1)}{2!(2+1)} \cdot \frac{1}{n+2} + \dots$$

$$= \frac{|n-1| |x+y+n|}{|x+n| |y+n|} \left\{ 1 + \frac{x}{4} \cdot \frac{y}{2} + \frac{x(x-1)}{1!} \cdot \frac{y(y-1)}{2(2+1)} + \dots \right.$$

to  $x+y+n+1$  terms  $\left. \right\}$ .

$$34. \frac{\sqrt{\frac{x+y-1}{x+y}}}{\sqrt{\frac{x-1}{x}} \sqrt{\frac{y-1}{y}}} \sqrt{\pi} = 1 + \frac{x}{4} \cdot \frac{y}{x+y+1} + \frac{x(x+1)}{1!} \cdot \frac{y(y+1)}{(x+y+1)(x+y+3)} + \dots$$

$$+ \frac{x(x+1)(x+2)}{1!} \cdot \frac{y(y+1)(y+3)}{(x+y+1)(x+y+3)(x+y+5)} + \dots$$

Cor.  $\frac{\sqrt{\frac{x-1}{x+y-2}}}{\sqrt{\frac{x-1}{x}} \sqrt{\frac{y-1}{y}}} \sqrt{\frac{2\pi}{2^n}} = 1 + \frac{1^2 - 2^2}{4(n+1)} + \frac{(1^2 - 2^2)(3^2 - 2^2)}{4 \cdot 8(n+1)(n+3)} + \dots$

e.g.  $\frac{\sqrt{\frac{n-1}{(n-2)^2}}}{\left(\sqrt{\frac{n-2}{n}}\right)^2} \sqrt{\frac{2\pi}{2^n}} = 1 + \frac{1^2}{4(n+1)} + \frac{1^2 \cdot 3^2}{4 \cdot 8(n+1)(n+3)} + \dots$

2.  $\frac{\sqrt{\frac{n-1}{\frac{2n-3}{8} \cdot \frac{2n-5}{8}}}}{\sqrt{\frac{2n-3}{8} \cdot \frac{2n-5}{8}}} \sqrt{\frac{2\pi}{2^n}} = 1 + \frac{1 \cdot 3}{16(n+1)} + \frac{4 \cdot 3 \cdot 5 \cdot 7}{16 \cdot 32(n+1)(n+3)} + \dots$

35. If  $\phi(n) = 1 + \left(\frac{1}{2}\right)^n + \left(\frac{1 \cdot 1}{2 \cdot 4}\right)^n + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^n + \dots$  to  $n$  terms, then

i.  $\pi \phi\left(\frac{n+1}{4}\right) = 3 \log 2 + \pi \frac{1}{2^{n/2}} + \frac{3}{4n} - \frac{99}{32n^2} + \frac{999}{32n^3} - \dots$

ii.  $1 + \left(\frac{2}{3}\right)^n + \left(\frac{2 \cdot 4}{3 \cdot 9}\right)^n + \dots$  to  $n$  terms  $= \frac{\pi^2}{4} \phi\left(\frac{n+2}{2}\right) - 2\left(\frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots\right)$

iii.  $1 + \frac{16}{\pi^2} \left(\frac{1-\frac{1}{2}}{2}\right)^n \left\{ 1 + \left(\frac{2}{3}\right)^n + \left(\frac{2 \cdot 4}{3 \cdot 9}\right)^n + \dots \right.$  to  $n$  terms  $\left. \right\} = 2 \phi\left(\frac{n+2}{2}\right)$

iv.  $\phi\left(\frac{1}{2}\right) = \frac{1}{2}$  and  $\frac{\pi^2}{8} \phi\left(\frac{1}{2}\right) = \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots$

# CHAPTER XI

$$1. \frac{1}{(1+x)^a} \left[ 1 + \frac{a}{1} \cdot \frac{m}{2m} \left\{ 1 - \frac{d(-x)}{d(x)} \right\} + \frac{a(a+1)}{1 \cdot 2} \cdot \frac{m(m+1)}{2m(2m+1)} \left\{ 1 - \frac{d(-x)}{d(x)} \right\}^2 + \dots \right]$$

is always an even function of  $x$ .

$$2. 1 + \frac{a}{1} \cdot \frac{m}{2m} \cdot \frac{2x}{1+x} + \frac{a(a+1)}{1 \cdot 2} \cdot \frac{m(m+1)}{2m(2m+1)} \left( \frac{2x}{1+x} \right)^2 + \dots$$

$$= (1+x)^{-a} \left\{ 1 + \frac{a(a+1)}{2(2m+1)} x^2 + \frac{a(a+1)(a+2)(a+3)}{2 \cdot 4 \cdot (2m+1)(2m+3)} x^4 + \dots \right\}$$

$$3. 1 + \frac{a}{1} \cdot \frac{m}{2m} \cdot \frac{4x}{(1+x)^2} + \frac{a(a+1)}{1 \cdot 2} \cdot \frac{m(m+1)}{2m(2m+1)} \left\{ \frac{4x}{(1+x)^2} \right\}^2 + \dots$$

$$= (1+x)^{-2a} \left\{ 1 + \frac{a}{1} \cdot \frac{a-m+\frac{1}{2}}{m+\frac{1}{2}} x^2 + \frac{a(a+1)}{1 \cdot 2} \cdot \frac{(a-m+\frac{1}{2})(a-m+\frac{1}{2})}{(m+\frac{1}{2})(m+\frac{1}{2})} x^4 + \dots \right\}$$

$$4. 1 + \frac{a(a+1)}{2(2m+1)} \cdot \frac{4x}{(1+x)^2} + \frac{a(a+1)(a+2)(a+3)}{2 \cdot 4 \cdot (2m+1)(2m+3)} \left\{ \frac{4x}{(1+x)^2} \right\}^2 + \dots$$

$$= (1+x)^{-a} \left\{ 1 + \frac{a}{1} \cdot \frac{a-m+\frac{1}{2}}{m+\frac{1}{2}} x + \frac{a(a+1)}{1 \cdot 2} \cdot \frac{(a-m+\frac{1}{2})(a-m+\frac{1}{2})}{(m+\frac{1}{2})(m+\frac{1}{2})} x^2 + \dots \right\}$$

$$5. 1 + \frac{1}{2} \cdot \frac{a}{1} \cdot \frac{4x}{(1+x)^2} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{a(a+1)}{1 \cdot 2} \cdot \left\{ \frac{4x}{(1+x)^2} \right\}^2 + \dots$$

$$= (1+x)^{-2a} \left\{ 1 + \left( \frac{a}{1} \right)^2 x^2 + \left[ \frac{a(a+1)}{1 \cdot 2} \right]^2 x^4 + \dots \right\}$$

$$6. 1 + \frac{a(a+1)}{2^2} \cdot \frac{4x}{(1+x)^2} + \frac{a(a+1)(a+2)(a+3)}{2^2 \cdot 4^2} \left\{ \frac{4x}{(1+x)^2} \right\}^2 + \dots$$

$$= (1+x)^{-a} \left\{ 1 + \left( \frac{a}{1} \right)^2 x + \left[ \frac{a(a+1)}{1 \cdot 2} \right]^2 x^2 + \dots \right\}$$

$$7. 1 + \frac{2x}{1} \cdot \frac{m}{2m} + \frac{(2x)^2}{1 \cdot 2} \cdot \frac{m(m+1)}{2m(2m+1)} + \frac{(2x)^3}{1 \cdot 3} \cdot \frac{m(m+1)(m+2)}{2m(2m+1)(2m+2)} + \dots$$

$$= e^x \left\{ 1 + \frac{x^2}{2} \cdot \frac{1}{2m+1} + \frac{x^4}{3 \cdot 4} \cdot \frac{1}{(2m+1)(2m+3)} + \dots \right\}$$

$$\text{Cor. } 1 + \frac{1}{2} \cdot \frac{a}{1} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{x^2}{1} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{x^3}{1} + \dots =$$

$$e^{\frac{x}{2}} \left\{ 1 + \frac{x^2}{4!} + \frac{x^4}{2! \cdot 8!} + \frac{x^6}{4! \cdot 8! \cdot 12!} + \dots \right\}$$

$$9. \phi(x) + \frac{2\phi'(0)}{1!} \cdot \frac{x}{2m} + \frac{2^2 \phi''(0)}{2!} \cdot \frac{x^2}{2m(2m+1)} + \dots$$

$$= \phi(1) + \frac{\phi''(0)}{2 \cdot 1!} \cdot \frac{1}{2m+1} + \frac{\phi^{IV}(0)}{2^2 \cdot 2!} \cdot \frac{1}{(2m+1)(2m+3)} + \dots$$

$$9. 1 + \frac{x^2}{2} \cdot \frac{1}{2n+1} + \frac{x^4}{2 \cdot 4} \cdot \frac{1}{(2n+1)(2n+3)} + \dots$$

$$= \frac{e^x \sqrt{\pi}}{x^n \sqrt{\pi}} \left[ e^x \left\{ 1 - \frac{n(n-1)}{2} \cdot \frac{1}{x} + \frac{(n+1)n(n-1)(n-2)}{2 \cdot 4} \cdot \frac{1}{x^2} - \dots \right\} \right.$$

$$\left. + e^{-x} \cos \pi n \left\{ 1 + \frac{n(n-1)}{2} \cdot \frac{1}{x} + \frac{(n+1)n(n-1)(n-2)}{2 \cdot 4} \cdot \frac{1}{x^2} + \dots \right\} \right]$$

$$\text{Cos. } 1 + \frac{x^2}{2!} + \frac{x^4}{2! \cdot 4!} + \frac{x^6}{2! \cdot 4! \cdot 6!} + \dots$$

$$= \frac{e^x}{\sqrt{2\pi x}} \left( 1 + \frac{1^2}{8x} + \frac{1^2 \cdot 3^2}{8 \cdot 16 x^2} + \frac{1^2 \cdot 3^2 \cdot 5^2}{8 \cdot 16 \cdot 24 x^3} + \dots \right)$$

$$10. 1 - \frac{x^2}{2} \cdot \frac{1}{2n+1} + \frac{x^4}{2 \cdot 4} \cdot \frac{1}{(2n+1)(2n+3)} - \dots$$

$$= \frac{e^x \sqrt{\pi}}{x^n \sqrt{\pi}} \left[ \cos\left(\frac{\pi x}{2} - x\right) \left\{ 1 - \frac{(n+1)n(n-1)(n-2)}{2 \cdot 4} \cdot \frac{1}{x^2} + \dots \right\} \right.$$

$$\left. + \sin\left(\frac{\pi x}{2} - x\right) \left\{ \frac{n(n-1)}{2x} - \frac{(n+2)(n+1)n(n-1)(n-2)(n-3)}{2 \cdot 4 \cdot 6 x^3} + \dots \right\} \right]$$

$$\text{Cos. } \int \left( 1 - \frac{x^2}{2} \cdot \frac{1}{2n+1} + \frac{x^4}{2 \cdot 4} \cdot \frac{1}{(2n+1)(2n+3)} - \dots \right) - \dots = 0$$

$$\text{then } x = \frac{\pi(\mu + \pi)}{2} - \frac{\pi(n-1)}{\pi(\mu + \pi)} - \frac{n(n-1)(7\pi, \pi-1-6)}{3\pi^3(\mu + \pi)^2} - \dots$$

where  $\mu$  is any odd integer.

$$11. \int_0^{\pi} \frac{\sin x}{x} dx = \frac{\pi}{2} - n \cos(x-\theta), \text{ then}$$

$$\int_0^{\pi} \frac{1 - \cos x}{x} dx = c + \log x - n \sin(x-\theta)$$

$$\text{where } n^2 = \frac{11}{2x} - \frac{13}{2x^3} + \frac{15}{3x^5} - \frac{17}{4x^7} + \dots$$

$$\cos \theta = \frac{1}{x} - \frac{12}{x^3} + \frac{16}{x^5} - \frac{16}{x^7} + \dots \text{ and}$$

$$\sin \theta = \frac{11}{x^2} - \frac{12}{x^4} + \frac{15}{x^6} - \frac{12}{x^8} + \dots$$

Ex. 1.  $\int_0^{\frac{\pi}{2}} \cos(\pi \sin^2 \theta) d\theta = 0.$

2.  $\int_0^{\frac{\pi}{2}} \cos(2\pi \sin^2 \theta) d\theta = - \int_0^{\frac{\pi}{2}} \cos(\pi \sin^2 \theta) d\theta.$

3.  $\int_0^{\frac{\pi}{2}} \cos\left(\frac{2\pi}{3} \sin^2 \theta\right) d\theta = \frac{1}{2} \int_0^{\frac{\pi}{2}} \cos\left(\frac{\pi}{3} \sin^2 \theta\right) d\theta.$

12. If  $x + y + z = \frac{1}{2}$ , then

$$1 + \frac{x}{4} \cdot \frac{y}{2} p + \frac{x(x-1)}{12} \cdot \frac{y(y-1)}{2(2+1)} p^2 + \frac{x(x-1)(x-2)}{12} \cdot \frac{y(y-1)(y-2)}{2(2+1)(2+2)} p^3 + \dots$$

$$= 1 + \frac{2x}{12} \cdot \frac{2y}{2} \cdot \frac{1-\sqrt{1-p}}{2} + \frac{2x(2x-1)}{12} \cdot \frac{2y(2y-1)}{2(2+1)} \left(\frac{1-\sqrt{1-p}}{2}\right)^2 + \dots$$

$$\text{ex. } 1 + \frac{1+n}{4^2} x + \frac{(1+n)(5^2+n)}{4^2 \cdot 8^2} x^2 + \frac{(1+n)(5^2+n)(9^2+n)}{4^2 \cdot 8^2 \cdot 11^2} x^3 + \dots$$

$$+ \dots = 1 + \frac{1+n}{2^2} \cdot \frac{1-\sqrt{1-x}}{2} + \frac{(1+n)(3^2+n)}{2^2 \cdot 4^2} \left(\frac{1-\sqrt{1-x}}{2}\right)^2 + \dots$$

e.g.  $1 + \frac{1}{2}(1+\frac{1}{2}) \frac{1-\sqrt{1-x}}{2} + \frac{1}{3}(1+\frac{1}{2})(1+\frac{1}{2} \cdot 2) \left(\frac{1-\sqrt{1-x}}{2}\right)^2 + \dots$

$$= 1 + \frac{x}{2^2} + \frac{2}{1} \cdot \left(1 + \frac{1}{2}\right) \frac{x^2}{4^2} + \frac{2 \cdot 4}{1 \cdot 3} \left(1 + \frac{1}{2}\right) \left(1 + \frac{1}{2} \cdot 2\right) \frac{x^3}{6^2} + \dots$$

ex.  $\left(\frac{1+\sqrt{1-x}}{2}\right)^4 \left\{ 1 + \frac{(1+\gamma)(\beta+\gamma)}{4 \cdot (1+1)} x + \frac{(1+\gamma)(1+\gamma+2)(\beta+\gamma)(\beta+\gamma+2)}{4 \cdot 8 \cdot (1+1)(1+2)} x^2 + \dots \right\}$

$$= 1 + \frac{\alpha}{12} \cdot \frac{\beta}{1+1} \cdot \frac{1-\sqrt{1-x}}{2} + \frac{\alpha(\alpha+1)}{12} \cdot \frac{\beta(\beta+1)}{(1+1)(1+2)} \left(\frac{1-\sqrt{1-x}}{2}\right)^2 + \dots$$

13. If  $\alpha + \beta + \gamma = 0$ , then

$$\left\{ 1 + \frac{\alpha}{12} \cdot \frac{\beta}{1+\frac{1}{2}} x + \frac{\alpha(\alpha-1)}{12} \cdot \frac{\beta(\beta-1)}{(1+\frac{1}{2})(1+\frac{1}{2})} x^2 + \dots \right\}^2$$

$$= 1 + \frac{2\alpha}{12} \cdot \frac{2\beta}{1+\frac{1}{2}} \cdot \frac{\gamma}{2\gamma} x + \frac{2\alpha(2\alpha-1)}{12} \cdot \frac{2\beta(2\beta-1)}{(1+\frac{1}{2})(1+\frac{1}{2})} \cdot \frac{\gamma(\gamma+1)}{2(1+\frac{1}{2})} x^2 + \dots$$

$$\text{Cor. 1. } \left\{ 1 + \frac{(1+n)}{4} x + \frac{(1+n)(3+n)}{4^2 \cdot 8} x^2 + \dots \right\}^2 =$$

$$1 + \frac{1}{2} \cdot \frac{(1+n)}{2} x + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{(1+n)(3+n)}{2^2 \cdot 4} x^2 + \dots$$

$$\text{Cor. 2. } \left\{ 1 + \frac{x}{2} + \frac{x^2}{2 \cdot 4} + \frac{x^3}{2^2 \cdot 4^2 \cdot 8} + \dots \right\}^2$$

$$= 1 + \frac{1}{2} \cdot \frac{x}{(1)} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{x^2}{(2)^2} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{x^3}{(12)^2} + \dots$$

14. If  $\alpha + \beta + 1 = \gamma + \delta,$

$$\left\{ 1 + \frac{\alpha}{u} \cdot \frac{\beta}{\gamma} \cdot \frac{1 - \sqrt{1-x}}{2} + \frac{\alpha(\alpha+1)}{u^2} \cdot \frac{\beta(\beta+1)}{\gamma(\gamma+1)} \cdot \left( \frac{1 - \sqrt{1-x}}{2} \right)^2 + \dots \right\}$$

$$\times \left\{ 1 + \frac{\alpha}{u} \cdot \frac{\beta}{\delta} \cdot \frac{1 - \sqrt{1-x}}{2} + \frac{\alpha(\alpha+1)}{u^2} \cdot \frac{\beta(\beta+1)}{\delta(\delta+1)} \cdot \left( \frac{1 - \sqrt{1-x}}{2} \right)^2 + \dots \right\}$$

$$= 1 + \frac{\alpha}{\gamma} \cdot \frac{\beta}{\delta} \cdot \frac{(\alpha+\beta)(\gamma+\delta)}{2 \cdot (2\alpha+2\beta)} x + \frac{\alpha(\alpha+1)}{\gamma(\gamma+1)} \cdot \frac{\beta(\beta+1)}{\delta(\delta+1)} \cdot \frac{(\alpha+\beta)(\alpha+\beta+2)}{2 \cdot 4} x^2$$

$$+ \frac{(\alpha+\delta)(\gamma+\delta+2)}{(2\alpha+2\beta)(2\alpha+2\beta+2)} x^2 + \dots$$

$$15. \left\{ 1 + \frac{x}{u} \cdot \frac{1}{\gamma} + \frac{x^2}{u^2} \cdot \frac{1}{\gamma(\gamma+1)} + \frac{x^3}{u^3} \cdot \frac{1}{\gamma(\gamma+1)(\gamma+2)} + \dots \right\}$$

$$\times \left\{ 1 + \frac{x}{u} \cdot \frac{1}{\delta} + \frac{x^2}{u^2} \cdot \frac{1}{\delta(\delta+1)} + \frac{x^3}{u^3} \cdot \frac{1}{\delta(\delta+1)(\delta+2)} + \dots \right\}$$

$$= 1 + \frac{x}{u} \cdot \frac{\gamma+\delta}{\gamma\delta} + \frac{x^2}{u^2} \cdot \frac{(\gamma+\delta+1)(\gamma+\delta+2)}{\gamma(\gamma+1)\delta(\delta+1)} + \frac{x^3}{u^3} \cdot \frac{(\gamma+\delta+3)(\gamma+\delta+2)}{\gamma(\gamma+1)(\gamma+2)\delta(\delta+2)}$$

$$+ \frac{\gamma+\delta+3}{(\delta+1)(\delta+2)} + \frac{x^4}{u^4} \cdot \frac{(\gamma+\delta+3)(\gamma+\delta+4)(\gamma+\delta+1)(\gamma+\delta+2)}{\gamma(\gamma+1)(\gamma+2)(\gamma+3)\delta(\delta+1)(\delta+2)(\delta+3)} + \dots$$

$$16. \left\{ 1 + \frac{x}{u} \cdot \frac{1}{m+1} \cdot \frac{1}{n+1} + \frac{x^2}{u^2} \cdot \frac{1}{(m+1)(m+2)} \cdot \frac{1}{(n+1)(n+2)} + \dots \right\} \times$$

$$\left\{ 1 - \frac{x}{u} \cdot \frac{1}{m+1} \cdot \frac{1}{n+1} + \frac{x^2}{u^2} \cdot \frac{1}{(m+1)(m+2)} \cdot \frac{1}{(n+1)(n+2)} - \dots \right\}$$

$$= 1 - \frac{x^2}{u^2} \cdot \frac{m+n+3}{(m+1)(m+2)} \cdot \frac{1}{(m+1)(m+2)} \cdot \frac{1}{(n+1)(n+2)}$$

$$+ \frac{x^4}{u^4} \cdot \frac{(m+n+5)(m+n+6)}{(m+1)(m+2)(m+3)(m+4)(n+1)(n+2)(n+3)(n+4)}$$

$$x \frac{1}{(n+2)(n+4)} - \frac{x^6}{13} \frac{(m+n+1)(m+n+2)(m+n+3)}{(m+1)(m+2)(m+3)(m+4)(m+5)(m+6)} \times \frac{(n+1)(n+2)(n+3)(n+4)(n+5)(n+6)}{(n+1)(n+2)(n+3)(n+4)(n+5)(n+6)} + \dots$$

$$17. \left\{ 1 + \frac{x}{2} \cdot \frac{1}{m+n+1} \cdot \frac{1}{n+1} + \frac{x^2}{2} \cdot \frac{(m+n+1)(m+n+2)}{(n+1)(n+2)} + \dots \right\} \\ + \dots \left\{ 1 + \frac{x}{2} \cdot \frac{1}{m+1} \cdot \frac{1}{n+1} + \frac{x^2}{2} \cdot \frac{(m+1)(m+2)}{(m+1)(m+2)} + \dots \right\} \\ = 1 + \frac{x}{2} \cdot \frac{2m+n+3}{m+n+1} \cdot \frac{1}{m+1} \cdot \frac{1}{n+1} + \frac{x^2}{2} \cdot \frac{(2m+n+4)(2m+n+6)}{(m+n+1)(m+n+2)} \times \\ \frac{1}{(m+1)(m+2)} \cdot \frac{1}{n^2-2} + \frac{x^3}{2} \cdot \frac{(2m+n+5)(2m+n+7)(2m+n+9)}{(m+n+1)(m+n+2)(m+n+3)} \times \\ \frac{1}{(m+1)(m+2)(m+3)} \cdot \frac{1}{(n^2-1)(n^2-3)} + \dots$$

$$18. \left\{ 1 + \frac{\beta}{\gamma} \cdot \frac{x}{2} + \frac{\beta(\beta-1)}{\gamma(\gamma+1)} \cdot \frac{x^2}{2} + \frac{\beta(\beta-1)(\beta-2)}{\gamma(\gamma+1)(\gamma+2)} \cdot \frac{x^3}{2} + \dots \right\} \times \\ \left\{ 1 - \frac{\beta}{\gamma} \cdot \frac{x}{2} + \frac{\beta(\beta-1)}{\gamma(\gamma+1)} \cdot \frac{x^2}{2} - \frac{\beta(\beta-1)(\beta-2)}{\gamma(\gamma+1)(\gamma+2)} \cdot \frac{x^3}{2} + \dots \right\} \\ = 1 - \frac{\beta}{\gamma} \cdot \frac{\beta+\gamma}{\gamma(\gamma+1)} \cdot \frac{x^2}{2} + \frac{\beta(\beta-1)}{\gamma(\gamma+1)} \cdot \frac{\beta+\gamma}{\gamma(\gamma+1)(\gamma+2)(\gamma+3)} \cdot \frac{x^4}{2} - \dots$$

$$19. \left\{ 1 + \frac{x}{2} d\beta + \frac{x^2}{2} d(d-1)\beta(\beta-1) + \dots \right\} \times \\ \left\{ 1 - \frac{x}{2} d\beta + \frac{x^2}{2} d(d-1)\beta(\beta-1) - \dots \right\} \\ = 1 - \frac{x^2}{2} d\beta(d+\beta-1) + \frac{x^4}{2} d(d-1)\beta(\beta-1)(d+\beta-1)(d+\beta-3) \\ - \frac{x^6}{2} d(d-1)(d-2)\beta(\beta-1)(\beta-2)(d+\beta-3)(d+\beta-4)(d+\beta-5) + \dots$$

$$20. \left\{ 1 + \frac{x}{2} \cdot \frac{m}{n+1} + \frac{x^2}{2} \cdot \frac{m(m-1)}{(n+1)(n+2)} + \dots \right\} \times \\ \left\{ 1 + \frac{x}{2} \cdot \frac{m+n}{n-1} + \frac{x^2}{2} \cdot \frac{(m+n)(m+n-1)}{(n-1)(n-2)} + \dots \right\} \\ = 1 + \frac{x}{2} \cdot (2m+n+1) \frac{1}{n^2-1} + \frac{x^2}{2} \cdot (2m+n)(2m+n+2) \frac{1}{n^2-2} + \\ \frac{x^3}{2} \cdot (2m+n-1)(2m+n+1)(2m+n+3) \frac{1}{(n^2-1)(n^2-3)} + \dots$$

$$21. \text{e.g. } (1 + \frac{x^3}{2} + \frac{x^6}{2} + \frac{x^9}{2} + \dots) (1 - \frac{x^3}{2} + \frac{x^6}{2} - \frac{x^9}{2} + \dots) =$$



130.  $\frac{1}{3} + \frac{x}{3} \left\{ 1 - \frac{(3x^2)^2}{16} + \frac{(3x^2)^6}{112} - \frac{(3x^2)^9}{118} + \dots \right\}$

2.  $\left\{ 1 + \frac{x}{(4)}^2 + \frac{x^2}{(12)^2} + \frac{x^3}{(12)^2} + \dots \right\} \left\{ 1 - \frac{x}{(12)^2} + \frac{x^2}{(12)^2} - \frac{x^3}{(12)^2} + \dots \right\}$   
 $= 1 - \frac{x^3 13}{(12)^3} + \frac{x^4 16}{(12 \cdot 14)^3} - \frac{x^6 12}{(12 \cdot 16)^3} + \dots$

3.  $(x + \frac{x^2}{16} + \frac{x^2}{12} + \frac{x^{10}}{110} + \dots)(x - \frac{x^4}{15} + \frac{x^7}{12} - \frac{x^{10}}{110} + \dots)$   
 $= \frac{x}{3} \left\{ \frac{3x^2}{12} - \frac{(3x^2)^6}{118} + \frac{(3x^2)^7}{114} - \dots \right\}$

4.  $\cos x \cosh x = 1 - \frac{(2x^2)^2}{15} + \frac{(2x^2)^6}{18} - \frac{(2x^2)^6}{112} + \dots$

5.  $\sin x \sinh x = \frac{2x^2}{12} - \frac{(2x^2)^3}{16} + \frac{(2x^2)^5}{110} - \dots$

6.  $\left\{ 1 + \frac{x}{(12)} + \frac{x^2}{(12)^2} + \frac{x^3}{(12)^2} + \dots \right\} \left\{ 1 - \frac{x}{(12)} + \frac{x^2}{(12)} - \frac{x^3}{(12)^2} + \dots \right\}$   
 $= 1 - \frac{x^2}{(12)^2 12} + \frac{x^4}{(12)^2 16} - \frac{x^6}{(12)^2 16} + \dots$

7.  $\left\{ 1 + \frac{1}{2} \cdot \frac{x}{12} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{x^2}{12^2} + \dots \right\} \left\{ 1 - \frac{1}{2} \cdot \frac{x}{12} + \frac{1 \cdot 1}{2 \cdot 4} \cdot \frac{x^2}{12^2} - \dots \right\}$   
 $= 1 + \frac{1}{2} \cdot \frac{x^2}{12} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{x^2}{2 \cdot 4} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{x^6}{2 \cdot 4 \cdot 6} + \dots$

8.  $(1 + \frac{x}{1 \cdot 3} + \frac{x^2}{1 \cdot 3 \cdot 5} + \frac{x^3}{1 \cdot 3 \cdot 5 \cdot 7} + \dots)(1 - \frac{x}{1 \cdot 3} + \frac{x^2}{1 \cdot 3 \cdot 5} - \frac{x^3}{1 \cdot 3 \cdot 5 \cdot 7} + \dots)$   
 $= 1 + \frac{x^2}{1 \cdot 3 \cdot 5} \cdot \frac{1}{3} + \frac{x^4}{1 \cdot 2 \cdot 5 \cdot 7 \cdot 9} \cdot \frac{1}{5} + \dots$

9.  $\left\{ \frac{1}{n} + \frac{x}{n(n+1)} + \frac{x^2}{n(n+1)(n+2)} + \dots \right\} \left\{ \frac{1}{n} - \frac{x}{n(n+1)} + \dots \right\}$   
 $= \frac{1}{n} \cdot \frac{1}{n} + \frac{x^2}{n(n+1)(n+2)} \cdot \frac{1}{n+1} + \frac{x^4}{n(n+1)(n+2)(n+3)(n+4)} \cdot \frac{1}{n+2} + \dots$

10.  $\left\{ 1 + xn + x^2 n(n-1) + x^3 n(n-1)(n-2) + \dots \right\} \left\{ 1 - xn + x^2 n(n-1) - \dots \right\}$   
 $= \frac{1}{2} \cdot n + \frac{x^2}{n!} \cdot n(n-1)(n-2) + \frac{x^4}{n!} \cdot n(n-1)(n-2)(n-3)(n-4) + \dots$

21.  $1 + \frac{1+x}{4} \cdot \frac{m \cdot n}{m+n+1} + \frac{(1+x)^2}{12} \cdot \frac{m(m+1) \cdot n(n+1)}{(m+n+1)(m+n+3)} + \dots$

$$= \sqrt{\frac{m+1}{2}} \left\{ 1 + \frac{x^2}{2} m(m+1) + \frac{x^4}{24} m(m+1)(m+2)(m+3) + \dots \right\} +$$

$$2 \sqrt{\frac{m+1}{2}} \left\{ \frac{x}{2} + \frac{x^3}{12} (m+1)(m+2) + \frac{x^5}{120} (m+1)(m+2)(m+3)(m+4) + \dots \right\}$$

29.  $e^{-mx} = 1 + \frac{1}{2} \cdot \frac{m^2}{2!} (1 - e^{-2x}) + \frac{1}{24} \cdot \frac{m^3}{3!} (1 - e^{-2x})^2 + \dots$

$$= 1 + \frac{A_1}{2} \cdot \left(\frac{x}{2}\right)^2 + \frac{A_2}{2^2} \cdot \left(\frac{x}{2}\right)^4 + \frac{A_3}{2^3} \cdot \left(\frac{x}{2}\right)^6 + \dots$$

where  $A_n = p^n - \frac{n(n-1)}{2} p^{n-1} + \frac{n(n-1)(n-2)(3n-1)}{24} p^{n-2}$   
 $\dots + (-1)^{n-1} 2p \cdot \frac{n-1}{1 \cdot 3 \cdot 5 \dots (2n-1)} B_{2n-2}$  and  $p = \frac{m(m-1)}{2}$ .

Cor. If  $A_n = K p^n$ , then  $K_1 = \frac{1}{1 \cdot 3 \cdot 5 \dots (2n-1)}$ ;  $K_3 = 3 \cdot 2^{2(n-1)} K_1$ ; &c.

23. If  $\phi(x)$  can be expressed in  $n$  different ways, the apparent values in the  $n$ th way being  $C_n + V_n$  and if  $c_1, c_2, c_3, \dots, c_n$  appear to be similar and  $v_1, v_2, v_3, \dots, v_n$  are known to be dissimilar, then  $c_1, c_2, c_3, \dots, c_n$  must be identically equal (say equal to  $c$ ) and the real value of  $\phi(x) = c + v_1 + v_2 + v_3 + \dots + v_n$ .

24. If  $\phi(x) = \frac{1}{x^n} \left\{ P_0 x^n + n P_1 x^{n-1} + \frac{n(n-1)}{2} P_2 x^{n-2} + \dots \right\}$   
 and  $Q_n = \phi(x) + \frac{n+1}{x} \phi(x+1) + \frac{(n+1)(n+2)}{2} \phi(x+2) + \dots$ , then

$$\phi(0) + (1-x)\phi(1) + (1-x)^2\phi(2) + (1-x)^3\phi(3) + \dots =$$

$$Q_0 - Q_1 x + Q_2 x^2 - Q_3 x^3 + \dots + \frac{1}{(\log \frac{1}{1-x})^{n+1}} \left\{ P_0 + P_1 \log \frac{1}{1-x} + P_2 (\log \frac{1}{1-x})^2 + \dots \right\}.$$

Cor. If  $Q_n' = \frac{1}{(m-n)!} \phi(m) + \frac{1}{(m-n+1)!} \phi(m+1) + \frac{1}{(m-n+2)!} \phi(m+2) + \dots$   
 then  $\phi(m)(1-x)^m + \phi(m+1)(1-x)^{m+1} + \phi(m+2)(1-x)^{m+2} + \dots$   
 $= Q_0' - Q_1' x + Q_2' x^2 - Q_3' x^3 + Q_4' x^4 - \dots +$

$$132. \frac{1}{(\log \frac{1}{1-x})^{n+1}} \{ P_0 + P_1 \log \frac{1}{1-x} + P_2 (\log \frac{1}{1-x})^2 + \dots \}$$

Cor. 2. If  $\alpha + \beta + \gamma + 1 = \delta + \epsilon$ , then when  $x$  vanishes

$$\frac{\alpha}{10} \frac{\beta}{15} \frac{\gamma}{16} + (1-x) \frac{\alpha+1}{11} \frac{\beta+1}{16} \frac{\gamma+1}{17} + (1-x)^2 \frac{\alpha+2}{12} \frac{\beta+2}{17} \frac{\gamma+2}{18} + \dots$$

$$+ \log x + \dots \frac{1}{\alpha} + \dots \frac{1}{\beta} =$$

$$1. \frac{(\gamma - \delta)(\gamma - \epsilon)}{(\alpha+1)(\beta+1)} + \frac{1}{2} \frac{(\gamma - \delta)(\gamma - \delta - 1)(\gamma - \epsilon)(\gamma - \epsilon - 1)}{(\alpha+1)(\alpha+2)(\beta+1)(\beta+2)} + \dots$$

$$25. \alpha \beta \left\{ \frac{\alpha+n}{2} \frac{\beta+n}{2} \frac{1}{\alpha+\beta+n+1} + \frac{1-x}{11} \frac{\alpha+n+1}{2} \frac{\beta+n+1}{2} + \dots \right\}$$

$$= \left\{ \frac{\alpha+n}{2} \frac{\beta+n}{2} \frac{1}{n-1} - \frac{x}{11} \frac{\alpha+n+1}{2} \frac{\beta+n+1}{2} \frac{1}{n-2} + \dots \right\}$$

$$+ \frac{1}{2^n} \left\{ \alpha \beta \frac{1}{n-1} - \frac{x}{11} \frac{\alpha+1}{2} \frac{\beta+1}{2} \frac{1}{n-2} + \dots \right\}$$

N.B. Though the above theorem is true for all values of  $n$  yet if  $n$  is an integer it assumes the form  $\infty - \infty$ ; so we must write  $n+h$  for  $n$  and then after simplification  $h$  should be made to vanish.

Cor. 1. If  $n$  is a positive integer,

$$\alpha \beta \left\{ \frac{\alpha+n}{2} \frac{\beta+n}{2} \frac{1}{\alpha+\beta+n+1} + \frac{1-x}{11} \frac{\alpha+n+1}{2} \frac{\beta+n+1}{2} + \dots \right\}$$

$$+ (-1)^n \log x \left\{ \frac{\alpha+n}{2} \frac{\beta+n}{2} \frac{1}{n} + \frac{x}{11} \frac{\alpha+n+1}{2} \frac{\beta+n+1}{2} \frac{1}{n+1} + \dots \right\}$$

$$+ (-1)^n \left\{ \frac{\alpha+n}{2} \frac{\beta+n}{2} \frac{1}{n} \left( \frac{1}{\alpha+n} + \frac{1}{\beta+n} - \frac{1}{n} - \delta \right) + \right.$$

$$\frac{x}{11} \frac{\alpha+n+1}{2} \frac{\beta+n+1}{2} \frac{1}{n+1} \left( \frac{1}{\alpha+n+1} + \frac{1}{\beta+n+1} - \frac{1}{n+1} - \delta \right) +$$

$$\left. \frac{x^2}{12} \frac{\alpha+n+2}{2} \frac{\beta+n+2}{2} \frac{1}{n+2} \left( \frac{1}{\alpha+n+2} + \frac{1}{\beta+n+2} - \frac{1}{n+2} - \delta \right) + \dots \right\}$$

$$= \frac{1}{2^n} \left\{ \alpha \beta \frac{1}{n-1} - \frac{x}{11} \frac{\alpha+1}{2} \frac{\beta+1}{2} \frac{1}{n-2} + \dots \text{to } n \text{ terms} \right\}$$

Cor. 2. If  $n$  is a negative integer,

$$\begin{aligned} & \log x \left\{ \frac{|a+n| |b+n|}{|a+b+n+1|} + 1-x \frac{|a+n+1| |b+n+1|}{|a+b+n+2|} + \&c \right\} \\ & + (x)^{-n} \log x \left\{ \frac{|a| |b|}{|a-b|} + x \frac{|a+1| |b+1|}{|a-b+1|} + x^2 \frac{|a+2| |b+2|}{|a-b+2|} + \&c \right\} \\ & + (x)^{-n} \left\{ \frac{|a| |b|}{|a-b|} \left( \leq \frac{1}{a} + \leq \frac{1}{b} - \leq \frac{1}{a-b} - 0 \right) + \right. \\ & \quad x \frac{|a+1| |b+1|}{|a-b+1|} \left( \leq \frac{1}{a+1} + \leq \frac{1}{b+1} - \leq \frac{1}{a-b+1} - \leq \frac{1}{1} \right) + \\ & \quad \left. x^2 \frac{|a+2| |b+2|}{|a-b+2|} \left( \leq \frac{1}{a+2} + \leq \frac{1}{b+2} - \leq \frac{1}{a-b+2} - \leq \frac{1}{2} \right) + \&c \right\} \\ & = |a+n| |b+n| |a-b-n-1| - x \frac{|a+n+1| |b+n+1| |a-b-n-2|}{|a-b-n-2|} + \&c \quad b-n \end{aligned}$$

terms. N.B. We may put  $n=0$  either in Cor. 1 or Cor. 2.

$$\begin{aligned} 16. & \log x \left\{ \frac{|a| |b|}{|a+b+1|} + (1-x) \frac{|a+1| |b+1|}{|a+b+2|} + \frac{(1-x)^2}{2} \frac{|a+2| |b+2|}{|a+b+3|} + \&c \right\} \\ & + \log x \left\{ |a| |b| + x \frac{|a+1| |b+1|}{|a-b|} + x^2 \frac{|a+2| |b+2|}{|a-b|} + \&c \right\} \\ & + |a| |b| \left( \leq \frac{1}{a} + \leq \frac{1}{b} \right) + x \frac{|a+1| |b+1|}{|a-b|} \left( \leq \frac{1}{a+1} + \leq \frac{1}{b+1} - 2 \times \frac{1}{1} \right) \\ & + x^2 \frac{|a+2| |b+2|}{|a-b|} \left( \leq \frac{1}{a+2} + \leq \frac{1}{b+2} - 2 \leq \frac{1}{2} \right) + \&c = 0. \end{aligned}$$

$$\text{Cor. } \pi \left\{ 1 + \left(\frac{1}{2}\right)^2 (1-x) + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)^2 (1-x)^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 2 \cdot 2}\right)^2 (1-x)^3 + \&c \right\}$$

$$\begin{aligned} & = \log x \left\{ 1 + \left(\frac{1}{2}\right)^2 x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)^2 x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 2 \cdot 2}\right)^2 x^3 + \&c \right\} \\ & - 4 \left\{ \left(\frac{1}{2}\right)^2 \frac{1}{1 \cdot 2} x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)^2 \left( \frac{1}{1 \cdot 2} + \frac{1}{3 \cdot 2} \right) x^2 + \&c \right\}. \end{aligned}$$

$$\begin{aligned} \text{ex. } \int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{\tan \frac{\phi}{2}}{\sqrt{1-x \cos^2 \theta \cos^2 \phi}} d\theta d\phi &= \frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-(1-x) \sin^2 \phi}} \\ &+ \frac{1}{2} \log x \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}. \end{aligned}$$

$$27. \left(\frac{1}{2}\right)^2 x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)^2 (1+x) x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 2 \cdot 2}\right)^2 (1+x+\frac{1}{2}) x^3 + \&c =$$

$$- \frac{1}{4} \left\{ 1 + \binom{1}{2} x + \binom{1 \cdot 2}{2 \cdot 2} x^2 + \binom{1 \cdot 2 \cdot 1}{2 \cdot 2 \cdot 2} x^3 + \dots \right\} \log(1-x)$$

$$\text{ex. 1. } e^{-\pi} \frac{1 + \binom{1}{2} (1-x) + \dots}{1 + \binom{1}{2} x + \dots} = \frac{1}{16} (x + \frac{x^2}{2} + \frac{2!}{64} x^3 + \dots)$$

$$2. e^{-\frac{2\pi}{\sqrt{3}}} \frac{1 + \frac{1 \cdot 2}{3} (1-x) + \dots}{1 + \frac{1 \cdot 2}{3} x + \dots} = \frac{1}{27} (x + \frac{5}{9} x^2 + \dots)$$

$$3. e^{-\pi\sqrt{3}} \frac{1 + \frac{1 \cdot 2}{6} (1-x) + \dots}{1 + \frac{1 \cdot 2}{6} x + \dots} = \frac{1}{64} (x + \frac{5}{8} x^2 + \dots)$$

$$4. e^{-2\pi} \frac{1 + \frac{1 \cdot 2}{6} (1-x) + \dots}{1 + \frac{1 \cdot 2}{6} x + \dots} = \frac{1}{432} (x + \frac{13}{18} x^2 + \dots)$$

$$28. \phi(0) \frac{\underline{a} \underline{b}}{\underline{a} \underline{b}} \underline{m-1} - \frac{\phi(1)}{\underline{1}} \frac{\underline{a+1} \underline{b+1}}{\underline{a+1} \underline{b+1}} \underline{m-2} + \dots$$

$$+ \phi(2) \frac{\underline{a+2} \underline{b+2}}{\underline{a+2} \underline{b+2}} \underline{m-1} - \frac{\phi(n+1)}{\underline{1}} \frac{\underline{a+n+1} \underline{b+n+1}}{\underline{a+n+1} \underline{b+n+1}} \underline{m-2} + \dots$$

$$= \underline{a+n} \underline{b+n} \left\{ \phi(0) \frac{\underline{a} \underline{b}}{\underline{a+b+n+1}} + \frac{\phi(0) - \phi(1)}{\underline{1}} \frac{\underline{a+1} \underline{b+1}}{\underline{a+b+n+1}} \right.$$

$$\left. + \frac{\phi(0) - 2\phi(1) + \phi(2)}{\underline{1}} \frac{\underline{a+2} \underline{b+2}}{\underline{a+b+n+1}} + \dots \right\}$$

$$\text{Cor: } \underline{a} \underline{b} \left\{ \phi(0) \frac{\underline{a} \underline{b}}{\underline{a+b+1}} + \frac{\phi(0) - \phi(1)}{\underline{1}} \frac{\underline{a+1} \underline{b+1}}{\underline{a+b+1}} + \dots \right\}$$

$$+ \phi'(0) \underline{a} \underline{b} + \phi'(1) \frac{\underline{a+1} \underline{b+1}}{\underline{1} \underline{1}} + \phi'(2) \frac{\underline{a+2} \underline{b+2}}{\underline{1} \underline{1}} + \dots$$

$$+ \phi(0) \underline{a} \underline{b} \left( \varepsilon \frac{1}{\underline{a}} + \varepsilon \frac{1}{\underline{b}} \right) + \phi(1) \frac{\underline{a+1} \underline{b+1}}{\underline{1} \underline{1}} \left( \varepsilon \frac{1}{\underline{a+1}} + \varepsilon \frac{1}{\underline{b+1}} - \varepsilon \right)$$

$$+ \phi(2) \frac{\underline{a+2} \underline{b+2}}{\underline{1} \underline{1}} \left( \varepsilon \frac{1}{\underline{a+2}} + \varepsilon \frac{1}{\underline{b+2}} - 2\varepsilon \frac{1}{\underline{2}} \right) + \dots = 0.$$

$$29. \int F(\alpha, \beta, \gamma, \delta, \epsilon) = 1 + \frac{\alpha}{\underline{1}} \cdot \frac{\beta}{\underline{1}} \cdot \frac{\gamma}{\underline{1}} \cdot \frac{1}{\epsilon} + \frac{\alpha(\alpha+1)}{\underline{1}} \cdot \frac{\beta(\beta+1)}{\underline{1}} \cdot \frac{\gamma(\gamma+1)}{\underline{1}} \cdot \frac{1}{\epsilon(\epsilon+1)} + \dots, \text{ then}$$

$$i. F(\alpha, \beta, \gamma, \delta, \epsilon) = \frac{\underline{\delta-1} \underline{\delta-\alpha-\beta-1}}{\underline{\delta-\alpha-1} \underline{\delta-\beta-1}} F(\alpha, \beta, \epsilon-\gamma, \alpha+\beta-\delta+1, \epsilon)$$

$$+ \frac{\underline{\delta-1} \underline{\epsilon-1} \underline{\alpha+\beta-\delta-1} \underline{\delta+\epsilon-\alpha-\beta-\gamma-1}}{\underline{\alpha-1} \underline{\beta-1} \underline{\epsilon-\gamma-1} \underline{\delta+\epsilon-\alpha-\beta-1}} F(\delta-\alpha, \delta-\beta, \delta+\epsilon-\alpha-\beta-\gamma, \delta-\alpha-\beta+1, \delta+\epsilon-\alpha-\beta).$$

ii For integral values of  $\alpha, \beta$  or  $\gamma$ ,

$$F(-2\alpha, -2\beta, -\gamma, -\alpha - \beta + \frac{1}{2}, \delta)$$

$$= F(-\alpha, -\beta, -\gamma, \gamma + \delta, -\alpha - \beta + \frac{1}{2}, \frac{\delta}{2}, \frac{\delta+1}{2})$$

30. If  $\alpha + \beta + 1 = \gamma + \delta$

$$\text{and } y = \frac{1 - \alpha - 1 \cdot \beta - 1}{\gamma - 1 \cdot \delta - 1} \cdot \frac{1 + \frac{\alpha}{u} \cdot \frac{\beta}{\delta} (1-x) + \frac{\alpha(\alpha+1)}{u^2} \cdot \frac{\beta(\beta+1)}{\delta(\delta+1)} (1-x)^2 + \dots}{1 + \frac{\alpha}{u} \cdot \frac{\beta}{\gamma} x + \frac{\alpha(\alpha+1)}{u^2} \cdot \frac{\beta(\beta+1)}{\gamma(\gamma+1)} x^2 + \dots}$$

then,

$$\frac{dy}{dx} = - \frac{1}{\left\{ 1 + \frac{\alpha}{u} \cdot \frac{\beta}{\gamma} x + \frac{\alpha(\alpha+1)}{u^2} \cdot \frac{\beta(\beta+1)}{\gamma(\gamma+1)} x^2 + \dots \right\}^2} \cdot \frac{1}{x^\gamma (1-x)^\delta}$$

$$\text{Cor. If } y = \frac{\pi}{\text{Sin} \pi n} \cdot \frac{1 + \frac{n}{u} \cdot \frac{1-n}{u} (1-x) + \frac{n(n+1)(1-n)(2-n)}{u^2 u^2} (1-x)^2 + \dots}{1 + \frac{n}{u} \cdot \frac{1-n}{u} x + \frac{n(n+1)(1-n)(2-n)}{u^2 u^2} x^2 + \dots}$$

then  $\frac{dy}{dx} =$

$$- \frac{1}{x(1-x) \left\{ 1 + \frac{n}{u} \cdot \frac{1-n}{u} x + \frac{n(n+1)(1-n)(2-n)}{u^2 u^2} x^2 + \dots \right\}^2}$$

31. If  $y = 1 + \frac{\alpha}{u} \cdot \frac{\beta}{\gamma} x + \frac{\alpha(\alpha+1)}{u^2} \cdot \frac{\beta(\beta+1)}{\gamma(\gamma+1)} x^2 + \dots$ , then

$$i. (\alpha-1)(\beta-1) \int y dx - x(1-x) \frac{dy}{dx} = (\gamma-1)(y-1) (\alpha+\beta-1) xy$$

$$ii. y \int \frac{x^{n-2} y dx}{x^\gamma (1-x)^\delta y^2} dx \text{ (where } \delta = \alpha + \beta + 1 - \gamma)$$

$$= \frac{x^{n-\gamma} (1-x)^{1-\delta}}{(n-\gamma)(n-1)} \cdot \left[ 1 + \frac{(n-\alpha)(n-\beta)}{n(n-\gamma+1)} x + \frac{(n-\alpha)(n-\alpha+1)(n-\beta)(n-\beta+1)}{n(n+1)(n-\gamma+1)(n-\gamma+2)} x^2 + \dots \right]$$

Cor. If  $y = 1 + \frac{n}{u} \cdot \frac{1-n}{u} x + \frac{n(n+1)}{u^2} \cdot \frac{(1-n)(2-n)}{u^2} x^2 + \dots$ , then

$$x(x-1) \frac{dy}{dx} = n(n-1) \int y dx$$

$$2. i. 1 + \left(\frac{1}{2}\right)^2 \left\{ 1 - \frac{\phi(x)}{\phi(x)} \right\} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left\{ 1 - \frac{\phi(x)}{\phi(x)} \right\}^2 + \dots$$

=  $\sqrt{\phi(x)}$  x an even function of  $x$  whatever be  $\phi(x)$ .

$$\text{ii. } 1 + \left(\frac{1}{2}\right)^x \left(1 - \frac{x}{2}\right) + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(1 - \frac{x}{2}\right)^2 + \&c$$

$$= \sqrt{x} \left\{ 1 + \left(\frac{1}{2}\right)^x \left(1 - x\right) + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(1 - x\right)^2 + \&c \right\}$$

$$\text{iii. } 1 + \left(\frac{1}{2}\right)^x \left\{ 1 - \left(\frac{x}{1+x}\right)^2 \right\} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left\{ 1 - \left(\frac{x}{1+x}\right)^2 \right\}^2 + \&c$$

$$= (1+x) \left\{ 1 + \left(\frac{1}{2}\right)^x x^2 + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x x^4 + \&c \right\}$$

$$\text{iv. } 1 + \left(\frac{1}{2}\right)^x \left\{ 1 - \left(\frac{x}{1+x}\right)^4 \right\} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left\{ 1 - \left(\frac{x}{1+x}\right)^4 \right\}^2 + \&c$$

$$= (1+x)^2 \left\{ 1 + \left(\frac{1}{2}\right)^x x^4 + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x x^8 + \&c \right\}$$

$$\text{v. } \sqrt{1+x^2} \left\{ 1 + \left(\frac{1}{2}\right)^x \frac{1+\frac{1}{2}x}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(\frac{1+\frac{1}{2}x}{2}\right)^2 + \&c \right\}$$

$$= \frac{1+\frac{1}{2}x}{2} \left\{ 1 + \left(\frac{1}{2}\right)^x \frac{1+\frac{x}{\sqrt{1+x^2}}}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(\frac{1+\frac{x}{\sqrt{1+x^2}}}{2}\right)^2 + \&c \right\}$$

$$+ \frac{1-\frac{1}{2}x}{2} \left\{ 1 + \left(\frac{1}{2}\right)^x \frac{1-\frac{x}{\sqrt{1+x^2}}}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(\frac{1-\frac{x}{\sqrt{1+x^2}}}{2}\right)^2 + \&c \right\}$$

$$\text{33. i. } 1 + \left(\frac{1}{2}\right)^x \frac{2x}{1+x} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(\frac{2x}{1+x}\right)^2 + \&c$$

$$= \sqrt{1+x} \left\{ 1 + \frac{1 \cdot 3}{4} x^2 + \frac{1 \cdot 3 \cdot 5 \cdot 7}{4 \cdot 8} x^4 + \&c \right\}$$

$$\text{ii. } 1 + \left(\frac{1}{2}\right)^x \frac{1-\sqrt{1-x}}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x \left(\frac{1-\sqrt{1-x}}{2}\right)^2 + \&c$$

$$= 1 + \left(\frac{1}{2}\right)^x x + \left(\frac{1 \cdot 3}{4 \cdot 8}\right)^x x^2 + \left(\frac{1 \cdot 3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12}\right)^x x^3 + \&c$$

$$\text{iii. } 1 + \left(\frac{1}{2}\right)^3 x + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^3 x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^3 x^3 + \&c$$

$$= \left\{ 1 + \left(\frac{1}{2}\right)^x x + \left(\frac{1 \cdot 3}{4 \cdot 8}\right)^x x^2 + \left(\frac{1 \cdot 3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12}\right)^x x^3 + \&c \right\}^2$$

$$\text{iv. } 1 + \frac{1 \cdot 3}{4} \frac{4x}{(1+x)^2} + \frac{1 \cdot 3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12} \left\{ \frac{4x}{(1+x)^2} \right\}^2 + \&c$$

$$= \sqrt{1+x} \left\{ 1 + \left(\frac{1}{2}\right)^x x + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^x x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^x x^3 + \&c \right\}$$

$$\text{v. } 1 + \left(\frac{1}{4}\right)^x x + \left(\frac{1 \cdot 3}{4 \cdot 8}\right)^x x^2 + \left(\frac{1 \cdot 3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12}\right)^x x^3 + \&c$$

$$= \sqrt{1-x} \left\{ 1 + \left(\frac{3}{4}\right)^x x + \left(\frac{3 \cdot 7}{4 \cdot 8}\right)^x x^2 + \left(\frac{3 \cdot 7 \cdot 11}{4 \cdot 8 \cdot 12}\right)^x x^3 + \&c \right\}$$

$$\text{ex. i. } 1 - \frac{1 \cdot 3}{4} \frac{4x}{(1-x)^2} + \frac{1 \cdot 3 \cdot 5 \cdot 7}{4 \cdot 8 \cdot 12} \left\{ \frac{4x}{(1-x)^2} \right\}^2 - \&c =$$

$$\sqrt{\frac{x}{1+x}} \left\{ 1 + \left(\frac{1}{2}\right)^2 \frac{x}{1+x} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{x}{1+x}\right)^2 + \dots \right\}$$

$$\text{ii. } 1 - \left(\frac{1}{2}\right)^2 \frac{x}{1-x} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left\{ \frac{x}{1-x} \right\}^2 - \dots$$

$$= \sqrt{1-x} \left\{ 1 + \left(\frac{1}{2}\right)^2 x + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 x^3 + \dots \right\}$$

$$\text{iii. } 1 - \left(\frac{1}{2}\right)^2 \frac{x}{1-x} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left\{ \frac{x}{1-x} \right\}^2 - \dots$$

$$= \frac{(1-x)\sqrt{1-x}}{1+x} \left\{ 1 + \left(\frac{1}{2}\right)^2 x + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 x^2 + \dots \right\}$$

34. If  $\pi \mu \eta = 1$  and  $\mu = \frac{\sqrt{\pi}}{\left(\frac{1}{2}\right)^{\mu}}$  such that

$$\sqrt{\mu} = 1.0864348112, 1380801457, 531612$$

$$\frac{1}{\sqrt{2}\eta} = 1.3110287771, 46060$$

$$\mu = 1.1803405990, 16092$$

$$\eta = .2696763005, 94191$$

$$\frac{1}{\eta} = 3.7081493546, 02731, \text{ then}$$

$$\text{i. } 1 + \left(\frac{1}{2}\right)^2 \frac{1+x}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{1+x}{2}\right)^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 \left(\frac{1+x}{2}\right)^3 + \dots$$

$$= \mu \left\{ 1 + \frac{1^2}{2 \cdot 4} x^2 + \frac{1^2 \cdot 3^2}{2 \cdot 4 \cdot 6 \cdot 8} x^4 + \frac{1^2 \cdot 3^2 \cdot 5^2}{2 \cdot 4 \cdot 6 \cdot 8 \cdot 10 \cdot 12} x^6 + \dots \right\}$$

$$+ \eta \left\{ x + \frac{3^2}{4 \cdot 6} x^3 + \frac{3^2 \cdot 7^2}{4 \cdot 6 \cdot 8 \cdot 10} x^5 + \frac{3^2 \cdot 7^2 \cdot 11^2}{4 \cdot 6 \cdot 8 \cdot 10 \cdot 12 \cdot 14} x^7 + \dots \right\}$$

$$\text{ii. } 1 + \left(\frac{1}{2}\right)^2 \left(\frac{1}{2} + \frac{x}{1+x}\right) + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{1}{2} + \frac{x}{1+x}\right)^2 + \dots$$

$$= \mu \sqrt{1+x} \left\{ 1 + \frac{1}{2} \cdot \frac{1}{3} x^2 + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{1 \cdot 5}{3 \cdot 7} x^4 + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{1 \cdot 5 \cdot 7}{3 \cdot 7 \cdot 11} x^6 + \dots \right\}$$

$$+ \eta \sqrt{1+x} \left\{ x + \frac{1}{2} \cdot \frac{3}{5} x^3 + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{3 \cdot 7}{5 \cdot 9} x^5 + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{3 \cdot 7 \cdot 11}{5 \cdot 9 \cdot 13} x^7 + \dots \right\}$$

$$\text{iii. } \frac{\pi}{4} \left\{ 1 + \left(\frac{1}{2}\right)^2 \frac{1+x}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{1+x}{2}\right)^2 + \dots \right\}^2$$

$$- \frac{\pi}{4} \left\{ 1 + \left(\frac{1}{2}\right)^2 \frac{1-x}{2} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{1-x}{2}\right)^2 + \dots \right\}^2$$

$$= x + \frac{2}{3} x^3 \left(1 - \frac{1^2}{2^2}\right) + \frac{2 \cdot 4}{3 \cdot 5} x^5 \left(1 - 2 \cdot \frac{1^2}{2^2} + \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2}\right) + \dots$$



$$\frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7} x^7 \left( 1 - 3 \cdot \frac{1^2}{2^2} + 3 \cdot \frac{1^2 \cdot 3^2}{2^2 \cdot 4^2} - \frac{1^2 \cdot 3^2 \cdot 5^2}{2^2 \cdot 4^2 \cdot 6^2} \right) + \dots$$

$$= x + \frac{x^3}{2} + \frac{61x^5}{120} + \frac{21x^7}{80} + \dots = \frac{x}{1-x^2} - \frac{1}{2} \cdot \frac{x^3}{(1-x^2)^2} + \frac{61}{120} \cdot \frac{x^5}{(1-x^2)^3} - \dots$$

ex. i.  $1 + \left(\frac{x}{2}\right)^{1+\frac{n}{2}} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{x}{2}\right)^{1+\frac{n}{2}} + \dots$

$$= \frac{1}{(1-x^2)^{\frac{1}{2}}} \left\{ 1 - \frac{1^2}{2 \cdot 4} \cdot \frac{x^2}{1-x^2} + \frac{1^2 \cdot 3^2}{2 \cdot 4 \cdot 6 \cdot 8} \left(\frac{x^2}{1-x^2}\right)^2 - \dots \right\}$$

$$+ \frac{7x}{(1-x^2)^{\frac{3}{2}}} \left\{ 1 - \frac{3^2}{4 \cdot 6} \cdot \frac{x^2}{1-x^2} + \frac{3^2 \cdot 7^2}{4 \cdot 6 \cdot 8 \cdot 10} \left(\frac{x^2}{1-x^2}\right)^2 - \dots \right\}$$

ii.  $1 + \left(\frac{x}{2}\right)^2 \left(\frac{x}{2} + \frac{x}{1+x^2}\right) + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{x}{2} + \frac{x}{1+x^2}\right)^2 + \dots$

$$= \frac{1}{\sqrt{1-x^2}} \left\{ 1 - \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{x^2}{1-x^2} + \frac{1 \cdot 3 \cdot 2 \cdot 6}{2 \cdot 4 \cdot 3 \cdot 7} \left(\frac{x^2}{1-x^2}\right)^2 - \dots \right\}$$

$$+ \frac{2 \cdot 7 \cdot x}{\sqrt{1-x^2}} \left\{ 1 - \frac{1}{2} \cdot \frac{2}{5} \cdot \frac{x^2}{1-x^2} + \frac{1 \cdot 3 \cdot 2 \cdot 6}{2 \cdot 4 \cdot 5 \cdot 7 \cdot 9} \left(\frac{x^2}{1-x^2}\right)^2 - \dots \right\}$$

35. i.  $\cos(2n \sin^{-1} x) = 1 - \frac{n}{1!} \cdot \frac{x}{2} x^2 + \frac{n(n-1)}{2!} \cdot \frac{n(n+1)}{2 \cdot 1 \cdot 2} x^4 - \dots$

ii.  $\frac{\sin(2n \sin^{-1} x)}{2n x} = 1 + \frac{\frac{1}{2} - n}{1!} \cdot \frac{\frac{1}{2} + n}{1 \cdot 2} x^2 + \frac{(\frac{1}{2} - n)(\frac{1}{2} - n - 1)}{2!} \cdot \frac{(\frac{1}{2} + n)(\frac{1}{2} + n + 1)}{1 \cdot 2 \cdot 2 \cdot 2} x^4 + \dots$

iii.  $\frac{\cos(2n \sin^{-1} x)}{\sqrt{1-x^2}} = 1 + \frac{\frac{1}{2} - n}{1!} \cdot \frac{\frac{1}{2} + n}{1 \cdot 2} x^2 + \frac{(\frac{1}{2} - n)(\frac{1}{2} - n - 1)}{2!} \cdot \frac{(\frac{1}{2} + n)(\frac{1}{2} + n + 1)}{2 \cdot 1 \cdot 2} x^4 + \dots$

36. i.  $(1+x)^n = 1 + n x (1+x)^{\frac{n-1}{2}} + \frac{n(n-1)}{4 \cdot 1^2} x^2 (1+x)^{\frac{n-2}{2}} + \frac{n(n-1)(n-3)}{4^2 \cdot 1^2} x^4 (1+x)^{\frac{n-4}{2}} + \dots$

ii.  $\frac{1 + (1+x)^n}{2} = (1+x)^{\frac{n}{2}} + \frac{n^2}{4 \cdot 1^2} x^2 (1+x)^{\frac{n-2}{2}} + \frac{n^2(n^2-2^2)}{4^2 \cdot 1^2} x^4 (1+x)^{\frac{n-4}{2}} + \dots$

iii.  $\left(\frac{1 + \sqrt{1+4x}}{2}\right)^n = 1 + n x (1+x)^{\frac{n-2}{2}} + \frac{n(n-2)(n-4)}{4 \cdot 1^2} x^2 (1+x)^{\frac{n-4}{2}} + \frac{n(n-2)(n-4)(n-6)(n-8)}{4^2 \cdot 1^2} x^4 (1+x)^{\frac{n-6}{2}} + \dots$

iv.  $\frac{1}{2} + \frac{1}{2} \left(\frac{1 + \sqrt{1+4x}}{2}\right)^n = (1+x)^{\frac{n}{2}} + \frac{n(n-4)}{4 \cdot 1^2} x^2 (1+x)^{\frac{n-4}{2}} + \frac{n(n-6)(n-10)}{4^2 \cdot 1^2} x^4 (1+x)^{\frac{n-6}{2}} + \dots$

$$1. \frac{a_1}{b_1} + \frac{a_2}{b_2} + \dots + \frac{a_n}{b_n} = a_1 \frac{N_{n-1}}{D_n} = \frac{a_1}{D_0 D_1} - \frac{a_1 a_2}{D_1 D_2} + \frac{a_1 a_2 a_3}{D_2 D_3} - \dots \&c$$

to  $n$  terms, where

$$N_{n-1} = b_n N_{n-2} + a_n N_{n-3} \text{ and } D_n = b_n D_{n-1} + a_n D_{n-2}.$$

$$\text{Cor. } a_1 + a_2 + a_3 + \dots \text{ to } n \text{ terms} = \frac{a_1}{1} - \frac{a_1 a_2}{a_1 + a_2} - \frac{a_1 a_2 a_3}{a_2 + a_3} - \frac{a_2 a_3 a_4}{a_3 + a_4} -$$

$$\frac{a_3 a_4 a_5}{a_4 + a_5} - \dots \&c \text{ to } n \text{ terms.}$$

$$2. x = (x - a_1) + \frac{x a_1}{x - a_2} + \frac{x a_2}{x - a_3} + \frac{x a_3}{x - a_4} + \dots \&c.$$

$$3. x = a_1 + \sqrt{x^2 + a_1(a_1 + 2a_2)} - 2a_1 \sqrt{x^2 + a_2(a_2 + 2a_3)} - 2a_3 \sqrt{\dots} \&c$$

$$4. x + n + a = \sqrt{ax + (n+a)^2} + x \sqrt{a(x+n) + (n+a)^2} + (x+n) \sqrt{\dots} \&c$$

e.g. i.  $3 = 1\sqrt{1+2} + 2\sqrt{1+3} + 3\sqrt{1+4} + \dots \&c$

ii.  $4 = 1\sqrt{6+2} + 2\sqrt{7+3} + 3\sqrt{8+4} + 4\sqrt{9+5} + \dots \&c.$

$$5. i. 2 \cos \theta = \sqrt{2 + 2 \cos 2\theta} = \sqrt{2 + \sqrt{2 + 2 \cos 4\theta}} = \sqrt{2 + \sqrt{2 + \sqrt{2 + 2 \cos 8\theta}}} \dots \&c.$$

$$ii. 2 \cos \theta = \sqrt[3]{2 \cos 3\theta} + \sqrt[3]{2 \cos 3\theta} + \sqrt[3]{2 \cos 3\theta} + \dots \&c.$$

$$= \sqrt[3]{6 \cos \theta} + \sqrt[3]{6 \cos 3\theta} + \sqrt[3]{6 \cos 9\theta} + \sqrt[3]{6 \cos 27\theta} + \dots \&c$$

$$6. \sqrt{\frac{a(a-2)}{h}} + \sqrt{\frac{a(a-2)}{h}} + \sqrt{\frac{a(a-4)}{h}} + \dots \text{ to } n \text{ terms} + h.$$

$$= \frac{a}{2} \left\{ 1 - \frac{v}{ax} + \frac{(v/ax)^2}{2(a-1)} - \frac{(v/ax)^3}{2(a-1)(a^2-1)} + \frac{(v/ax)^4(a+5)}{8(a-1)(a^2-1)(a^2-1)} \right.$$

$$\left. - \frac{(v/ax)^5(2a^2+3a+7)}{8(a-1)(a^2-1)(a^2-1)(a^2-1)} + \dots \right\} \text{ where } v \text{ is a function}$$

of  $a$  and  $h$  independent of  $x$  defined by the relation

$$\frac{2h}{a} = 1 - v + \frac{v^2}{2(a-1)} - \frac{v^3}{2(a-1)(a^2-1)} + \frac{v^4(a+5)}{8(a-1)(a^2-1)(a^2-1)} - \dots \&c$$

the coefft. of  $v^{n+1} = \frac{1}{2(a^n-1)}$  x the coefft. of  $v^n$  in the square of the series

$$7. x = \frac{x+1}{x} + \frac{x+2}{x+1} + \frac{x+2}{x+2} + \dots \&c. \text{ Cor. } 1 = \frac{2}{1} + \frac{2}{2} + \frac{4}{3} + \frac{5}{4} + \dots \&c.$$

14.0.  
8.  $\frac{1}{x+a} - \frac{1}{(x+a)(x+2a)} + \frac{1}{(x+a)(x+2a)(x+3a)} - \dots$  to  $n$  terms

$$= \frac{1}{x+a} + \frac{x+a}{x+2a-1} + \frac{x+2a}{x+3a-1} + \frac{x+3a}{x+4a-1} + \dots$$

Cor.  $\frac{1}{e-1} = \frac{1}{1} + \frac{2}{2} + \frac{3}{3} + \frac{4}{4} + \dots$

9.  $\frac{x+a+1}{x+1} = \frac{x+a}{x-1} + \frac{x+2a}{x+1} + \frac{x+3a}{x+2a-1} + \dots$

e.g. 1.  $\frac{4}{3} = \frac{3}{1} + \frac{4}{2} + \frac{5}{3} + \frac{6}{4} + \dots$

2.  $\frac{5}{3} = \frac{4}{1} + \frac{6}{3} + \frac{8}{5} + \frac{10}{7} + \dots$

10. If  $n$  is a positive integer,

$$n = \frac{1}{1-n} + \frac{2}{2-n} + \frac{3}{3-n} + \dots + \frac{n}{0} + \frac{n+1}{1} + \frac{n+2}{2} + \frac{n+3}{3} + \dots$$

11. If  $a$  is a positive integer and  $D = \phi(n-1)$  where  $\phi(n) = N$  where  $N_{n+1}$  and  $N_n$  are the numerator and the denominator in the fraction

$$n+2-a + \frac{a-1}{n+3-a} + \frac{a-2}{n+4-a} + \frac{a-3}{n+5-a} + \dots$$

Cor. 1.  $\frac{n^4+n+1}{n^2-n+1} = \frac{n}{n-3} + \frac{n+1}{n-2} + \frac{n+2}{n-1} + \frac{n+3}{n} + \dots$

2.  $\frac{x^3+2x+1}{(x-1)^2+(x-1)+1} = \frac{x}{x-4} + \frac{x+1}{x-3} + \frac{x+2}{x-2} + \frac{x+3}{x-1} + \dots$

12.  $1 = \frac{x+a}{a} + \frac{(x+a)^2-a^2}{a} + \frac{(x+2a)^2-a^2}{a} + \frac{(x+3a)^2-a^2}{a} + \dots$

13. If  $a < b$ ,  $a = \frac{ab}{a+b+d} - \frac{(a+d)(b+d)}{a+b+3d} + \frac{(a+2d)(b+2d)}{a+b+5d} - \dots$

14.  $\frac{a_1}{x} + \frac{a_2}{1} + \frac{a_3}{x} + \frac{a_4}{1} + \dots$  to  $2n$  terms

$$= \frac{a_1}{x+a_2} - \frac{a_2 a_3}{x+a_2+a_4} + \frac{a_4 a_5}{x+a_2+a_6} - \dots$$

15.  $\frac{a_1+h}{1} + \frac{a_1}{x} + \frac{a_2+h}{1} + \frac{a_2}{x} + \frac{a_3+h}{1} + \dots$

$$= h + \frac{a_1}{1} + \frac{a_1+h}{x} + \frac{a_2}{1} + \frac{a_2+h}{x} + \frac{a_3}{1} + \dots$$

16.  $\frac{1}{(m+1)(m+2)} - \frac{1}{(m+2)(m+3)} + \frac{1}{(m+3)(m+4)} - \dots$

$$= \frac{1}{m+1} + \frac{(m+1)^L (m+1)^L}{m+1+3+} + \frac{(m+1)^L (m+1)^L}{m+1+} \&c.$$

17.  $\frac{1}{1+} \frac{a_1 x}{1+} \frac{a_2 x}{1+} \frac{a_3 x}{1+} \&c = 1 - A_1 x + A_2 x^2 - A_3 x^3 + \&c$

let  $P_n = a_1 a_2 a_3 \dots a_{n-1} (a_1 + a_2 + \dots + a_n)$ , then

$$P_1 = A_1; P_2 = A_2; P_3 = A_3 - a_1 A_2; P_4 = A_4 - (a_1 + a_2) A_3$$

$$P_5 = A_5 - (a_1 + a_2 + a_3) A_4 + a_1 a_3 A_3$$

$$P_6 = A_6 - (a_1 + a_2 + a_3 + a_4) A_5 + (a_1 a_3 + a_2 a_4 + a_1 a_4) A_4$$

$$P_n = \phi_0(n) A_n - \phi_1(n) A_{n-1} + \phi_2(n) A_{n-2} - \&c$$

where  $\phi_n(m+1) - \phi_n(m) = a_{m-1} \phi_{n-1}(m-1)$ .

Cor. I. If  $\frac{1}{1+b_1 x} + \frac{a_1 x}{1+b_2 x} + \frac{a_2 x}{1+b_3 x} + \&c = 1 - A_1 x + A_2 x^2 - \&c$

$$P_n = a_1 a_2 a_3 \dots a_{n-1} (a_1 + b_1 + a_2 + b_2 + \dots + a_n + b_n)$$

$$= \phi_0(n) A_n - \phi_1(n) A_{n-1} + \phi_2(n) A_{n-2} - \&c \text{ where}$$

$$\phi_n(m+1) - \phi_n(m) = b_m \phi_{n-1}(m) + a_{m-1} \phi_{n-1}(m-1)$$

Cor. II. In the above results  $D_{n-1} = \phi_0(n) + x \phi_1(n) + x^2 \phi_2(n) + \&c$

ex.  $\left\{ 1 + \left(\frac{1}{2}\right)^n x + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^n x^2 + \&c \right\}^2 = \frac{1}{1} - \frac{x}{2} - \frac{3x}{8} - \frac{5x}{2} - \frac{17x}{40}$

N.B. The peculiarity in this continued fraction is if  $x=1$

it assumes the form  $1 + 1 + \frac{3}{5} + \frac{3}{5} + \&c$ .

18.  $\frac{(x+1)^n - (x-1)^n}{(x+1)^n + (x-1)^n} = \frac{x}{x} + \frac{n^2-1^2}{3x} + \frac{n^2-3^2}{5x} + \frac{n^2-5^2}{7x} + \&c$

N.B. If  $V_n$  denotes the above fraction, then  $V_n + \frac{1}{V_n} = \frac{2}{\sqrt{2}}$

Cor. I.  $\tan^{-1} x = \frac{x}{1} + \frac{(x)^2}{3} + \frac{(2x)^2}{5} + \frac{(3x)^2}{7} + \frac{6x^2}{9} + \&c$

Cor. 2.  $\log \frac{1+x}{1-x} = \frac{2x}{1} - \frac{(x)^2}{3} + \frac{(2x)^2}{5} - \frac{(3x)^2}{7} + \dots$

3.  $\tan x = \frac{x}{1} - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$

4.  $\frac{e^x}{e^x+1} = \frac{x}{2} + \frac{x^3}{6} + \frac{x^5}{10} + \frac{x^7}{14} + \dots$

19. 
$$\frac{\frac{x}{n} + \frac{x^2}{4} \cdot \frac{1}{n(n+1)} + \frac{x^3}{8} \cdot \frac{1}{n(n+1)(n+2)} + \dots}{1 + \frac{x}{4} \cdot \frac{1}{n} + \frac{x^2}{8} \cdot \frac{1}{n(n+1)} + \dots}$$

$$= \frac{x}{n} + \frac{x}{n+1} + \frac{x}{n+2} + \frac{x}{n+3} + \dots$$

20. d. 
$$\frac{\frac{\beta}{\gamma}x + \frac{d-\gamma}{4} \cdot \frac{\beta(\beta+1)}{\gamma(\gamma+1)}x^2 + \frac{(\alpha-\gamma)(\alpha-\gamma-1)}{12} \cdot \frac{\beta(\beta+1)(\beta+2)}{\gamma(\gamma+1)(\gamma+2)}x^3 + \dots}{1 + \frac{d-\gamma}{4} \cdot \frac{\beta}{\gamma}x + \frac{(\alpha-\gamma)(\alpha-\gamma-1)}{12} \cdot \frac{\beta(\beta+1)}{\gamma(\gamma+1)}x^2 + \dots}$$

$$= \frac{\alpha \beta}{\gamma} + \frac{\alpha(d-\gamma)(\beta-\gamma)}{\gamma+1} + \frac{\alpha(\alpha+1)(\beta+1)}{\gamma+2} + \frac{\alpha(\alpha-\gamma-1)(\beta-\gamma-1)}{\gamma+3} + \dots$$

21. 
$$\frac{\beta}{\gamma}x - \frac{\beta(\beta+1)}{\gamma(\gamma+1)}x^2 + \frac{\beta(\beta+1)(\beta+2)}{\gamma(\gamma+1)(\gamma+2)}x^3 - \dots$$

$$= \frac{\beta x}{\gamma} + \frac{\gamma(\beta+1)x}{\gamma+1} + \frac{1(\gamma-\beta)x}{\gamma+2} + \frac{(\gamma+1)(\beta+2)x}{\gamma+3} + \frac{2(\gamma-\beta+1)x}{\gamma+4} + \dots$$

$$= \frac{\beta x}{\gamma} + \frac{(\beta+1)x}{1} + \frac{1(1+x)}{\gamma} + \frac{(\beta+2)x}{1} + \frac{2(1+x)}{\gamma} + \dots$$

$$= \frac{\beta x}{\gamma + x(\beta+1)} - \frac{1(\beta+1)x(1+x)}{\gamma+1+x(\beta+3)} - \frac{2(\beta+2)x(1+x)}{\gamma+2+x(\beta+5)} - \dots$$

Cor. 1 
$$\frac{x}{n} + \frac{x^2}{n(n+1)} + \frac{x^3}{n(n+1)(n+2)} + \dots$$

$$= \frac{x}{n} - \frac{nx}{n+1} + \frac{x}{n+2} - \frac{(n+1)x}{n+3} + \frac{2x}{n+4} - \dots$$

$$= \frac{x}{n-x} + \frac{x}{n+1-x} + \frac{2x}{n+2-x} + \frac{3x}{n+3-x} + \dots$$

Cor. 2. 
$$1 + \frac{x}{2+1} + \frac{x^2}{(2+1)(2+2)} + \frac{x^3}{(2+1)(2+2)(2+3)} + \dots$$

$$= 1 + \frac{2x}{2} + \frac{3x}{3} + \frac{4x}{4} + \frac{5x}{5} + \frac{6x}{6} + \dots$$

$$22. \frac{\frac{3}{7}x + \frac{\alpha}{\Gamma} \cdot \frac{\beta(\beta+1)}{\Gamma(\Gamma+1)} x^\Gamma + \frac{\alpha(\alpha-1)}{\Gamma^2} \cdot \frac{\beta(\beta+1)(\beta+2)}{\Gamma(\Gamma+1)(\Gamma+2)} x^{\Gamma+2} + \dots}{1 + \frac{\alpha}{\Gamma} \cdot \frac{\beta}{\Gamma} x + \frac{\alpha(\alpha-1)}{\Gamma^2} \cdot \frac{\beta(\beta+1)}{\Gamma(\Gamma+1)} x^2 + \dots}$$

$$= \frac{\beta x}{\Gamma - (\alpha + \beta + 1)x + \Gamma + 1 - (\alpha + \beta + 2)x + \frac{(\beta+1)(\alpha + \Gamma + 1)x}{\Gamma + 1} - (\alpha + \beta + 3)x + \dots}$$

$$23. \frac{a_n}{b_n x} + \frac{a_{n+1}}{b_{n+1} x} + \frac{a_{n+2}}{b_{n+2} x} + \dots = C_n(1 - P_n x + Q_n x^2 - R_n x^3 + \dots)$$

where  $C_n C_{n+1} = a_n$ ;  $P_n + P_{n+1} = \frac{b_n}{C_{n+1}}$  or  $\frac{b_n C_n}{a_n}$ ;  
 $Q_n + Q_{n+1} = (P_n)^2$ ;  $R_n + R_{n+1} = P_n(Q_n - Q_{n+1})$ ;  
 $S_n + S_{n+1} = P_n(R_n - R_{n+1}) - Q_n Q_{n+1}$ ; generally  
 $Z_n + Z_{n+1} = P_n(Y_n - Y_{n+1}) - Q_n X_{n+1} - R_n W_{n+1}$   
 $- S_n V_{n+1} - \dots - X_n Q_{n+1}$ .

N.B. In some cases the above theorem is only approximately true.

ex.  $\sqrt{\frac{2x}{\pi}} = \frac{x}{1} + \frac{2x}{2} + \frac{3x}{3} + \frac{4x}{4} + \dots = \frac{2}{3\pi}$  when  $x = \infty$ .

$$24. \frac{n}{n} + \frac{x}{n+1} + \frac{x}{n+2} + \frac{x}{n+3} + \dots + \frac{x}{n+r}$$

$$= \left\{ 1 + \frac{x}{\Gamma} \cdot \frac{n-1}{(n+1)(n+r)} + \frac{x^2}{\Gamma^2} \cdot \frac{(n-2)(n-3)}{(n+1)(n+2)(n+r)(n+r-1)} \right.$$

$$\left. + \frac{x^3}{\Gamma^3} \cdot \frac{(n-3)(n-4)(n-5)}{(n+1)(n+2)(n+3)(n+r)(n+r-1)(n+r-2)} + \dots \right\}$$

$$\div \left\{ 1 + \frac{x}{\Gamma} \cdot \frac{n}{n(n+r)} + \frac{x^2}{\Gamma^2} \cdot \frac{(n-1)(n-2)}{n(n+1)(n+2)(n+r)} + \dots \right\}$$

the no. of terms being limited.

$$25. \frac{\frac{x+n-3}{4} \frac{x-n-3}{4}}{\frac{x+n-1}{4} \frac{x-n-1}{4}} = \frac{4}{x} \cdot \frac{x^2-1}{2x} \cdot \frac{x^2-9}{2x} \cdot \frac{x^2-25}{2x} \dots$$

164.

$$\text{Cor. 1. } \left( \frac{\frac{x-3}{4}}{\frac{x-1}{4}} \right)^2 = \frac{4}{x} + \frac{12}{2x} + \frac{3^2}{2x} + \frac{5^2}{2x} + \frac{7^2}{2x} + \&c$$

$$\text{Cor. 2. } \frac{\frac{x-5}{8} \frac{x-7}{8}}{\frac{x-1}{8} \frac{x-3}{8}} = \frac{8}{x} + \frac{1 \cdot 3}{2x} + \frac{5 \cdot 7}{2x} + \frac{9 \cdot 11}{2x} + \&c$$

$$26. \left\{ \frac{\frac{x+n-3}{4} \frac{x-n-3}{4}}{\frac{x+n-1}{4} \frac{x-n-1}{4}} \right\}^2 = \frac{8}{x^2 + \frac{n^2-1}{2}} + \frac{1^2 - n^2}{1} + \frac{1^4}{x^2 - 1} + \frac{8^2 - n^2}{1} + \frac{3^2}{x^2 - 1}$$

$$= \frac{8}{x^2 - \frac{n^2-1}{2}} + \frac{1^2}{1} + \frac{1^2 - n^2}{x^2 - 1} + \frac{8^2}{1} + \frac{3^2 - n^2}{x^2 - 1} + \&c$$

$$\text{Cor. } \left\{ \frac{\frac{x-1}{4}}{\frac{x-1}{4}} \right\}^2 = \frac{8}{x^2 - 1} + \frac{1^2}{1} + \frac{1^2}{x^2 - 1} + \frac{3^2}{1} + \frac{3^2}{x^2 - 1} + \&c$$

$$27. x + \frac{(1+y)^2 + n}{2x} + \frac{(3+y)^2 + n}{2x} + \frac{(5+y)^2 + n}{2x} + \&c$$

$$= y + \frac{(1+x)^2 + n}{2y} + \frac{(3+x)^2 + n}{2y} + \frac{(5+x)^2 + n}{2y} + \&c$$

$$28. x + \frac{n^2 + 1^2}{2x} + \frac{n^2 + 3^2}{2x} + \frac{n^2 + 5^2}{2x} + \&c$$

$$= n + \frac{x^2 - 1^2}{2n} + \frac{x^2 - 3^2}{2n} + \frac{x^2 - 5^2}{2n} + \&c \text{ approximately if } n \text{ is great.}$$

$$29. \left( \frac{1}{x+n+1} - \frac{1}{x+n+3} + \frac{1}{x+n+5} - \&c \right)$$

$$+ \left( \frac{1}{x-n+1} - \frac{1}{x-n+3} + \frac{1}{x-n+5} - \&c \right)$$

$$= \frac{1}{x} + \frac{1^2 - n^2}{x} + \frac{2^2}{x} + \frac{3^2 - n^2}{x} + \frac{4^2}{x} + \frac{5^2 - n^2}{x} + \&c$$

$$\text{Cor. } 2 \left( \frac{1}{x+1} - \frac{1}{x+3} + \frac{1}{x+5} - \&c \right) = \frac{1}{x} + \frac{1^2}{x} + \frac{2^2}{x} + \frac{3^2}{x} + \&c$$

$$30. \left( \frac{1}{x-n+1} + \frac{1}{x-n+3} + \frac{1}{x-n+5} + \&c \right)$$

$$- \left( \frac{1}{x+n+1} + \frac{1}{x+n+3} + \frac{1}{x+n+5} + \&c \right)$$

$$= \frac{n}{x} + \frac{1^2(1^2 - n^2)}{3x} + \frac{2^2(2^2 - n^2)}{5x} + \frac{3^2(3^2 - n^2)}{7x} + \&c$$

$$\text{Cor. } 2 \left\{ \frac{1}{(x+1)^2} + \frac{1}{(x+3)^2} + \frac{1}{(x+5)^2} + \&c \right\} = \frac{1}{x} + \frac{1^2}{x} + \frac{2^2}{5x} + \frac{3^2}{7x} + \&c$$

$$31. \left( \frac{1}{x-n+1} - \frac{1}{x-n+3} + \frac{1}{x-n+5} - \dots \right) \\ - \left( \frac{1}{x+n+1} - \frac{1}{x+n+3} + \frac{1}{x+n+5} - \dots \right) \\ = \frac{n}{x^2-1} + \frac{2^2 \cdot n^2}{1} + \frac{2^2}{x^2-1} + \frac{4^2 \cdot n^2}{1} + \frac{4^2}{x^2-1} + \dots$$

$$\text{Cor } 2 \left\{ \frac{1}{(x+1)^2} - \frac{1}{(x+3)^2} + \frac{1}{(x+5)^2} - \dots \right\} \\ = \frac{1}{x^2-1} + \frac{2^2}{1} + \frac{2^2}{x^2-1} + \frac{4^2}{1} + \frac{4^2}{x^2-1} + \dots$$

$$32. i. 2x \left( \frac{1}{2x} - \frac{1}{x+2} + \frac{1}{2+4} - \frac{1}{x+6} + \dots \right) \\ = \frac{1}{x} + \frac{1 \cdot 2}{x} + \frac{2 \cdot 3}{x} + \frac{3 \cdot 4}{x} + \frac{4 \cdot 5}{x} + \dots$$

$$ii. 2x^2 \left\{ \frac{1}{2x^2} - \left( \frac{1}{x+1} \right)^2 + \left( \frac{1}{x+2} \right)^2 - \left( \frac{1}{x+3} \right)^2 + \dots \right\} \\ = \frac{1}{x} + \frac{1 \cdot 2}{x} + \frac{1 \cdot 2}{x} + \frac{2^2}{x} + \frac{2 \cdot 3}{x} + \frac{3^2}{x} + \dots$$

$$iii. \frac{1}{(x+1)^3} + \frac{1}{(x+2)^3} + \frac{1}{(x+3)^3} + \dots \\ = \frac{1}{2x(x+1)} + \frac{1^3}{1} + \frac{1^3}{6x(x+1)} + \frac{2^3}{1} + \frac{2^3}{10x(x+1)} + \dots \\ = \frac{1}{2x^2+2x+1} - \frac{1^3}{3(2x^2+2x+3)} - \frac{2^3}{5(2x^2+2x+5)} - \frac{3^3}{7(2x^2+2x+7)} - \dots$$

$$A3. \frac{\left[ \frac{x+m+n-1}{2} \right] \left[ \frac{x-m-n-1}{2} \right] - \left[ \frac{x+m-n-1}{2} \right] \left[ \frac{x-m+n-1}{2} \right]}{\left[ \frac{x+m+n-1}{2} \right] \left[ \frac{x-m-n-1}{2} \right] + \left[ \frac{x+m-n-1}{2} \right] \left[ \frac{x-m+n-1}{2} \right]} \\ = \frac{mn}{x} + \frac{(m^2-1^2)(n^2-1^2)}{3x} + \frac{(m^2-2^2)(n^2-2^2)}{5x} + \frac{(m^2-3^2)(n^2-3^2)}{7x} + \dots$$

$$34. \text{Pf } P = \frac{\left[ \frac{x+l+n-3}{4} \right] \left[ \frac{x+l-n-3}{4} \right] \left[ \frac{x-l+n-1}{4} \right] \left[ \frac{x-l-m-1}{4} \right]}{\left[ \frac{x-l+n-3}{4} \right] \left[ \frac{x-l-n-3}{4} \right] \left[ \frac{x+l+n-1}{4} \right] \left[ \frac{x+l-m-1}{4} \right]}$$



146.

$$\text{then } \frac{1-P}{1+P} = \frac{l}{x+l} + \frac{l^2-n^2}{x+l} + \frac{2^2-l^2}{x+l} + \frac{3^2-n^2}{x+l} + \frac{4^2-l^2}{x+l} + \dots$$

$$\text{Cor. If } F(\alpha, \beta) = \tan^{-1} \frac{\alpha}{x+\alpha} + \frac{\beta^2+\gamma^2}{x+\beta} + \frac{\alpha^2+(2\gamma)^2}{x+\alpha} + \frac{\beta^2+(3\gamma)^2}{x+\beta} + \dots$$

and  $A$  be the average of  $\alpha$  &  $\beta$ , then  $F(A, A)$  is the average of  $F(\alpha, \beta)$  and  $F(\beta, \alpha)$ .

$$35. \text{ If } P = \frac{\left| \frac{x+l+m+n-1}{2} \right| \left| \frac{x+l-m-n-1}{2} \right| \left| \frac{x+m-n-l-1}{2} \right| \left| \frac{x+n-l-m-1}{2} \right|}{\left| \frac{x-l-m-n-1}{2} \right| \left| \frac{x-l+m+n-1}{2} \right| \left| \frac{x-m+n+l-1}{2} \right| \left| \frac{x-n+l+m-1}{2} \right|}$$

$$\text{then } \frac{1-P}{1+P} = \frac{2lmn}{x^2-l^2-m^2-n^2+1} + \frac{4(l^2-1^2)(m^2-1^2)(n^2-1^2)}{3(x^2-l^2-m^2-n^2+5)} + \frac{4(l^2-2^2)(m^2-2^2)(n^2-2^2)}{5(x^2-l^2-m^2-n^2+9)} + \dots$$

$$= \frac{2lmn}{y+l-2l^2m} + \frac{2(1-m)(1^2-n^2)}{1+l} + \frac{2(1+m)(1^2-l^2)}{3y+l} + \frac{2(l-m)(l^2-n^2)}{1+l} + \frac{2(2+m)(2^2-l^2)}{5y+l} + \dots \quad \text{where } y = x^2 - (1-m)^2 \text{ \& } l = (x^2-l^2)(1-2m)$$

$$36. \text{ If } P = \frac{\left| \frac{x+l+n-1}{4} \right| \left| \frac{x+l-n-3}{4} \right| \left| \frac{x-l+n-3}{4} \right| \left| \frac{x-l-n-1}{4} \right|}{\left| \frac{x-l+n-1}{4} \right| \left| \frac{x-l-n-3}{4} \right| \left| \frac{x+l-n-1}{4} \right| \left| \frac{x+l+n-3}{4} \right|}$$

$$\text{then } \frac{1-P}{1+P} = \frac{ln}{x^2-1-l^2} + \frac{2^2-n^2}{1+l} + \frac{2^2-l^2}{x^2-1+l} + \frac{4^2-n^2}{1+l} + \frac{4^2-l^2}{x^2-1+l} + \dots$$

$$37. \text{ If } \phi(y) = \frac{1}{y+1} + \frac{1}{y+3} + \frac{1}{y+5} + \dots, \text{ then}$$

$$\phi(x-l-n) - \phi(x+l-n) + \phi(x+l+n) - \phi(x-l+n)$$

$$= \frac{2ln}{x^2-1+n^2-l^2} + \frac{2(1^2-n^2)}{1+l} + \frac{2(1^2-l^2)}{3(x^2-1)+m^2-l^2} + \frac{4(2^2-n^2)}{1+l}$$

$$38. \left\{ \frac{1}{(x-n)^2} + \frac{1}{(x-n+1)^2} + \frac{1}{(x-n+2)^2} + \frac{1}{(x-n+3)^2} + \dots \right\}$$

$$- \left\{ \frac{1}{(x+n)^2} + \frac{1}{(x+n+1)^2} + \frac{1}{(x+n+2)^2} + \frac{1}{(x+n+3)^2} + \dots \right\}$$

$$= \frac{\pi}{x^2-n^2+1} + \frac{2(1^2-n^2)}{1} + \frac{\pi}{3(x^2-n^2)+1} + \frac{4(2^2-n^2)}{1} + \dots$$

$$= \frac{\pi}{x^2-n^2+1} - \frac{4(1^2-n^2)1^4}{3(x^2-n^2+5)} - \frac{4(2^2-n^2)2^4}{5(x^2-n^2+9)} - \dots$$

$$39. \frac{\left| \frac{x+l+n-3}{4} \right| \left| \frac{x-l+n-3}{4} \right| \left| \frac{x+l-n-3}{4} \right| \left| \frac{x-l-n-3}{4} \right|}{\left| \frac{x+l+n-1}{4} \right| \left| \frac{x-l+n-1}{4} \right| \left| \frac{x+l-n-1}{4} \right| \left| \frac{x-l-n-1}{4} \right|}$$

$$= \frac{8}{x^2-l^2+n^2-1} + \frac{1^2-n^2}{1} + \frac{1^2-l^2}{x^2+1} + \frac{3^2-n^2}{1} + \frac{3^2-l^2}{x^2-1} + \dots$$

$$Q. \frac{P}{Q} = \frac{\left| \frac{\alpha+\beta+\gamma+\delta+\epsilon-1}{2} \right| \left| \frac{\alpha+\beta+\gamma-\delta-\epsilon-1}{2} \right|}{\left| \frac{\alpha+\beta-\gamma-\delta+\epsilon-1}{2} \right| \left| \frac{\alpha-\beta-\gamma+\delta+\epsilon-1}{2} \right| \left| \frac{\alpha-\beta+\gamma+\delta-\epsilon-1}{2} \right|} \times$$

$$\frac{\left| \frac{\alpha-\beta+\gamma-\delta+\epsilon-1}{2} \right| \left| \frac{\alpha+\beta-\gamma+\delta-\epsilon-1}{2} \right| \left| \frac{\alpha-\beta-\gamma-\delta-\epsilon-1}{2} \right|}{\left| \frac{\alpha+\beta+\gamma+\delta-\epsilon-1}{2} \right| \left| \frac{\alpha+\beta+\gamma-\delta+\epsilon-1}{2} \right|} \times$$

$$\text{and } Q = \frac{\left| \frac{\alpha+\beta+\gamma+\delta-\epsilon-1}{2} \right| \left| \frac{\alpha+\beta+\gamma-\delta+\epsilon-1}{2} \right|}{\left| \frac{\alpha+\beta-\gamma+\delta+\epsilon-1}{2} \right| \left| \frac{\alpha-\beta-\gamma+\delta+\epsilon-1}{2} \right| \left| \frac{\alpha+\beta-\gamma-\delta-\epsilon-1}{2} \right|} \times$$

$$\frac{\left| \frac{\alpha-\beta+\gamma-\delta-\epsilon-1}{2} \right| \left| \frac{\alpha-\beta-\gamma+\delta-\epsilon-1}{2} \right| \left| \frac{\alpha-\beta-\gamma-\delta+\epsilon-1}{2} \right|}{\left| \frac{\alpha+\beta+\gamma+\delta-\epsilon-1}{2} \right| \left| \frac{\alpha+\beta+\gamma-\delta+\epsilon-1}{2} \right|} \text{, then}$$

$$\frac{P \cdot Q}{P+Q} = \frac{8\alpha\beta\gamma\delta\epsilon}{\left\{ 2(\alpha^4+\beta^4+\gamma^4+\delta^4+\epsilon^4+1) - (\alpha^2+\beta^2+\gamma^2+\delta^2+\epsilon^2-1)^2 - 2^2 \right\} + 3\left\{ 2(\alpha^4+\beta^4+\gamma^4+\delta^4+\epsilon^4+1) - (\alpha^2+\beta^2+\gamma^2+\delta^2+\epsilon^2-5)^2 - 6^2 \right\} + 64(\alpha^2-1)(\beta^2-1)(\gamma^2-1)(\delta^2-1)(\epsilon^2-1)}$$

$$64(d^2-2^2)(\beta^2-2^2)(\gamma^2-2^2)(\delta^2-2^2)(\epsilon^2-2^2)$$

$$5\{2(d^2+\beta^2+\gamma^2+\delta^2+\epsilon^2+1) - (d^2+\beta^2+\gamma^2+\delta^2+\epsilon^2-9)^2 - 10^2\} + \&c$$

N.B. If any one of  $d, \beta, \gamma, \delta, \epsilon$  be an integer the theorem is true. The result will be permanently true if  $d$  is removed from the numerators or if it is expanded in powers of  $\frac{1}{2}$ .

$$41. 1 + \frac{\beta}{\gamma+1}x + \frac{\beta(\beta-1)}{(\gamma+1)(\gamma+2)}x^2 + \&c = \frac{\sqrt{\beta}\sqrt{\gamma}}{\beta+\gamma} \cdot \frac{(1+x)^{\beta+\gamma}}{x^\gamma}$$
  
$$\frac{\gamma}{(\beta+1)x+1-\gamma} - \frac{1(1-\gamma)(1+x)}{(\beta+2)x+3-\gamma} - \frac{2(2-\gamma)(1+x)}{(\beta+3)x+5-\gamma} - \&c.$$

$$42. 1 + \frac{x}{n+1} + \frac{x^2}{(n+1)(n+2)} + \frac{x^3}{(n+1)(n+2)(n+3)} + \&c$$
  
$$= \frac{e^x \Gamma(n)}{x^n} - \frac{x}{x+1} - \frac{1-n}{1+x} - \frac{1}{x+2} - \frac{2-n}{1+x} - \frac{2}{x+3} - \frac{2-n}{1+x} - \&c.$$
  
$$= \frac{e^x \Gamma(n)}{x^n} - \frac{n}{x+1-n} - \frac{1(1-n)}{x+3-n} - \frac{2(2-n)}{x+5-n} - \frac{3(3-n)}{x+7-n} - \&c.$$

$$\text{Cor. } \frac{1}{n} - \frac{x}{2} \cdot \frac{1}{n+1} + \frac{x^2}{12} \cdot \frac{1}{n+2} - \frac{x^3}{12} \cdot \frac{1}{n+3} + \&c$$
  
$$= \frac{\Gamma(n)}{x^n} - \frac{e^{-x}}{x+1} - \frac{1-n}{1+x} - \frac{1}{x+2} - \frac{2-n}{1+x} - \frac{2}{x+3} - \&c$$

$$43. 1 + \frac{x}{1.3} + \frac{x^2}{1.3.5} + \frac{x^3}{1.3.5.7} + \frac{x^4}{1.3.5.7.9} + \&c$$
  
$$= \sqrt{\frac{\pi}{2x}} e^x - \frac{1}{x+1} - \frac{1}{1+x} - \frac{2}{2+x} - \frac{3}{1+x} - \frac{4}{x+3} - \frac{5}{1+x} - \&c$$
  
$$= \sqrt{\frac{\pi}{2x}} e^x - \frac{1}{x+1} - \frac{1.2}{x+5} - \frac{3.4}{x+7} - \frac{5.6}{x+9} - \&c$$

$$\text{Cor. } 1. \int_0^x e^{-x^2} dx = \frac{\sqrt{\pi}}{2} - \frac{e^{-x^2}}{2x} - \frac{1}{x} - \frac{2}{2x} - \frac{3}{x} - \frac{4}{2x} - \&c$$

$$2. \int_0^x \int_0^x \frac{e^{-x^2}}{x} dx dx = \frac{\sqrt{\pi}}{2} (\frac{1}{2} + \log 2x) \text{ when } x \text{ is very great.}$$

$$44. \int_0^x \frac{1-e^{-x}}{x} dx = \frac{x}{12} - \frac{x^2}{24} + \frac{x^3}{72} - \&c = C + \log x + e^{-x} \phi(x)$$

$$i. \phi(x) = \frac{1}{x} - \frac{1}{x^2} + \frac{1}{2x^3} - \frac{1}{24x^4} + \&c$$

ii.  $\phi(x)$  lies between  $\frac{1}{x}$  &  $\frac{1}{x+1}$  and very nearly equals  $\sqrt{\frac{\phi(x+1)}{x}}$

iii.  $\phi(x) = \frac{1}{x+1} - \frac{1}{x+2} + \frac{1}{x+3} - \frac{1}{x+4} + \frac{1}{x+5} - \frac{1}{x+6} + \dots$   
 $= \frac{1}{x+1} - \frac{1^2}{x+3} + \frac{1^2}{x+5} - \frac{1^2}{x+7} + \dots$

iv.  $\phi(x) = \frac{1}{x} - \frac{1!}{x^2} + \frac{1!}{x^3} - \frac{2!}{x^4} + \dots \pm \frac{1!}{x^n} \frac{1}{x+n+1} - \frac{1!(1+n)}{x+n+3} - \dots$   
 $= \frac{2(2+n)}{x+n+5} - \frac{3(3+n)}{x+n+7} - \dots$

Ex. 1.  $\frac{x}{1!} + \frac{x^2}{2!}(1+\frac{1}{x}) + \frac{x^3}{3!}(1+\frac{1}{x}+\frac{1}{x^2}) + \dots = e^x(c_0 + \log x) + \phi(x)$ .

Ex. 2. If  $\int_0^{n(1-h)} \frac{1-e^{-x}}{x} dx = C + \log n$ , then

$$\phi(n) = h(e^n - 1) + \frac{h^2}{2}(e^n - 1 - \frac{n}{h}) + \frac{h^3}{6}(e^n - 1 - \frac{n}{h} - \frac{n^2}{2h^2}) + \dots$$

$\phi(1) = .5763474$ ;  $\phi(10) = .9229106$ .

45. i. Den<sup>n</sup> in  $\frac{1}{1+x} + \frac{x}{1+x^2} + \frac{x^2}{1+x^3} + \frac{x^3}{1+x^4} + \dots = \frac{(n-1)x}{1+x} + \frac{nx}{1+x^2}$   
 $= 1 + \frac{n^2}{2}x + \frac{n^2(n-1)^2}{12}x^2 + \frac{n^2(n-1)^2(n-2)^2}{24}x^3 + \dots$

ii. Den<sup>n</sup> in  $\frac{1}{1+x} + \frac{x}{1+x^2} + \frac{x^2}{1+x^3} + \frac{x^3}{1+x^4} + \dots + \frac{(n-1)x}{1+x} + \frac{(n-1)x}{1+x^2}$   
 $= 1 + \frac{n^2}{2}(1-\frac{1}{n})x + \frac{n^2(n-1)^2}{12}(1-\frac{2}{n})x^2 + \frac{n^2(n-1)^2(n-2)^2}{24}(1-\frac{3}{n})x^3 + \dots$

46. i.  $\frac{x}{1!} - \frac{x^2}{2^2!} + \frac{x^3}{3^2!} - \dots = \phi_n(x) + (-1)^{n-1} \psi_n(x) e^{-x}$ .

where  $\phi_n(x)$  is the term independent of  $p$  in  $\frac{x^p 1-p}{p^n}$  and  $\psi_n(x) - \psi_n'(x) = \frac{\psi_{n-1}(x)}{x}$ .

ii.  $\phi_n(x) = \frac{1}{12} \{ A_0 (\log x)^n + \frac{n}{2} A_1 (\log x)^{n-1} + \frac{n(n-1)}{2} A_2 (\log x)^{n-2} + \dots + A_n \}$  where  $12 = A_0 - A_1 \frac{x}{2} + A_2 \frac{x^2}{12} - A_3 \frac{x^3}{24} + \dots$

$A_n = S_1 A_{n-1} + (n-1) S_2 A_{n-2} + (n-1)(n-2) S_3 A_{n-3} + \dots$

iii.  $12 = 1 - .5772156649x + .9890560173x^2 - \dots$

$$.9074790803x^3 + .9817280965 \frac{x^4}{1+\theta x}$$

$$\theta_0 = 1.00027; \theta_1 = \frac{51}{52}; \theta_2 = \frac{77}{82}; \theta_3 = \frac{5}{28}; \theta_4 = -\frac{1}{38} \text{ nearly}$$

$$i.v. \Psi_n(x) = \frac{x}{\left(x + \frac{x}{2} + \frac{5x+10}{6x} + \frac{41x+58}{10+2x}\right)^{n+1}}$$

$$ex. \int \frac{1-e^{-x}}{x} dx - \frac{1}{2} \left\{ \int \frac{1-e^{-x}}{x} dx \right\}^2 = \frac{\pi^2}{12} \text{ when } x \text{ is great.}$$

$$47. \int_0^{\infty} e^{-x} \left(1 + \frac{x}{n}\right)^n dx = 1 + \frac{x}{1} + \frac{1(n-1)}{3} + \frac{2(n-2)}{5} + \frac{3(n-3)}{7} + \dots$$

$$= 2 + \frac{n-1}{2} + \frac{1(n-2)}{4} + \frac{2(n-3)}{6} + \frac{3(n-4)}{8} + \dots$$

$$= \frac{e^n \ln n}{2n^n} - \frac{2n}{2} + \frac{3n}{3} - \frac{4n}{4} + \frac{5n}{5} + \dots$$

$$48. \int_0^{\infty} e^{-x} \left(1 + \frac{x}{n}\right)^n dx = \frac{e^n \ln n}{2n^n} + \frac{2}{3} - \frac{4}{135n} + \frac{8}{27 \cdot 105 n^2}$$

$$+ \frac{16}{105 \cdot 81 n^3} - \frac{32281}{3^8 \cdot 5^2 \cdot 7 \cdot 11 n^4} - \dots$$

$$Corr. 1 + \frac{x}{2} + \frac{x^2}{6} + \dots + \frac{x^n}{n!} \theta = \frac{e^x}{2}$$

$$\text{where } \theta = \frac{4+15n}{8+45n} \text{ very nearly.}$$

N.B.	$x=0$	Real value of $\theta$	App. value of $\theta$
		.50000	.50000
	$x = \frac{1}{3}$	.37750	.37705
	$x = 1$	.35914	.35849
	$x = 1\frac{1}{2}$	.35146	.35099
	$x = 2$	.34726	.34694
	$x = \infty$	.33333	.33333

$$49. C_0 + \log n + \frac{x}{12} + \frac{x^2}{24} + \frac{x^3}{36} + \frac{x^4}{48} + \dots$$

$$= e^n \left( \frac{1}{n} + \frac{1}{2n^2} + \frac{1}{3n^3} + \dots + \frac{1(n-1)}{n^n} \theta \right)$$

$$\text{where } \theta = \frac{2}{3} + \frac{4}{135n} + \frac{8}{27 \cdot 105 n^2} - \dots$$

1. If  $n$  is the integer just greater than  $n$  or equal to  $n$ ,

$$\int_0^{\infty} \frac{A_1 x + A_2 x^2 + A_3 x^3 + \dots}{x^{n+1}} dx = \cos \pi N \int_0^{\infty} \frac{A_N x^N + A_{N+1} x^{N+1} + \dots}{x^{n+1}} dx$$

e.g.  $\int_0^{\infty} \frac{e^{-x^2}}{x^4} dx = \frac{2}{3} \sqrt{\pi}$  really means that

$$\int_0^{\infty} \frac{e^{-x^2} - 1 + x^2}{x^4} dx = \frac{2}{3} \sqrt{\pi}$$

Cor. Thus the meanings of the integrals  $\int_0^{\infty} e^{-ax} x^{n-1} \frac{\cos bx}{\sin bx} dx$

$$= \frac{\Gamma(n)}{(a^2 + b^2)^{\frac{n}{2}}} \frac{\cos(n \tan^{-1} \frac{b}{a})}{\sin(n \tan^{-1} \frac{b}{a})} \text{ for negative values of } n \text{ are known.}$$

$$2i. \int \phi(x) e^{-nx} dx = -e^{-nx} \left\{ \frac{\phi(x)}{n} + \frac{\phi'(x)}{n^2} + \frac{\phi''(x)}{n^3} + \dots \right\}$$

$$ii. \int \phi(x) \cos nx dx = \sin nx \left\{ \frac{\phi(x)}{n} - \frac{\phi''(x)}{n^3} + \dots \right\} \\ + \cos nx \left\{ \frac{\phi'(x)}{n^2} - \frac{\phi'''(x)}{n^4} + \dots \right\}$$

$$iii. \int \phi(x) \sin nx dx = \sin nx \left\{ \frac{\phi'(x)}{n^2} - \frac{\phi'''(x)}{n^4} + \dots \right\} \\ - \cos nx \left\{ \frac{\phi(x)}{n} - \frac{\phi''(x)}{n^3} + \dots \right\}$$

$$3. \int_x^{\infty} e^{-x^2} \cos 2nx dx = e^{-x^2} \left\{ \frac{\cos(2nx + \theta)}{2n} - \frac{1 \cos(2nx + 2\theta)}{2^2 n^2} \right. \\ \left. + \frac{1.3 \cos(2nx + 5\theta)}{2^3 n^3} - \frac{1.3.5 \cos(2nx + 7\theta)}{2^4 n^4} + \dots \right\}$$

where  $\tan \theta = \frac{n}{x}$  and  $n = \sqrt{x^2 + x^2}$ .

$$4. \int_0^{\infty} e^{-x^2} \left\{ e^{2nx} \phi(x) + e^{-2nx} \phi(-x) \right\} dx$$

$$= \int_0^{\infty} e^{n^2 - x^2} \left\{ \phi(n+x) + \phi(n-x) \right\} dx =$$

$$\sqrt{\pi} e^{x^2} \left\{ \phi(x) + \frac{\phi'(x)}{4} + \frac{\phi''(x)}{4 \cdot 8} + \frac{\phi'''(x)}{4 \cdot 8 \cdot 12} + \dots \right\}$$

$$5. \int_0^{\infty} e^{-\frac{x^2}{2}} \left\{ A_1 - \frac{x^2}{12} A_3 + \frac{x^4}{12} A_5 - \dots \right\} dx$$

$$= \frac{\sqrt{\pi}}{2} \left\{ A_0 - \frac{2}{12} A_2 + \frac{2^2}{12} A_4 - \frac{2^3}{12} A_6 + \dots \right\}$$

$$6. \int_0^{\infty} e^{-x} \left(1 + \frac{x}{n}\right)^{m-h} dx = 1 + \left(1 - \frac{h}{n}\right) + \left(1 - \frac{h}{n}\right)\left(1 - \frac{h+1}{n}\right) + \left(1 - \frac{h}{n}\right)\left(1 - \frac{h+1}{n}\right)\left(1 - \frac{h+2}{n}\right) + \dots$$

$$= \frac{e^n \Gamma(m-h)}{2 n^{m-h}} + A_0 - \frac{A_1}{n} + \frac{A_2}{n^2} - \dots \text{ where}$$

$$A_0 = \frac{2}{3} - h; \quad A_1 = \frac{4}{135} - \frac{h^2(h-h)}{3};$$

$$A_2 = \frac{8}{2835} + \frac{2h(1-h)}{135} - \frac{h(1-h^2)(2-2h^2)}{45} \quad \&c.$$

$$7. (m-n-1) \int_0^{\infty} \frac{\left(1 + \frac{x}{n}\right)^m}{\left(1 + \frac{x}{m}\right)^m} dx = \frac{m}{2} \cdot \frac{m \Gamma(m)}{n^m \Gamma(m)} \cdot \frac{\Gamma(m-n)}{(m-n)^{m-n}}$$

$$+ \frac{2}{3} (m+n) - \frac{4(m+n)(m-2n)(m-\frac{n}{2})}{135 m n (m-n)}$$

$$+ \frac{8(m^3+n^3)(m-2n)(m-\frac{n}{2})}{2835 m^2 n^2 (m-n)^2}$$

$$+ \frac{16(m^3+n^3)(m-2n)(m-\frac{n}{2})(m^2-mn+n^2)}{8505 m^3 n^3 (m-n)^3} - \dots \&c.$$

$$8. \int_0^{\infty} \left\{ \frac{x^m \Gamma(m)}{\Gamma(m+x)} + e^{-x} \left(1 + \frac{x}{n}\right)^m \right\} dx = \frac{e^n \Gamma(m)}{n^m} + \frac{6n}{12n+1}$$

very very nearly.

9. Let  $\int_0^{\infty} \frac{e^{-m^2 x^2}}{1+x^2} dx = \phi(m)$  and if  $m \ll m$ , then

$$\int_0^{\infty} \frac{e^{-mx} x^2}{1+x^2} \cos 2mnx dx = \frac{e^{-n^2}}{2} \{ \phi(2n+i) + \phi(2n-i) \}$$

10. 
$$= \frac{\phi(h \cdot \alpha + \delta)}{\phi(h \cdot \beta + \gamma)} + \frac{\phi(h \cdot \alpha + \delta)}{\phi(h \cdot \beta + \gamma)} \cdot \frac{\phi(h \cdot \alpha + 2\delta)}{\phi(h \cdot \beta + 2\gamma)} + \dots$$

$$= \sqrt{\frac{\pi \phi(0)}{2h(\gamma - \delta) \phi'(0)}} + \frac{1}{3} \cdot \frac{\gamma + \delta}{\gamma - \delta} \left\{ 1 - \frac{\phi(0) \cdot \phi''(0)}{\phi'(0)^2} \right\} + \frac{\alpha - \beta}{\gamma - \delta}$$

if  $h$  is very small.

Cor. i. 
$$1 + \left(\frac{x}{x+1}\right)^n + \left\{ \frac{x^2}{(x+1)(x+2)} \right\}^n + \left\{ \frac{x^3}{(x+1)(x+2)(x+3)} \right\}^n + \dots$$

$$= \sqrt{\frac{\pi x}{2n}} + \frac{1}{3n}$$

when  $x$  is very great

ii. 
$$1 + \left(\frac{x}{2}\right)^n + \left(\frac{x^2}{4}\right)^n + \left(\frac{x^3}{8}\right)^n + \dots$$

$$= \frac{e^{nx} + \frac{x^2-1}{2^2} \left( \frac{1}{n!x} + \frac{1}{2n!x^2} + \dots \right)}{\sqrt{n} \cdot (2\pi x)^{\frac{n-1}{2}}}$$

11. i. 
$$1 + \left(\frac{e^n}{1}\right) + \left(\frac{e^{2n}}{2}\right)^2 + \left(\frac{e^{3n}}{3}\right)^3 + \left(\frac{e^{4n}}{4}\right)^4 + \dots$$

$$= \sqrt{2\pi n} e^{2n} - \frac{1}{24n} - \frac{1}{48n^2} - \left(\frac{1}{36} + \frac{1}{5760}\right) \frac{1}{n^3} - \dots$$

if  $n$  is great

ii. 
$$\int_0^{\infty} \frac{x^{n-1} dx}{1 + \frac{x}{2} + \left(\frac{x}{2}\right)^2 + \left(\frac{x}{2}\right)^3 + \left(\frac{x}{2}\right)^4 + \dots}$$

$$= n^n \left( \frac{1}{n} + \frac{1}{2n^2} + \frac{1}{3n^3} + \frac{2}{3n^4} + \dots \right)$$

if  $n$  is great.

iii. 
$$\log 2 \left( \frac{1}{2 \log 2} - \frac{1}{3 \log 3} + \frac{1}{4 \log 4} - \frac{1}{5 \log 5} + \dots \right)$$

$$+ (\log 2)^2 \left( \frac{1}{2 \log 2 \log 4} + \frac{1}{3 \log 3 \log 6} + \frac{1}{4 \log 4 \log 8} + \dots \right) = 1$$

12. The approximate value of  $e^{-x} \left\{ \phi(0) + \frac{x}{1!} \phi(1) + \frac{x^2}{2!} \phi(2) + \dots \right\}$  when  $x$  is great can be found by successive differentiation, and by transforming the result applying III & ex. 1. if necessary.



157,  
 e.g.  $\log 1 + \frac{x}{11} \log 2 + \frac{x^2}{11} \log 3 + \frac{x^3}{11} \log 4 + \dots$

$$= e^x \left( \log x + \frac{1}{2x} + \frac{1}{12x^2} + \frac{1}{12x^3} + \frac{19}{120x^4} + \frac{9}{20}x^5 + \dots \right)$$

13.  $\int_0^{\infty} \frac{dx}{(x^2+a^2)(x^2+b^2)(x^2+c^2)(x^2+d^2)}$

$$= \frac{\pi}{4} \frac{(a+b+c+d)^3 - (a^3+b^3+c^3+d^3)}{abcd(a+b)(b+c)(c+a)(a+d)(b+d)(c+d)}$$

Cor. If  $\alpha, \beta, \gamma, \delta$  are the roots of  $x^4 - px^3 + qx^2 - rx + s = 0$

then  $\int_0^{\infty} \frac{dx}{(x^2+\alpha^2)(x^2+\beta^2)(x^2+\gamma^2)(x^2+\delta^2)} = \frac{\pi}{2\beta} \cdot \frac{1}{\alpha} - \frac{p\delta}{q - \frac{r}{\beta}}$

14.  $\frac{1}{a + \frac{x^2}{a}} - \frac{2a}{11} \cdot \frac{1}{a+1 + \frac{x^2}{a+1}} + \frac{2a(2a+1)}{12} \cdot \frac{1}{a+2 + \frac{x^2}{a+2}} - \dots$

$$= \frac{\Delta (1a-1)^2 / 2(2a-1)}{\left\{1 + \left(\frac{x}{a}\right)^2\right\} \left\{1 + \left(\frac{x}{a+1}\right)^2\right\} \left\{1 + \left(\frac{x}{a+2}\right)^2\right\} \left\{1 + \left(\frac{x}{a+3}\right)^2\right\} \dots}$$

and

$$\int_0^{\infty} \frac{\cos nx}{a + \frac{x^2}{a}} dx = \frac{\pi}{2} e^{-na}; \text{ Combining these results}$$

15.  $\int_0^{\infty} \frac{\cos 2nx dx}{\left\{1 + \left(\frac{x}{a}\right)^2\right\} \left\{1 + \left(\frac{x}{a+1}\right)^2\right\} \left\{1 + \left(\frac{x}{a+2}\right)^2\right\} \left\{1 + \left(\frac{x}{a+3}\right)^2\right\} \dots} = \frac{\sqrt{\pi}}{2} \cdot \frac{1a-1}{|a-1|} \text{Sech}^2$

16. i  $\int_0^{\infty} \frac{\sinh ax}{\sinh \pi x} \cos nx dx = \frac{1}{2} \cdot \frac{\sin a}{\cosh n + \cos a}$

ii.  $\int_0^{\infty} \frac{\cosh ax}{\sinh \pi x} \sin nx dx = \frac{1}{2} \cdot \frac{\sinh n}{\cosh n + \cos a}$

iii.  $\int_0^{\infty} \frac{\sin^2 nx}{e^{2\pi x} - 1} dx = \frac{1}{2} \left( \frac{1}{e^{2n}} + \frac{1}{2} - \frac{1}{2n} \right);$

iv.  $\int_0^{\infty} \frac{x^{n-1}}{e^{2\pi x} - 1} dx = \frac{B_n}{2n} \cdot \int_0^{\infty} \frac{x^{n-1}}{\cosh \frac{\pi x}{2}} dx = F_n$

17.  $\phi(0) + \phi(l) + \phi(x) + \dots + \phi(n)$   
 $\int_0^l \phi(x) dx + \frac{1}{2} \phi(n) + \int_0^\infty \frac{\phi(n+x) - \phi(n-x)}{i(e^{2\pi x} - 1)} dx$

Con.  $\log n = n \log n - n + \frac{1}{2} \log(2\pi n) + 2 \int_0^\infty \frac{\tan^{-1} \frac{x}{n}}{e^{2\pi x} - 1} dx$

18. i. If  $f(x) + \phi(x) = f(x+l)$ , then

$$f(x) + \frac{1}{2} \phi(x) = \frac{1}{2} \int_0^x \phi(\xi) d\xi + 2 \int_0^\infty \frac{\phi(x+i\xi) - \phi(x-i\xi)}{(e^{2\pi \xi} - 1)i} d\xi$$

ii. If  $f(x+l) + f(x-l) = \phi(x)$ , then

$$2f(x) = \int_0^\infty \frac{\phi(x+i\xi) + \phi(x-i\xi)}{e^{\pi \xi} + e^{-\pi \xi}} d\xi$$

19. i. If  $\int_0^h \phi(x) \cos nx dx = \psi(n)$   $m < \leq h$

then  $\int_0^\infty \psi(x) \cos mx dx = \frac{\pi}{2} \phi(m), \frac{\pi}{4} \phi(m), 0$

ii. If  $\int_0^h \phi(x) \sin nx dx = \psi(n)$

then  $\int_0^\infty \psi(x) \sin mx dx = \frac{\pi}{2} \phi(m), \frac{\pi}{4} \phi(m), 0$   $m < \leq h$

Con.  $\int_0^\infty \operatorname{sech}^{2a} x \cos 2\pi x dx$   
 $= \frac{\sqrt{\pi} |a-1/2|}{\{1+(\frac{m}{a})^2\} \{1+(\frac{m}{a+1})^2\} \{1+(\frac{m}{a+2})^2\} \&c}$

20.  $\int_0^\infty \frac{\sinh ax}{\sinh \pi x} \cdot \frac{dx}{1+n^2x^2}$  (a lying between 0 and  $\pi$ )

$$= \frac{\sin a}{1+n} - \frac{\sin 2a}{1+2n} + \frac{\sin 3a}{1+3n} - \frac{\sin 4a}{1+4n} + \&c$$

21.  $\int_{\alpha_1}^{\beta_1} \phi_1(p, x) F(x) dx = \psi_1(p, n)$

&  $\int_{\alpha_2}^{\beta_2} \phi_2(p, x) F(x) dx = \psi_2(p, n)$ , then

$$\int_{\alpha_1}^{\beta_1} \phi_1(p, x) \psi_2(q, n(x)) dx = \int_{\alpha_2}^{\beta_2} \phi_2(q, x) \psi_1(p, n(x)) dx$$

Cor.  $\int_0^{\infty} \phi(p, x) \cos nx dx = \psi(p, n)$ , then

$$\frac{\pi}{2} \int_0^{\infty} \phi(p, x) \phi(q, bx) dx = \int_0^{\infty} \psi(p, x) \psi(q, bx) dx$$

ex.  $\int_0^{\infty} dx = \pi$ , then  $\sqrt{a} \int_0^{\infty} \frac{e^{-x^2}}{e^{ax} + e^{-ax}} dx = \sqrt{b} \int_0^{\infty} \frac{e^{-x^2}}{e^{bx} + e^{-bx}} dx$

N.B. This can also be got from the theorem: - if  $d\beta = \frac{2}{\beta}$

$$\sqrt{a} \{ E_1 - E_3 \frac{a^2}{a} + E_5 \frac{a^4}{a} - \dots \} = \sqrt{b} \{ E_1 - E_3 \frac{b^2}{b} + E_5 \frac{b^4}{b} - \dots \}$$

which is obtained from the theorem: -

$$\phi(1) - \phi(2) + \phi(3) - \dots = \phi(1) - \phi(1) + \phi(2) - \dots$$

22.i.  $\int_0^{\infty} \frac{1}{\sqrt{1+(\frac{x}{a})^2}} \sqrt{1+(\frac{x}{a+1})^2} \sqrt{1+(\frac{x}{a+2})^2} \dots \sqrt{1+(\frac{x}{a+n})^2} dx$

$$= \frac{\sqrt{\pi}}{2} \cdot \frac{|a-\frac{1}{2}| |a-\frac{1}{2}| |a+b-1|}{|a-1| |b-1| |a+b-\frac{1}{2}|}$$

ii.  $\int_0^{\infty} \frac{1+(\frac{x}{b+2})^2}{1+(\frac{x}{a})^2} \frac{1+(\frac{x}{b+2})^2}{1+(\frac{x}{a+1})^2} \frac{1+(\frac{x}{b+2})^2}{1+(\frac{x}{a+2})^2} \dots dx$

$$= \frac{\sqrt{\pi}}{2} \cdot \frac{|a-\frac{1}{2}| |b-\frac{1}{2}| |b-a-\frac{1}{2}|}{|a-1| |b-\frac{1}{2}| |b-\frac{1}{2}|}$$

$$23. \int_0^{\infty} \frac{x^{l+a-1}}{(x+a)^n} \cdot \frac{dx}{x^m}$$

$$= \frac{\pi}{\Gamma(n)} \operatorname{Cosec} \pi m \left\{ \frac{1}{a^m} - \frac{n}{\Gamma} \cdot \frac{1}{(a+1)^m} + \frac{n(n-1)}{\Gamma^2} \cdot \frac{1}{(a+2)^m} - \dots \right\}$$

24. i.  $A_0 + A_1 + A_2 + \dots + A_n$   
 $= A_n + A_{n-1} + \dots + A_0$  &c to infinity  $- (A_{-1} + A_{-2} + A_{-3} + \dots)$

Cor.  $A_0 + \frac{A_1}{\Gamma} + \frac{A_2}{\Gamma^2} + \dots + \frac{A_n}{\Gamma^n}$   
 $= \frac{A_n}{\Gamma^n} + \frac{A_{n-1}}{\Gamma^{n-1}} + \dots$  ad. inf.

ii.  $\phi(x) + \{\phi(x+1) + \phi(x-1)\} + \{\phi(x+2) + \phi(x-2)\} + \dots$   
 $= \phi(y) + \{\phi(y+1) + \phi(y-1)\} + \{\phi(y+2) + \phi(y-2)\} + \dots$

Cor.  $\frac{x^h}{\Gamma} + \left( \frac{x^{h+n}}{\Gamma^{h+n}} + \frac{x^{h-n}}{\Gamma^{h-n}} \right) + \left( \frac{x^{h+2n}}{\Gamma^{h+2n}} + \frac{x^{h-2n}}{\Gamma^{h-2n}} \right) + \dots$   
 $= 1 + \left( \frac{x^n}{\Gamma^n} + \frac{x^{-n}}{\Gamma^{-n}} \right) + \left( \frac{x^{2n}}{\Gamma^{2n}} + \frac{x^{-2n}}{\Gamma^{-2n}} \right) + \dots = \frac{e^x}{\Gamma^n}$

for all values of  $x, n$  and  $h, n$  being  $\geq 1$ .

iii.  $\int_{-\infty}^{\infty} \frac{\phi(x)}{\Gamma x} dx = \phi(0) + \frac{\phi(1)}{\Gamma} + \frac{\phi(2)}{\Gamma^2} + \frac{\phi(3)}{\Gamma^3} + \dots$

Cor. 1.  $\int_{-\infty}^{\infty} \frac{a^x}{\Gamma x} dx = e^a$ , Cor. 2.  $\int_{-\infty}^{\infty} \frac{a^x \Gamma^n}{\Gamma x \Gamma^n x} dx = (1+a)^n$

Let  $n < \frac{1}{2}\pi$  in the following examples

25. i.  $\int_0^{\infty} \left( \frac{a^x}{\Gamma x} + \frac{a^{-x}}{\Gamma -x} \right) \operatorname{Cos} nx dx = e^{a \operatorname{Cos} n} \operatorname{Cos}(a \operatorname{Sin} n)$

&  $\int_0^{\infty} \left( \frac{a^x}{\Gamma x} - \frac{a^{-x}}{\Gamma -x} \right) \operatorname{Sin} nx dx = e^{a \operatorname{Cos} n} \operatorname{Sin}(a \operatorname{Sin} n)$

ii.  $\int_0^{\infty} \left( \frac{a^{b+x}}{\Gamma b+x} + \frac{a^{b-x}}{\Gamma b-x} \right) \operatorname{Cos} nx dx = e^{a \operatorname{Cos} n} \operatorname{Cos}(a \operatorname{Sin} n - \dots)$

$$\& \int_0^{\infty} \left( \frac{a^{b+x}}{b+x} - \frac{a^{b-x}}{b-x} \right) \sin nx \, dx = e^{a \cos n} \sin(a \sin n - n).$$

N.B. i. The maximum value of  $\frac{a^x}{b-x} = \frac{e^{\int \frac{x}{a} da}}{\sqrt{2\pi}}$

$$= \frac{a^{a-\frac{1}{2}}}{a-\frac{1}{2}} e^{\frac{1}{2} a^2 (3.6 a^2 + 10 \cdot 1)}$$

very nearly.

ii. The following theorem is very useful in evaluating definite integrals:—

$$\int_a^b \phi(x) dx = h \left\{ \frac{1}{2} \phi(a) + \phi(a+h) + \phi(a+2h) + \phi(a+3h) + \dots + \phi(b-2h) + \phi(b-h) + \frac{1}{2} \phi(b) \right\}$$

$$+ B \frac{h^2}{12} \{ \phi'(a) - \phi'(b) \} - B \frac{h^4}{45} \{ \phi'''(a) - \phi'''(b) \} + \&c.$$

26. i.  $\int_0^{\infty} \frac{\cos 2m x}{(1+x^2)^{m+1}} dx = \frac{\pi}{2} \cdot \frac{\pi^m}{\Gamma(m)} e^{-2m} \left\{ 1 + \frac{m}{4} \cdot \frac{m+1}{\pi} + \frac{m(m-1)}{4 \cdot 8} \cdot \frac{(m+1)(m+2)}{\pi^2} + \frac{m(m-1)(m-2)}{4 \cdot 8 \cdot 12} \cdot \frac{(m+1)(m+2)(m+3)}{\pi^3} + \dots \right\}$

ii  $\int_0^{\infty} \frac{x^{2m}}{(1+x^2)^{n+1}} \cos px \, dx = \frac{\pi}{2} \cdot (-1)^m \cdot \frac{e^{-p}}{2^n \Gamma(n)} \left\{ p^n + A_1 p^{n-1} + A_2 p^{n-2} + \dots \right\}$

where  $m$  is any positive integer and  $A_n = \frac{n+2}{\Gamma(n-2)} \cdot \frac{1}{2^n} \cdot \frac{1}{a}$

$$\left\{ 1 - \frac{4}{4} \cdot \frac{n m n}{(n+2)(n+1)} + \frac{4^2}{12} \cdot \frac{n(n-1)n(n-1)n(n-1)}{(n+2)(n+1)(n+2)(n+1)(n+2)} - \&c \right\}$$

27.  $\left\{ 1 + \left(\frac{1}{2}\right)^n \right\} \left\{ 1 + \left(\frac{1}{3}\right)^n \right\} \left\{ 1 + \left(\frac{1}{4}\right)^n \right\} \&c$   $n$  being even

$$= \prod \sqrt{\frac{\cosh(2\pi x \sin \frac{\pi a}{n}) - \cos(2\pi x \cos \frac{\pi a}{n})}{2\pi^2 x^2}} \quad \text{where}$$

$$a = 1, 3, 5 \text{ to } n-1.$$

Cor.  $\left\{ 1 + \left(\frac{2n}{n+1}\right)^3 \right\} \left\{ 1 + \left(\frac{2n}{n+2}\right)^3 \right\} \left\{ 1 + \left(\frac{2n}{n+3}\right)^3 \right\} \&c$

$$= \frac{(2n)^3}{8n} \cdot \frac{\sinh(\pi n \sqrt{3})}{\pi n \sqrt{3}}$$

N.B Thus it is possible to find the value of the product:

$$\{1 + (\frac{x}{a})^2\} \{1 + (\frac{x}{a+d})^2\} \{1 + (\frac{x}{a+2d})^2\} \&c$$

Cor. 2.  $\{1 + (\frac{2m+1}{n+1})^2\} \{1 + (\frac{2m+3}{n+2})^2\} \{1 + (\frac{2m+5}{n+3})^2\}$   
 $= \frac{(2m+1)^3}{16m+2} \frac{13m+14}{(2m+2)^3} \text{Cosh} \{ \pi(m+t) \sqrt{3} \} \cdot \frac{(2m)^3}{\pi(2m+1)}$

28.  $m \pi \{ 1 + (\frac{x^n}{1^n} + \frac{x^{-n}}{1^{-n}}) + (\frac{x^{2n}}{1^{2n}} + \frac{x^{-2n}}{1^{-2n}}) + \&c \}$   
 $= e^x + e^{x \cos \frac{2\pi}{n}} \text{Cos}(x \sin \frac{2\pi}{n}) + e^{x \cos \frac{4\pi}{n}} \text{Cos}(x \sin \frac{4\pi}{n})$   
 $+ e^{x \cos \frac{6\pi}{n}} \text{Cos}(x \sin \frac{6\pi}{n}) + \&c.$  to  $m \pi$  terms where  $m$  is any arbitrary integer.

29.  $i \int_0^\infty \frac{(-x^2)^l}{1+x^{2n}} \text{Cos } px \, dx = \frac{\pi}{2n} e^{-p} +$

$$\frac{\pi}{n} \sum e^{-p \cos \frac{\pi l}{n}} \text{Cos} \{ (2l+1) \frac{\pi l}{n} - p \sin \frac{\pi l}{n} \}$$

where  $p$  is any quantity,  $l$  any integer,  $n$  any odd integer and  $n = 1, 2, 3, 4$  up to  $\frac{n-1}{2}$ .

ii  $\int_0^\infty \frac{(-x^2)^l}{1+x^{2n}} \text{Cos } px \, dx$  where  $n$  is even &  $n = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}$  to  $\frac{n-1}{2}$   
 $= \frac{\pi}{n} \sum e^{-p \cos \frac{\pi l}{n}} \text{Cos} \{ (2l+1) \frac{\pi l}{n} - p \sin \frac{\pi l}{n} \}$

30. i.  $\int_0^\infty \frac{\text{Sin}^{2n+1} x}{x} \, dx = \int_0^\infty \frac{\text{Sin}^{2n+2} x}{x^2} \, dx = \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n)}$

ii. If  $\int_0^\infty \frac{\text{Sin}^n x}{x^p} \, dx = \phi(n, p)$ , then  $(p-1)(p-2) \phi(n, p) = n(n-1) \phi(n-2, p-2) - n^2 \phi(n, p-2)$ . Thus it is possible

to find  $\int_0^\infty \frac{\sin^{2n+1} x}{x^b} dx$   $b$  being any integer.

Cor. 1.  $\int_0^\infty \frac{\sin^{2n+3} x}{x^3} dx = \frac{\sqrt{\pi}}{4} \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n+1)} (n+\frac{1}{2})$

Cor. 2.  $\int_0^\infty \frac{\sin^{2n+1} x}{x^4} dx = \frac{\sqrt{\pi}}{6} \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n+1)} (n+2)$  &c &c &c

N.B. The above theorems are obtained by combining Art. 22. II with the following theorems:—

i.  $\int_0^\infty \frac{\sin^n x}{x^p} dx = \frac{1}{\Gamma(p)} \int_0^\infty \int_0^\infty e^{-zx} z^{p-1} \sin^n x dx dz$

ii.  $\int_0^\infty e^{-ax} \sin^{2n+1} x dx = \frac{\Gamma(2n+1)}{(a^2+1^2)(a^2+3^2)(a^2+5^2)\dots(a^2+(2n+1)^2)}$

iii.  $\int_0^\infty e^{-ax} \sin^{2n} x dx = \frac{\Gamma(2n)}{a(a^2+2^2)(a^2+4^2)(a^2+6^2)\dots(a^2+2n^2)}$

31. i. If  $\int_0^h \phi(x) \cos \pi x dx = \psi(n)$  and  $\alpha\beta = 2\pi$ , then

$\alpha \{ \frac{1}{2} \phi(0) + \phi(\alpha) \cos \pi \alpha + \phi(2\alpha) \cos 2\pi \alpha + \dots + \phi(m\alpha) \cos m\pi \alpha \}$   
 $= \psi(n) + \psi(\beta-n) + \psi(\beta+n) + \psi(2\beta-n) + \psi(2\beta+n) + \dots$

where  $m\alpha$  is the greatest multiple of  $\alpha$  less than  $h$  and  $\pi$  lies between  $0$  &  $\beta$ . If  $h$  be a multiple of  $\alpha$  the last term is  $\frac{1}{2} \phi(h) \cos \pi h$ . (Such conditions are required in similar theorems)

ii.  $\int_0^h \frac{\sin \pi x}{\sin x} \phi(x) dx = \pi \{ \frac{1}{2} \phi(0) - \phi(\pi) \cos \pi + \phi(2\pi) \cos 2\pi - \phi(3\pi) \cos 3\pi + \dots \pm \phi(m\pi) \cos m\pi \}$

$- 2 \psi(m+1) - 2 \psi(m+2) - 2 \psi(m+3) - \dots$  ad. inf; the conditions

being similar to that of i.

Cor. i. If  $\int_0^\infty \phi(x) \cos nx dx = \psi(n)$  and  $\alpha\beta = 2\pi$ , then

$$\alpha \left\{ \frac{1}{2} \phi(0) + \phi(\alpha) + \phi(2\alpha) + \phi(3\alpha) + \dots \text{ad. inf.} \right\}$$

$$= \psi(0) + 2\psi(\beta) + 2\psi(2\beta) + 2\psi(3\beta) + \dots$$

Cor. ii. If  $n$  becomes infinitely great,  $\int_0^h \frac{\sin nx}{\sin x} \phi(x) dx$

$$= \pi \left\{ \frac{1}{2} \phi(0) - \phi(\pi) \cos n\pi + \phi(2\pi) \cos 2n\pi - \dots \pm \phi(m\pi) \cos m\pi \right\}$$

where  $m\pi$  is the greatest multiple of  $\pi$  less than  $h$ .

32. i. If  $\int_0^h \phi(x) \sin nx dx = \psi(n)$  and  $\alpha\beta = 2\pi$ , then

$$\alpha \left\{ \phi(\alpha) \sin n\alpha + \phi(2\alpha) \sin 2n\alpha + \phi(3\alpha) \sin 3n\alpha + \dots + \phi(m\alpha) \sin mn\alpha \right\}$$

$$= \psi(n) - \psi(\beta-n) + \psi(\beta+n) - \psi(2\beta-n) + \dots \text{ad. inf.}$$

with the same condition as in 31.

ii.  $\frac{1}{2} \phi(x) + \phi(x+\alpha) + \phi(x+2\alpha) + \dots \text{ad. inf.}$

$$= \frac{1}{2} \int_0^\infty \phi(x+z) dz - \frac{\beta z}{12} x \phi'(x) + \frac{\beta z}{12} x^2 \phi'''(x) - \dots$$

Cor. If  $\int_0^\infty \phi(x) \sin nx dx = \psi(n)$  and  $\alpha\beta = \frac{\pi}{2}$ , then

$$\alpha \left\{ \phi(\alpha) - \phi(3\alpha) + \phi(5\alpha) - \phi(7\alpha) + \dots \text{ad. inf.} \right\}$$

$$= \psi(\beta) - \psi(3\beta) + \psi(5\beta) - \psi(7\beta) + \dots \text{ad. inf.}$$

P. B. Just as in 31. ii. the following integrals can be found:

$$\int_0^h \frac{\cos nx}{\cos x} \phi(x) dx; \int_0^h \frac{\sin nx}{\cos x} \phi(x) dx; \int_0^h \frac{\cos nx}{\sin x} \phi(x) dx$$



$$33. i. \int_0^{\infty} \left\{ \frac{(-x^2)^{\ell}}{1-x^{2n}} + \frac{(-1)^{\ell}}{n(x^2-1)} \right\} \text{Cosp}x \, dx$$

$$= \frac{\pi}{2n} e^{-p} + \frac{\pi}{n} \sum e^{-p \cos \frac{\pi r}{n}} \cos \left\{ (2\ell+1) \frac{\pi r}{n} - p \sin \frac{\pi r}{n} \right\}$$

where  $n$  is even and  $r = 1, 2, 3, \dots$  up to  $\frac{n-2}{2}$ .

$$ii. \int_0^{\infty} \left\{ \frac{(-x^2)^{\ell}}{1-x^{2n}} + \frac{(-1)^{\ell}}{n(x^2-1)} \right\} \text{Cosp}x \, dx$$

$$= \frac{\pi}{n} \sum e^{-p \cos \frac{\pi r}{n}} \cos \left\{ (2\ell+1) \frac{\pi r}{n} - p \sin \frac{\pi r}{n} \right\}$$

where  $n$  is odd and  $r = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots$  up to  $\frac{n-2}{2}$ .

$$34. i. \frac{\pi \text{Cosp}x}{x \text{Sin} \pi x} = \frac{1}{x^2} + \frac{2 \text{Cosp}x}{1-x^2} - \frac{9 \text{Cosp}2x}{2^2-x^2} + \frac{2 \text{Cosp}3x}{3^2-x^2} - \dots$$

$$ii. \frac{\pi \text{Sin} \theta x}{4x \text{Cos} \frac{\pi x}{2}} = \frac{\text{Sin} \theta}{1-x^2} - \frac{\text{Sin} 3\theta}{3^2-x^2} + \frac{\text{Sin} 5\theta}{5^2-x^2} - \dots$$

$$\text{Cor. i. } \frac{\pi \text{Cosh} \theta x}{x \text{Sin} \pi x} = \frac{1}{x^2} - \frac{2 \text{Cosh} \theta}{1+x^2} + \frac{2 \text{Cosh} 2\theta}{2^2+x^2} - \dots$$

$$ii. \frac{\pi \text{Sin} \theta x}{4x \text{Cosh} \frac{\pi x}{2}} = \frac{\text{Sin} \theta}{1+x^2} - \frac{\text{Sin} 3\theta}{3^2+x^2} + \frac{\text{Sin} 5\theta}{5^2+x^2} - \dots$$

$$35. \sqrt{\alpha} \left\{ 1 + \frac{2}{(1+\alpha^2)^{n+1}} + \frac{2}{(1+4\alpha^2)^{n+1}} + \frac{2}{(1+9\alpha^2)^{n+1}} + \dots \right\}$$

$$= \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n)} \sqrt{\beta} \left\{ 1 + 2e^{-2/\beta} \phi(4/\beta) + 2e^{-4/\beta} \phi(9/\beta) + \dots \right\} \text{ with}$$

$$\alpha\beta = \pi \text{ \& } \phi(t) = \Gamma(n) \Gamma(n) + \frac{n}{\Gamma(n)} \Gamma(n+1) + \frac{n(n-1)}{\Gamma(n)} \Gamma(n-2) \Gamma(n) + \dots$$

$$36. m \left\{ \frac{1}{2(m^2+n^2)} + \frac{1}{m^2+(n+1)^2} + \frac{1}{m^2+(n+2)^2} + \dots \right\}$$

$$= \left( \tan^{-1} \frac{m}{n} + \frac{\beta_2}{2} \cdot \frac{\text{Sin}(2 \tan^{-1} \frac{m}{n})}{m^2+n^2} - \frac{\beta_4}{4} \cdot \frac{\text{Sin}(4 \tan^{-1} \frac{m}{n})}{(m^2+n^2)^2} + \dots \right)$$

$$\text{Cor. } n \left\{ \frac{1}{4n^2} + \frac{1}{n^2+(n+1)^2} + \frac{1}{n^2+(n+2)^2} + \frac{1}{n^2+(n+3)^2} + \dots \right\}$$

$$= \frac{\pi}{4} + \frac{\beta_2}{2} \cdot \frac{1}{2n^2} - \frac{\beta_6}{6} \cdot \frac{1}{8n^6} + \frac{\beta_{10}}{10} \cdot \frac{1}{32n^{10}} - \dots$$

$$1. \quad \frac{1}{x^2(1+\frac{x^2}{2})(1+\frac{x^2}{3})(1+\frac{x^2}{6})(1+\frac{x^2}{10}) \dots}$$

$$= \frac{1}{x^2} - \frac{3}{1+x^2} + \frac{5}{3+x^2} - \frac{7}{6+x^2} + \frac{9}{10+x^2} - \dots$$

$$\text{Cor.} \quad \frac{3}{x(e^{2\pi x\sqrt{2}} - 1)} - \frac{5}{\sqrt{6}(e^{2\pi x\sqrt{6}} - 1)} + \frac{7}{\sqrt{12}(e^{2\pi x\sqrt{12}} - 1)} - \dots$$

$$+ \frac{1}{x} \left\{ \text{sech}\left(\frac{\pi}{x}\sqrt{1-\frac{x^2}{2}}\right) + \text{sech}\left(\frac{\pi}{x}\sqrt{4-\frac{x^2}{3}}\right) + \text{sech}\left(\frac{\pi}{x}\sqrt{9-\frac{x^2}{6}}\right) + \dots \right\}$$

$$= \frac{1}{2\pi x} + \frac{\pi x}{6} - C. \quad \text{for all values of } x$$

$$\text{where } C = \frac{1}{2} + \frac{1}{3+\sqrt{8}} - \frac{1}{5+\sqrt{12}} + \frac{1}{7+\sqrt{18}} - \dots$$

$$= 1 - \frac{\pi}{\sqrt{2}} + \frac{1}{6(3+\sqrt{8})^2} - \frac{1}{10(5+\sqrt{12})^2} + \frac{1}{14(7+\sqrt{18})^2} - \dots$$

N.B. Similarly any function whose denominator is in the form of a product can be expressed as the sum of partial fractions and many other theorems may be deduced from the result.

$$2. \quad \frac{x^2 y^2}{(x+m)(y+n)} = \frac{1}{(m-1)} \left\{ \frac{1}{x+1} \cdot \frac{1-\frac{y}{x}}{1-\frac{y}{x}+n} - \frac{m-1}{1} \cdot \frac{1}{x+2} \cdot \frac{1-\frac{y}{x}}{1-\frac{y}{x}+n} \right.$$

$$\left. + \frac{(m-1)(m-2)}{2} \cdot \frac{1}{x+3} \cdot \frac{1-\frac{3y}{x}}{1-\frac{3y}{x}+n} - \dots \right\} +$$

$$\frac{1}{(n-1)} \left\{ \frac{1}{y+1} \cdot \frac{1-\frac{x}{y}}{1-\frac{x}{y}+m} - \frac{m-1}{1} \cdot \frac{1}{y+2} \cdot \frac{1-\frac{2x}{y}}{1-\frac{2x}{y}+m} + \dots \right\}.$$

$$\text{Cor.} \quad \frac{\pi}{\sin \pi x} \cdot \frac{1}{(m+x)(n-x)} = \frac{1}{x} + \frac{n}{m+1} \cdot \frac{1}{1-x} - \frac{n(n-1)}{(m+1)(m+2)} \cdot \frac{1}{2-x}$$

$$+ \frac{n(n-1)(n-2)}{(m+1)(m+2)(m+3)} \cdot \frac{1}{3-x} - \dots - \frac{m}{n+1} \cdot \frac{1}{1+x} + \frac{m(m-1)}{(n+1)(n+2)} \cdot \frac{1}{2+x}$$

$$- \frac{m(m-1)(m-2)}{(n+1)(n+2)(n+3)} \cdot \frac{1}{3+x} + \dots$$

Cor. 2. 
$$\frac{\frac{\pi}{2} \alpha \beta}{\alpha - \epsilon \beta - \epsilon} = \alpha \left\{ 1 - \frac{\alpha-1}{\beta+1} \cdot \frac{1}{3} + \frac{(\alpha-1)(\alpha-2)}{(\beta+1)(\beta+2)} \cdot \frac{1}{5} - \dots \right\}$$

$$+ \beta \left\{ 1 - \frac{\beta-1}{\alpha+1} \cdot \frac{1}{3} + \frac{(\beta-1)(\beta-2)}{(\alpha+1)(\alpha+2)} \cdot \frac{1}{5} - \dots \right\}$$

3. 
$$1 + \frac{\alpha}{\gamma+1} \cdot \frac{\beta}{\delta+1} + \frac{\alpha(\alpha-1)}{(\gamma+1)(\gamma+2)} \cdot \frac{\beta(\beta-1)}{(\delta+1)(\delta+2)} + \dots$$

$$+ \frac{\gamma}{\alpha+1} \cdot \frac{\delta}{\beta+1} + \frac{\gamma(\gamma-1)}{(\alpha+1)(\alpha+2)} \cdot \frac{\delta(\delta-1)}{(\beta+1)(\beta+2)} + \dots$$

$$= \frac{\alpha \beta \gamma \delta \alpha + \beta + \gamma + \delta}{\alpha + \gamma \beta + \gamma \alpha + \delta \beta + \delta \alpha}$$

4. 
$$\frac{1}{1^2+x^2+\frac{x^4}{12}} + \frac{1}{2^2+x^2+\frac{x^4}{24}} + \frac{1}{3^2+x^2+\frac{x^4}{36}} + \dots$$

$$= \frac{\pi}{2x\sqrt{3}} \frac{\sinh \pi x \sqrt{3} - \sqrt{3} \sin \pi x}{\cosh \pi x \sqrt{3} - \cos \pi x}$$

Cor. If  $n$  be any integer excluding 0,

$$\frac{1}{1^2+(2n)^2+\frac{(2n)^4}{12}} + \frac{1}{2^2+(2n)^2+\frac{(2n)^4}{24}} + \frac{1}{3^2+(2n)^2+\frac{(2n)^4}{36}} + \dots$$

$$= \frac{1}{12n^2} + \frac{1}{2} \left( \frac{1}{1^2+3n^2} + \frac{1}{2^2+3n^2} + \frac{1}{3^2+3n^2} + \dots \right)$$

N.B. A great number of theorems like the above can be got from XIII 29 & 33.

5. i If  $x$  is any integer greater than 0 and  $x$  lies between 0 and  $\frac{\pi}{2n+1}$ , then (both inclusive)

$$\frac{\sin 2nx}{1} + \frac{\sin 2nx}{2} + \frac{\sin 2nx}{3} + \dots = \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma(\frac{1}{2})}{\Gamma(n)}$$

ii. 
$$\frac{\sin 2nx}{x} + \frac{\sin 2nx}{4x} + \frac{\sin 2nx}{9x} + \dots = \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma(\frac{1}{2})}{\Gamma(n)}$$

if  $x$  lies between 0 and  $\frac{\pi}{2n+1}$ . (both inclusive).

N.B. Many series like the above can be got from XIII 30.

$$6. \sqrt{a} \left\{ \frac{1}{2} \operatorname{sech}^{2n} \alpha + \operatorname{sech}^{2n} 2\alpha + \operatorname{sech}^{2n} 3\alpha + \dots \right\}$$

$$= \frac{\sqrt{a}}{2} \sqrt{\beta} \left\{ \frac{1}{2} + \phi(\beta) + \phi(2\beta) + \phi(3\beta) + \dots \right\} \text{ with } \alpha/\beta = \pi$$

$$a = \beta \Rightarrow \sqrt{a} = \frac{1}{\sqrt{\beta}}$$

$$\left\{ 1 - \left(\frac{\beta}{m}\right)^2 \right\} \left\{ 1 + \left(\frac{\beta}{m+1}\right)^2 \right\} \left\{ 1 + \left(\frac{\beta}{m+2}\right)^2 \right\} \dots$$

$$7. e^{\frac{\alpha}{2}} \sqrt{a} \left\{ \frac{1}{2} + e^{-\alpha} \cos n\alpha + e^{-4\alpha} \cos 2n\alpha + e^{-9\alpha} \cos 3n\alpha + \dots \right\}$$

$$= \sqrt{a} \left\{ \frac{1}{2} + e^{-\beta} \cosh n\beta + e^{-4\beta} \cosh 2n\beta + e^{-9\beta} \cosh 3n\beta + \dots \right\}$$

with  $\alpha/\beta = \pi$ .

$$8. \sqrt{a} \left\{ \frac{1}{2} + e^{-\alpha} + e^{-4\alpha} + e^{-9\alpha} + \dots \right\}$$

$$= \sqrt{a} \left\{ \frac{1}{2} + e^{-\beta} + e^{-4\beta} + e^{-9\beta} + \dots \right\} \text{ with } \alpha/\beta = \pi.$$

8. i. If  $\alpha/\beta = \pi$ , then  $\frac{\alpha}{2} \coth n\alpha - \frac{\beta}{4} \cot n\beta$

$$= \frac{\alpha}{2} + \frac{\alpha \sinh n\alpha}{e^{2\alpha} - 1} + \frac{\alpha \sinh 4n\alpha}{e^{4\alpha} - 1} + \frac{\alpha \sinh 6n\alpha}{e^{6\alpha} - 1} + \dots$$

$$+ \frac{\beta \sin 2n\beta}{e^{2\beta} - 1} + \frac{\beta \sin 4n\beta}{e^{4\beta} - 1} + \frac{\beta \sin 6n\beta}{e^{6\beta} - 1} + \dots$$

ii. If  $\alpha/\beta = \pi$ , then  $\frac{\alpha^2}{2} + \frac{1}{2} \log \frac{\sin n\alpha}{\sinh n\beta}$

$$= \left\{ \frac{\alpha^2}{12} + \frac{\cos 2n\alpha}{e^{2\alpha} - 1} + \frac{\cos 4n\alpha}{2(e^{4\alpha} - 1)} + \frac{\cos 6n\alpha}{3(e^{6\alpha} - 1)} + \dots \right\}$$

$$- \left\{ \frac{\beta^2}{12} + \frac{\cosh 2n\beta}{e^{2\beta} - 1} + \frac{\cosh 4n\beta}{2(e^{4\beta} - 1)} + \frac{\cosh 6n\beta}{3(e^{6\beta} - 1)} + \dots \right\}$$

iii. If  $\alpha/\beta = \pi$ , then  $\frac{\alpha^2}{6} \phi(0) + \frac{\alpha}{2} \phi'(0) + \frac{\alpha^2}{24} \phi''(0) +$

$$\frac{\phi(\alpha) + \phi(2\alpha)}{e^{2\alpha} - 1} + \frac{\phi(2\alpha) + \phi(3\alpha)}{2(e^{6\alpha} - 1)} + \frac{\phi(3\alpha) + \phi(4\alpha)}{3(e^{6\alpha} - 1)} + \dots$$

$$+ \phi(\beta) + \frac{1}{2} \phi(2\beta) + \frac{1}{3} \phi(3\beta) + \dots$$

$$= \frac{\beta^2}{6} \phi(0) + \frac{\beta \pi}{2} \phi'(0) + \frac{\pi^2}{24} \phi''(0) +$$

$$\frac{\phi(\alpha) + \phi(-\alpha)}{e^{2\alpha} - 1} + \frac{\phi(2\alpha) + \phi(-2\alpha)}{2(e^{4\alpha} - 1)} + \dots$$

$$+ \phi(\alpha) + \frac{1}{2}\phi(2\alpha) + \frac{1}{3}\phi(3\alpha) + \dots$$

Cor. i. If  $d\beta = \pi^2$ , then  $\frac{\alpha + \beta}{12} = \frac{1}{2} + \frac{2\alpha}{e^{2\alpha} - 1} + \frac{4\alpha}{e^{4\alpha} - 1} + \frac{6\alpha}{e^{6\alpha} - 1}$   
 $+ \frac{8\alpha}{e^{8\alpha} - 1} + \dots + \frac{2\beta}{e^{2\beta} - 1} + \frac{e^{4\beta}}{e^{4\beta} - 1} + \frac{e^{6\beta}}{e^{6\beta} - 1} + \frac{e^{8\beta}}{e^{8\beta} - 1} + \dots$

ii. If  $d\beta = \pi^2$ , then

$$e^{\frac{\alpha - \beta}{12}} = \frac{\sqrt[3]{2} (1 - e^{-2\alpha})(1 - e^{-4\alpha})(1 - e^{-6\alpha}) \dots}{\sqrt[3]{\beta} (1 - e^{-2\beta})(1 - e^{-4\beta})(1 - e^{-6\beta}) \dots}$$

ex.  $\frac{1}{24} - \frac{1}{8\pi} = \frac{1}{e^{2\pi}} + \frac{2}{e^{4\pi}} + \frac{3}{e^{6\pi}} + \frac{4}{e^{8\pi}} + \dots$

9. i. If  $\int_0^h \phi(x) \cos nx dx = \psi(n)$  and  $d\beta = \frac{\pi}{2}$ , then

$$\alpha \{ \phi(\alpha) \sin n\alpha - \phi(3\alpha) \sin 3n\alpha + \dots \pm \phi(m\alpha) \sin mn\alpha \}$$

$$= \frac{\psi(\beta - n) - \psi(\beta + n)}{2} - \frac{\psi(3\beta - n) + \psi(3\beta + n)}{2} + \dots \text{ad. inf.}$$

where  $m\alpha$  is the greatest odd multiple of  $\alpha$  less than  $h$   
 and  $n$  lies between  $-\beta$  &  $\beta$

ii. If  $\int_0^h \phi(x) \sin nx dx = \psi(n)$  and  $d\beta = \frac{\pi}{2}$ , then

$$\alpha \{ \phi(\alpha) \cos n\alpha - \phi(3\alpha) \cos 3n\alpha + \dots \pm \phi(m\alpha) \cos mn\alpha \}$$

$$= \frac{\psi(\beta - n) + \psi(\beta + n)}{2} - \frac{\psi(3\beta - n) + \psi(3\beta + n)}{2} + \dots \text{ad. inf.}$$

with the conditions in the first part.

10.  $e^{\frac{\alpha^2}{2}} \{ e^{-\alpha^2} \sinh n\alpha - e^{-9\alpha^2} \sinh 3n\alpha + e^{-25\alpha^2} \sinh 5n\alpha - \dots \} \sqrt{\alpha}$   
 $= \sqrt{\beta} \{ e^{-\beta^2} \sinh n\beta - e^{-9\beta^2} \sinh 3n\beta + e^{-25\beta^2} \sinh 5n\beta - \dots \}$  with  $\alpha = \frac{\pi}{2}$

$$11. \alpha + \beta = \pi,$$

$$\alpha \left\{ \frac{\cos \alpha d}{e^{\alpha d}} + \frac{\cos 3\alpha d}{e^{3\alpha d}} + \frac{\cos 5\alpha d}{e^{5\alpha d}} + \dots \right\}$$

$$= \beta \left\{ \frac{\cosh 2n\beta}{e^{\beta^2 + e^{-\beta^2}}} + \frac{\cosh 6n\beta}{e^{3\beta^2 + e^{-3\beta^2}}} + \dots \right\}$$

$$12. \alpha + \beta = \frac{\pi}{2}$$

$$\alpha \left\{ \frac{\sin \alpha d}{e^{\alpha d} + e^{-\alpha d}} - \frac{\sin 3\alpha d}{e^{3\alpha d} + e^{-3\alpha d}} + \frac{\sin 5\alpha d}{e^{5\alpha d} + e^{-5\alpha d}} - \dots \right\}$$

$$= \beta \left\{ \frac{\sinh 2n\beta}{e^{\beta^2 + e^{-\beta^2}}} - \frac{\sinh 6n\beta}{e^{3\beta^2 + e^{-3\beta^2}}} + \frac{\sinh 10n\beta}{e^{5\beta^2 + e^{-5\beta^2}}} - \dots \right\}$$

$$13. \alpha + \beta = \frac{\pi}{2}$$

$$\alpha \left\{ \frac{\phi(\alpha) - \phi(-\alpha)}{e^{\alpha} + e^{-\alpha}} - \frac{\phi(3\alpha) - \phi(-3\alpha)}{e^{3\alpha} + e^{-3\alpha}} + \dots \right\}$$

$$+ i\beta \left\{ \frac{\phi(i\beta) - \phi(-i\beta)}{e^{\beta} + e^{-\beta}} - \frac{\phi(3i\beta) - \phi(-3i\beta)}{e^{3\beta} + e^{-3\beta}} + \dots \right\} = 0.$$

13. If  $\alpha + \beta = \pi$  and  $n$  is a positive integer greater than unity,

$$\alpha^n \left\{ \frac{B_{2n}}{4n} \cos \pi n + \frac{1^{2n-1}}{e^{\alpha}} + \frac{2^{2n-1}}{e^{4\alpha}} + \frac{3^{2n-1}}{e^{9\alpha}} + \dots \right\}$$

$$= (\beta)^n \left\{ \frac{B_{2n}}{4n} \cos \pi n + \frac{1^{2n-1}}{e^{\beta}} + \frac{2^{2n-1}}{e^{4\beta}} + \frac{3^{2n-1}}{e^{9\beta}} + \dots \right\}$$

$$\text{Case i. } \frac{1^5}{e^{\pi}} + \frac{2^5}{e^{4\pi}} + \frac{3^5}{e^{9\pi}} + \frac{4^5}{e^{16\pi}} + \dots = \frac{1}{504}$$

$$\text{ii. } \frac{1^9}{e^{\pi}} + \frac{2^9}{e^{4\pi}} + \frac{3^9}{e^{9\pi}} + \frac{4^9}{e^{16\pi}} + \dots = \frac{1}{264}$$

$$\text{iii. } \frac{1^{13}}{e^{\pi}} + \frac{2^{13}}{e^{4\pi}} + \frac{3^{13}}{e^{9\pi}} + \frac{4^{13}}{e^{16\pi}} + \dots = \frac{1}{24}$$

$$\text{iv. } \frac{1^{4n+1}}{e^{\pi}} + \frac{2^{4n+1}}{e^{4\pi}} + \frac{3^{4n+1}}{e^{9\pi}} + \frac{4^{4n+1}}{e^{16\pi}} + \dots = \frac{B_{4n+2}}{8n+4}$$

14. If  $d\beta = \pi^2$  and  $n$  is a positive integer,

$$d^{n+1} \left\{ \frac{1^{2n+1}}{e^{\frac{d}{2}} + e^{-\frac{d}{2}}} - \frac{3^{2n+1}}{e^{\frac{3d}{2}} + e^{-\frac{3d}{2}}} + \frac{5^{2n+1}}{e^{\frac{5d}{2}} + e^{-\frac{5d}{2}}} - \dots \right\}$$

$$+ (\beta)^{n+1} \left\{ \frac{1^{2n+1}}{e^{\frac{\beta}{2}} + e^{-\frac{\beta}{2}}} - \frac{3^{2n+1}}{e^{\frac{3\beta}{2}} + e^{-\frac{3\beta}{2}}} + \frac{5^{2n+1}}{e^{\frac{5\beta}{2}} + e^{-\frac{5\beta}{2}}} - \dots \right\} = 0.$$

Cor. If  $n$  is a positive integer excluding 0,

$$\frac{1^{4n-1}}{e^{\frac{d}{2}} + e^{-\frac{d}{2}}} - \frac{3^{4n-1}}{e^{\frac{3d}{2}} + e^{-\frac{3d}{2}}} + \frac{5^{4n-1}}{e^{\frac{5d}{2}} + e^{-\frac{5d}{2}}} - \dots = 0.$$

15. If  $d\beta = \frac{\pi^2}{4}$ , then

$$\frac{\text{sech } d}{1} - \frac{\text{sech } 3d}{3} + \frac{\text{sech } 5d}{5} - \frac{\text{sech } 7d}{7} + \dots$$

$$+ \frac{\text{sech } \beta}{1} - \frac{\text{sech } 3\beta}{3} + \frac{\text{sech } 5\beta}{5} - \frac{\text{sech } 7\beta}{7} + \dots$$

$$= 2 \left\{ \tan^{-1} e^{-d} - \tan^{-1} e^{-3d} + \tan^{-1} e^{-5d} - \dots \right.$$

$$\left. + \tan^{-1} e^{-\beta} - \tan^{-1} e^{-3\beta} + \tan^{-1} e^{-5\beta} - \dots \right\} = \frac{\pi}{4}.$$

Cor.  $\tan^{-1} e^{-\frac{\pi}{4}} - \tan^{-1} e^{-\frac{3\pi}{4}} + \tan^{-1} e^{-\frac{5\pi}{4}} - \dots = \frac{\pi}{16}.$

16. If  $m$  and  $n$  are positive integers,

$$i. \int_0^{\infty} \frac{\sin^{2m+1} x}{x} \cos^{2n} x \, dx = \frac{\Gamma(m-\frac{1}{2}) \Gamma(n-\frac{1}{2})}{2 \Gamma(m+n)}$$

$$= \int_0^{\infty} \frac{\sin^{2m+2} x}{x^2} \cos^{2n} x \, dx.$$

ii. If  $m, n$  and  $p$  are positive integers,  $(1)^p \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma(n) \Gamma(m-\frac{1}{2})}{\Gamma(n+p) \Gamma(m)}$

$$\int_0^{\infty} \frac{\sin^{2m+1} x}{x} \cos^{2n} x \, dx = \int_0^{\infty} \frac{\sin^{2m+2} x}{x^2} \cos^{2n} x \, dx =$$

17. i. If  $d\beta = 2\pi$  and  $m\alpha$  is the greatest multiple of  $\alpha$  less than  $\frac{\pi}{2}$ , then for all values of  $n$  and  $p$ ,

$$\alpha \left\{ \frac{1}{2} + \cos^2 d \cos p d + \cos^4 2d \cos 2p d + \dots + \cos^{m\alpha} m d \cos p m d \right\}$$

$$= \frac{\pi L}{2^{n+1}} \left\{ \frac{1}{\left| \frac{n+\beta}{2} \right| \left| \frac{n-\beta}{2} \right|} + \left( \frac{1}{\left| \frac{n+\beta-1}{2} \right| \left| \frac{n-\beta+1}{2} \right|} + \frac{1}{\left| \frac{n+\beta+1}{2} \right| \left| \frac{n-\beta-1}{2} \right|} \right) \right.$$

$$\left. + \left( \frac{1}{\left| \frac{n+2\beta-1}{2} \right| \left| \frac{n-2\beta+1}{2} \right|} + \frac{1}{\left| \frac{n+2\beta+1}{2} \right| \left| \frac{n-2\beta-1}{2} \right|} \right) + \&c \text{ to } \infty \right\}$$

ii.  $\alpha \left\{ \cos^m d \sin p d - \cos^m 3d \sin 3p d + \dots \pm \cos^m m d \sin m p d \right\}$

$$= \frac{\pi L m}{2^{n+2}} \left\{ \left( \frac{1}{\left| \frac{n+\beta-p}{2} \right| \left| \frac{n-\beta+p}{2} \right|} - \frac{1}{\left| \frac{n+\beta+p}{2} \right| \left| \frac{n-\beta-p}{2} \right|} \right) \right.$$

$$\left. - \left( \frac{1}{\left| \frac{n+3\beta-p}{2} \right| \left| \frac{n-3\beta+p}{2} \right|} - \frac{1}{\left| \frac{n+3\beta+p}{2} \right| \left| \frac{n-3\beta-p}{2} \right|} \right) + \&c \right\}$$

where  $d\beta = \frac{\pi}{2}$  and  $m\alpha$  is the greatest odd multiple of  $\alpha$  less than  $\frac{\pi}{2}$ . In both the cases if  $m\alpha$  be an exact multiple  $\frac{1}{2}$  the last term must be taken, but there is no such necessity here.

Cor. 1. If  $\alpha$  lies between  $0$  &  $\frac{\pi}{n+1}$  (both inclusive)

$$\alpha \left\{ \frac{1}{2} + \cos^{2\alpha} d + \cos^{2\alpha} 2d + \cos^{2\alpha} 3d + \dots + \cos^{2\alpha} n d \right\}$$

$$= \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma\left(\frac{n}{2}\right)} \text{ where } n \text{ is an integer and } m\alpha \neq \frac{\pi}{2}$$

Cor. 2. But if it lies between  $\frac{\pi}{n}$  &  $\frac{2\pi}{n+1}$  the value is



$$\frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n)} \left( 1 + \frac{2\Gamma(n)}{\Gamma(n+\frac{1}{2})\Gamma(n-\frac{1}{2})} \right)$$

18. If  $\phi(x) = \sum \frac{P_n}{b_n - a_n x}$  and  $\psi(y) = \sum \frac{Q_n}{\gamma_n - b_n y}$ , then

$$\phi(x) \psi(y) = \sum \frac{P_n}{b_n - a_n x} \psi\left(\frac{b_n}{a_n} \cdot \frac{y}{x}\right) + \sum \frac{Q_n}{\gamma_n - b_n y} \phi\left(\frac{\gamma_n}{b_n} \cdot \frac{x}{y}\right)$$

Cor. 1.  $\pi^2 x y n^2 \frac{\cos \theta n x}{\sin \pi n x} \cdot \frac{\cosh \phi n y}{\sinh \pi n y}$

$$= 1 - 2\pi x y n^2 \left\{ \frac{\cos \phi}{1^2 + n^2 y^2} \cdot \frac{\cosh \frac{\theta x}{y}}{\sinh \frac{\pi x}{y}} - \frac{2 \cos 2\phi}{2^2 + n^2 y^2} \cdot \frac{\cosh \frac{2\theta x}{y}}{\sinh \frac{2\pi x}{y}} \right.$$

$$\left. + \frac{3 \cos 3\phi}{3^2 + n^2 y^2} \cdot \frac{\cosh \frac{3\theta x}{y}}{\sinh \frac{3\pi x}{y}} - \dots \right\}$$

$$+ 2\pi x y n^2 \left\{ \frac{\cos \theta}{1^2 - n^2 x^2} \cdot \frac{\cosh \frac{\phi y}{x}}{\sinh \frac{\pi y}{x}} - \frac{2 \cos 2\theta}{2^2 - n^2 x^2} \cdot \frac{\cosh \frac{2\phi y}{x}}{\sinh \frac{2\pi y}{x}} \right.$$

$$\left. + \frac{3 \cos 3\theta}{3^2 - n^2 x^2} \cdot \frac{\cosh \frac{3\phi y}{x}}{\sinh \frac{3\pi y}{x}} - \dots \right\}$$

Cor. 2.  $\frac{\pi}{4 n^2} \cdot \frac{\sin \theta n x}{\cos \frac{\pi n x}{2}} \cdot \frac{\sinh \phi n y}{\cosh \frac{\pi n y}{2}}$

$$= y^2 \left\{ \frac{\sin \phi}{1^2 + n^2 y^2} \cdot \frac{\sinh \frac{\theta x}{y}}{\cosh \frac{\pi x}{2y}} - \frac{\sin 3\phi}{3^2 + n^2 y^2} \cdot \frac{\sinh \frac{3\theta x}{y}}{3 \cosh \frac{3\pi x}{2y}} + \dots \right\}$$

$$+ x^2 \left\{ \frac{\sin \theta}{1^2 - n^2 x^2} \cdot \frac{\sinh \frac{\phi y}{x}}{\cosh \frac{\pi y}{2x}} - \frac{\sin 3\theta}{3^2 - n^2 x^2} \cdot \frac{\sinh \frac{3\phi y}{x}}{3 \cosh \frac{3\pi y}{2x}} + \dots \right\}$$

Cor. 3.  $\frac{\pi}{4} \cdot \frac{\cos \theta n x}{\sin \frac{\pi n x}{2}} \cdot \frac{\sinh \phi n y}{\cosh \frac{\pi n y}{2}}$

$$= \frac{\phi y}{2x} \left\{ \frac{\sin \phi}{1^2 + y^2 n^2} \cdot \frac{\cosh \frac{\theta x}{y}}{\sinh \frac{\pi x}{2y}} - \frac{2 \sin 3\phi}{3^2 + y^2 n^2} \cdot \frac{\cosh \frac{3\theta x}{y}}{3 \sinh \frac{3\pi x}{2y}} + \dots \right\}$$

$$+ n^2 x^2 \left\{ \frac{\cos 2\theta}{2^2 - x^2 n^2} \cdot \frac{\sinh \frac{2\phi y}{x}}{2 \cosh \frac{\pi y}{x}} - \frac{\cos 4\theta}{4^2 - n^2 x^2} \cdot \frac{\sinh \frac{4\phi y}{x}}{4 \cosh \frac{2\pi y}{x}} + \dots \right\}$$

19. i.  $\pi = y \cot \pi x \coth \pi y$

$= 1 + 2\pi xy \left\{ \frac{\coth \frac{\pi x}{y}}{1+y^2} + \frac{2 \coth \frac{2\pi x}{y}}{2^2+y^2} + \frac{3 \coth \frac{3\pi x}{y}}{3^2+y^2} + \dots \right\}$

$- \pi xy \left\{ \frac{\coth \frac{\pi y}{x}}{1-x^2} + \frac{2 \coth \frac{2\pi y}{x}}{2^2-x^2} + \frac{3 \coth \frac{3\pi y}{x}}{3^2-x^2} + \dots \right\}$

ii.  $\frac{\pi^2 xy}{\sin \pi x \sinh \pi y}$

$= 1 - 2\pi xy \left\{ \frac{1}{1+y^2} \cdot \frac{1}{\sinh \frac{\pi x}{y}} - \frac{2}{2^2+y^2} \cdot \frac{1}{\sinh^2 \frac{\pi x}{y}} + \dots \right\}$

$+ 2\pi xy \left\{ \frac{1}{1-x^2} \cdot \frac{1}{\sinh \frac{\pi y}{x}} - \frac{2}{2^2-x^2} \cdot \frac{1}{\sinh^2 \frac{\pi y}{x}} + \dots \right\}$

iii.  $\frac{\pi}{4} \tan \frac{\pi x}{2} \tanh \frac{\pi y}{2}$

$= y^2 \left\{ \frac{\tanh \frac{\pi x}{2y}}{(1+y^2)} + \frac{\tanh \frac{3\pi x}{2y}}{3(3^2+y^2)} + \frac{\tanh \frac{5\pi x}{2y}}{5(5^2+y^2)} + \dots \right\}$

$+ x^2 \left\{ \frac{\tanh \frac{\pi y}{2x}}{(1-x^2)} + \frac{\tanh \frac{3\pi y}{2x}}{3(3^2-x^2)} + \frac{\tanh \frac{5\pi y}{2x}}{5(5^2-x^2)} + \dots \right\}$

iv.  $\frac{\pi}{4} \sec \frac{\pi x}{2} \operatorname{sech} \frac{\pi y}{2}$

$= \frac{\operatorname{sech} \frac{\pi y}{2y}}{1+y^2} - \frac{3 \operatorname{sech} \frac{3\pi y}{2y}}{3^2+y^2} + \frac{5 \operatorname{sech} \frac{5\pi y}{2y}}{5^2+y^2} - \dots$

$+ \frac{\operatorname{sech} \frac{\pi y}{2x}}{1-x^2} - \frac{3 \operatorname{sech} \frac{3\pi y}{2x}}{3^2-x^2} + \frac{5 \operatorname{sech} \frac{5\pi y}{2x}}{5^2-x^2} - \dots$

v.  $\frac{\pi}{4} \cot \frac{\pi x}{2} \operatorname{sech} \frac{\pi y}{2}$

$= \frac{1}{2x} - y \left\{ \frac{\coth \frac{\pi x}{2y}}{1+y^2} - \frac{\coth \frac{3\pi x}{2y}}{3^2+y^2} + \frac{\coth \frac{5\pi x}{2y}}{5^2+y^2} - \dots \right\}$

$- x \left\{ \frac{\operatorname{sech} \frac{\pi y}{x}}{2^2-x^2} + \frac{\operatorname{sech} \frac{2\pi y}{x}}{4^2-x^2} + \frac{\operatorname{sech} \frac{3\pi y}{x}}{6^2-x^2} + \dots \right\}$

N.B. Similarly for  $\tan \frac{\pi x}{2} \coth \frac{\pi y}{2}$  and  $\sec \frac{\pi x}{2} \coth \frac{\pi y}{2}$

$$20. i. \pi^2 x^2 \cot \pi x \operatorname{Coth} \pi x$$

$$= 1 - 4\pi x^4 \left\{ \frac{\operatorname{Coth} \pi}{1^4 - x^4} + \frac{2 \operatorname{Coth} 2\pi}{2^4 - x^4} + \frac{3 \operatorname{Coth} 3\pi}{3^4 - x^4} + \dots \right\}$$

$$\text{Cor. } (\pi x)^2 \frac{\operatorname{Cosh} \pi x \sqrt{x} + \cos \pi x \sqrt{x}}{\operatorname{Cosh} \pi x \sqrt{x} - \cos \pi x \sqrt{x}}$$

$$= 1 + 4\pi x^4 \left\{ \frac{\operatorname{Coth} \pi}{1^4 + x^4} + \frac{2 \operatorname{Coth} 2\pi}{2^4 + x^4} + \frac{3 \operatorname{Coth} 3\pi}{3^4 + x^4} + \dots \right\}$$

$$ii. \pi^2 x^2 \operatorname{Cosec} \pi x \operatorname{Cosech} \pi x$$

$$= 1 + 4\pi x^4 \left\{ \frac{\operatorname{Cosec} \pi}{1^4 - x^4} - \frac{2 \operatorname{Cosec} 2\pi}{2^4 - x^4} + \frac{3 \operatorname{Cosec} 3\pi}{3^4 - x^4} - \dots \right\}$$

$$\text{Cor. } \frac{2\pi^2 x^2}{\operatorname{Cosh} \pi x \sqrt{x} - \cos \pi x \sqrt{x}}$$

$$= 1 - 4\pi x^4 \left\{ \frac{\operatorname{Cosech} \pi}{1^4 + x^4} - \frac{2 \operatorname{Cosech} 2\pi}{2^4 + x^4} + \frac{3 \operatorname{Cosech} 3\pi}{3^4 + x^4} - \dots \right\}$$

$$iii. \frac{\pi}{8x^2} \tan \frac{\pi x}{2} \operatorname{Tanh} \frac{\pi x}{2}$$

$$= \frac{\operatorname{Tanh} \frac{\pi}{2}}{1^4 - x^4} + \frac{3 \operatorname{Tanh} \frac{3\pi}{2}}{3^4 - x^4} + \frac{5 \operatorname{Tanh} \frac{5\pi}{2}}{5^4 - x^4} + \dots$$

$$\text{Cor. } \frac{\pi}{8x^2} \frac{\operatorname{Cosh} \frac{\pi x}{\sqrt{x}} - \cos \frac{\pi x}{\sqrt{x}}}{\operatorname{Cosh} \frac{\pi x}{\sqrt{x}} + \cos \frac{\pi x}{\sqrt{x}}}$$

$$= \frac{\operatorname{Tanh} \frac{\pi}{2}}{1^4 + x^4} + \frac{3 \operatorname{Tanh} \frac{3\pi}{2}}{3^4 + x^4} + \frac{5 \operatorname{Tanh} \frac{5\pi}{2}}{5^4 + x^4} + \dots$$

$$iv. \frac{\pi}{8} \operatorname{Sec} \frac{\pi x}{2} \operatorname{Sech} \frac{\pi x}{2}$$

$$= \frac{1^3 \operatorname{sech} \frac{\pi}{2}}{1^4 - x^4} - \frac{3^3 \operatorname{sech} \frac{3\pi}{2}}{3^4 - x^4} + \frac{5^3 \operatorname{sech} \frac{5\pi}{2}}{5^4 - x^4} - \dots$$

$$\text{Cor. } \frac{\pi/4}{\operatorname{Cosh} \frac{\pi x}{\sqrt{x}} + \cos \frac{\pi x}{\sqrt{x}}}$$

$$= \frac{1^3 \operatorname{sech} \frac{\pi}{2}}{1^4 + x^4} - \frac{3^3 \operatorname{sech} \frac{3\pi}{2}}{3^4 + x^4} + \frac{5^3 \operatorname{sech} \frac{5\pi}{2}}{5^4 + x^4} - \dots$$

21. i. If  $d\alpha = \pi^2$  and  $n$  any integer,

$$\begin{aligned}
& \{ (-d)^{-n} \left\{ \frac{1}{2} S_{2n-1} + \frac{1}{1^{2n-1}} (e^{2\alpha}) + \frac{1}{2^{2n-1}} (e^{4\alpha}) + \dots \right\} \right. \\
& \left. - (-\beta)^{1-n} \left\{ \frac{1}{2} S_{2n-1} + \frac{1}{1^{2n-1}} (e^{2\beta}) + \frac{1}{2^{2n-1}} (e^{4\beta}) + \dots \right\} \right\} \\
& = \frac{B_{2n}}{2^n} \{ (-d)^n + \beta^n \} + \pi^2 \frac{B_2}{2} \frac{B_{2n-2}}{2^{n-2}} \{ (-d)^{n-2} + \beta^{n-2} \} \\
& - \pi^4 \frac{B_4}{2^2} \frac{B_{2n-4}}{2^{n-4}} \{ (-d)^{n-4} + \beta^{n-4} \} + \dots \text{the last term} \\
& \text{being } -\pi^n \frac{B_n}{1^n} \frac{B_n}{1^n} (-1)^{\frac{n}{2}} \text{ or } \pi^{n-1} \frac{B_{n-1}}{1^{n-1}} \frac{B_{n+1}}{1^{n+1}} (-1)^{\frac{n+1}{2}} \{ (-d) + \beta \} \\
& \text{according as } n \text{ is even or odd.}
\end{aligned}$$

ii. If  $d\beta = \pi^2$  and  $n$  any integer, then

$$\begin{aligned}
& d^{1-n} \left\{ \frac{1}{1^{2n-1}} (e^{\frac{d}{2}} + e^{-\frac{d}{2}}) - \frac{1}{3^{2n-1}} (e^{\frac{3d}{2}} + e^{-\frac{3d}{2}}) + \dots \right\} \frac{2^{2n+1}}{\pi} \\
& + (-\beta)^{1-n} \left\{ \frac{1}{1^{2n-1}} (e^{\frac{\beta}{2}} + e^{-\frac{\beta}{2}}) - \frac{1}{3^{2n-1}} (e^{\frac{3\beta}{2}} + e^{-\frac{3\beta}{2}}) + \dots \right\} \frac{2^{2n+1}}{\pi} \\
& = \frac{E_1 E_{2n-1}}{2^{n-2}} \{ (-d)^{n-1} + \beta^{n-1} \} - \frac{E_3 E_{2n-3}}{2^{n-4}} \{ (-d)^{n-3} + \beta^{n-3} \} \\
& + \frac{E_5 E_{2n-5}}{2^{n-6}} \{ (-d)^{n-5} + \beta^{n-5} \} - \dots \text{the last term being} \\
& (-1)^{\frac{n-1}{2}} \left( \frac{E_n}{1^n} \right)^2 \text{ or } (-1)^{\frac{n}{2}} \frac{E_{n-1}}{1^{n-2}} \frac{E_{n+1}}{1^n} (d-\beta) \text{ according as } n \text{ is} \\
& \text{odd or even.}
\end{aligned}$$

iii. If  $d\beta = \pi^2$  and  $n$  any integer,  $\frac{\sqrt{d}}{d^n} \left\{ \frac{1}{1^{2n}} - \frac{1}{3^{2n}} + \frac{1}{5^{2n}} - \dots \right\}$

$$+ \frac{1}{1^{2n}} (e^{-d}) - \frac{1}{3^{2n}} (e^{-3d}) + \frac{1}{5^{2n}} (e^{-5d}) - \dots \} =$$

$$\frac{\sqrt{\beta}}{\beta^n} \left[ (-1)^m \left\{ \frac{1}{2^{2n}(e^\beta + e^{-\beta})} + \frac{1}{4^{2n}(e^{2\beta} + e^{-2\beta})} + \frac{1}{6^{2n}(e^{3\beta} + e^{-3\beta})} + \dots \right. \right. \\ \left. \left. + \frac{1}{4} \left\{ \frac{(\frac{\beta}{2})^{2n}}{12n} E_{2n+1} + \frac{\beta_2}{12} \cdot \frac{E_{2n-1}}{12n-2} \left(\frac{\beta}{2}\right)^{2n-1} (2d) - \frac{\beta_4}{14} \cdot \frac{E_{2n-3}}{12n-4} \left(\frac{\beta}{2}\right)^{2n-2} (2d) \right. \right. \right. \\ \left. \left. \left. + \frac{\beta_6}{16} \cdot \frac{E_{2n-5}}{12n-6} \left(\frac{\beta}{2}\right)^{2n-3} (2d)^3 - \dots - \frac{(\alpha\beta)^m}{12n} B_{2n} E_1 \right\} \right] \right]$$

$$22. i. \frac{\pi^2 xy}{2} \cdot \frac{\cosh \pi(x+y)\sqrt{2} \cos \pi(x-y)\sqrt{2} - \cosh \pi(x-y)\sqrt{2} \cos \pi(x+y)\sqrt{2}}{(\cosh \pi x\sqrt{2} - \cos \pi x\sqrt{2})(\cosh \pi y\sqrt{2} - \cos \pi y\sqrt{2})}$$

$$= 1 + 2\pi x^3 y \left\{ \frac{\coth \frac{\pi y}{x}}{1^4 + x^4} + \frac{2 \coth \frac{2\pi y}{x}}{2^4 + x^4} + \frac{3 \coth \frac{3\pi y}{x}}{3^4 + x^4} + \dots \right\}$$

$$+ 2\pi x y^3 \left\{ \frac{\coth \frac{\pi x}{y}}{1^4 + y^4} + \frac{2 \coth \frac{2\pi x}{y}}{2^4 + y^4} + \frac{3 \coth \frac{3\pi x}{y}}{3^4 + y^4} + \dots \right\}$$

$$ii. \int_0^\infty \frac{\cos 2\pi x}{\cosh \pi\sqrt{x} + \cos \pi\sqrt{x}} dx = \frac{e^{-\pi}}{\cosh \frac{\pi}{2}} - \frac{3e^{-9\pi}}{\cosh \frac{3\pi}{2}} + \frac{5e^{-25\pi}}{\cosh \frac{5\pi}{2}} - \dots$$

Cor. If  $d\beta = \frac{\pi}{4}$ , then

$$\frac{1}{\cosh \sqrt{d} + \cos \sqrt{d}} - \frac{1}{3} \frac{1}{\cosh \sqrt{3d} + \cos \sqrt{3d}} + \frac{1}{5} \frac{1}{\cosh \sqrt{5d} + \cos \sqrt{5d}} - \dots$$

$$+ \frac{1}{\cosh \frac{\pi}{2} \cosh \beta} - \frac{1}{3} \frac{1}{\cosh \frac{3\pi}{2} \cosh 3\beta} + \frac{1}{5} \frac{1}{\cosh \frac{5\pi}{2} \cosh 5\beta} - \dots$$

$$= \frac{\pi}{8}$$

$$iii. \text{ If } d/\beta = 4\pi^2 \text{ and } k = \frac{C_0 + \log_2 \pi}{4} + \frac{1}{e^{2\pi}} + \frac{1}{3(e^{6\pi})} + \frac{1}{5(e^{10\pi})} + \dots$$

$$\frac{7d}{720} + \frac{\cos \sqrt{d}}{1(e^{\sqrt{d}} - 2\cos \sqrt{d} + e^{-\sqrt{d}})} + \frac{\cos \sqrt{3d}}{2(e^{\sqrt{3d}} - 2\cos \sqrt{3d} + e^{-\sqrt{3d}})} + \dots$$

$$= k + \frac{\beta}{48\pi} - \frac{1}{2} \log \beta + \frac{\coth \pi}{1(e^\beta - 1)} + \frac{\coth 2\pi}{2(e^{4\beta} - 1)} + \frac{\coth 3\pi}{3(e^{9\beta} - 1)} + \dots$$

i.e.  $4k = C_0 + 3 \log_2 2 - \frac{\pi}{8} + \log \left| -\frac{1}{2} \right|$ , where  $C_0$  is the constant of  $\frac{1}{2}$

$$23. i. \frac{1}{2\pi} + \coth \pi \left\{ \phi(0) - x^4 \phi(4) + x^8 \phi(8) - \dots \right\}$$

$$+ 2 \coth 2\pi \left\{ \phi(0) - (2x)^4 \phi(4) + (2x)^8 \phi(8) - \dots \right\}$$

$$+ 3 \coth 3\pi \left\{ \phi(0) - (3x)^4 \phi(4) + (3x)^8 \phi(8) - \dots \right\}$$

$$+ \dots \dots \dots = \frac{\pi}{2x^2} \left\{ \frac{1}{2} \phi(2) + h \right\}$$

where  $h$  the error is very nearly equal to  $\phi(2) - \frac{2\pi}{x^{11}} \frac{\phi(-1)}{\sqrt{2}} + \frac{(2\pi)^3}{x^{13}} \frac{\phi(-5)}{\sqrt{2}} + \dots$  the general term being  $\frac{(2\pi)^m}{x^{2m+1}} \cos \frac{3m\pi}{4}$  if  $x$  is small. Similarly

$$ii. \operatorname{sech} \frac{\pi}{2} \left\{ \phi(0) - x^4 \phi(4) + x^8 \phi(8) - \dots \right\}$$

$$- \frac{\operatorname{sech} \frac{3\pi}{2}}{3} \left\{ \phi(0) - (3x)^4 \phi(4) + (3x)^8 \phi(8) - \dots \right\}$$

$$+ \frac{\operatorname{sech} \frac{5\pi}{2}}{5} \left\{ \phi(0) - (5x)^4 \phi(4) + (5x)^8 \phi(8) - \dots \right\}$$

$$- \dots \dots \dots = \frac{\pi}{8} \phi(0) - \frac{\pi}{2} h$$

where  $h$  is very nearly equal to  $\phi(0) - \frac{\pi/\sqrt{2}}{x^{11}} \phi(-1) + \frac{(\pi/\sqrt{2})^3}{x^{13}} \phi(-5) - \dots$  if  $x$  is small.

$$24. \frac{1}{4n^2} + \frac{1}{n^2 + (n+1)^2} + \frac{1}{n^2 + (n+2)^2} + \frac{1}{n^2 + (n+3)^2} + \dots$$

$$= \frac{\pi}{4n} + \frac{1}{8\pi n^3} - \frac{\pi}{n} \cdot \frac{1}{e^{4\pi n} - 2e^{2\pi n} \cos 2\pi n + 1}$$

$$+ 4n \left\{ \frac{1}{e^{2\pi}} \cdot \frac{1}{1+4n^2} + \frac{2}{e^{4\pi}} \cdot \frac{1}{2^2+4n^2} + \frac{3}{e^{6\pi}} \cdot \frac{1}{3^2+4n^2} + \dots \right\}$$

N.B. i.  $\frac{1}{2n^2} + \frac{1}{n^2+1^2} + \frac{1}{n^2+2^2} + \dots = \frac{\pi}{2n} + \frac{\pi}{n} \cdot \frac{1}{e^{2\pi n} - 1}$

ii.  $\frac{1}{n^2+1^2} + \frac{1}{n^2+2^2} + \frac{1}{n^2+3^2} + \dots = \frac{\pi}{4n} - \frac{\pi}{2n} \cdot \frac{1}{e^{\pi n} + 1}$

$$25. \frac{1}{n^2 + (n+1)^2} + \frac{1}{n^2 + (n+3)^2} + \frac{1}{n^2 + (n+5)^2} + \dots$$

$$+ 4n \left\{ \frac{1}{e^{\pi} + 1} \cdot \frac{1}{1^2 + 4n^2} + \frac{3}{e^{3\pi} + 1} \cdot \frac{1}{3^2 + 4n^2} + \frac{5}{e^{5\pi} + 1} \cdot \frac{1}{5^2 + 4n^2} + \dots \right\}$$

$$= \frac{\pi}{8n} - \frac{\pi}{2n} \frac{1}{e^{2\pi n} + 2e^{\pi n} \cos \pi n + 1}$$

$$\text{ex. i. } \frac{\text{coth } \pi}{1^3} + \frac{\text{coth } 2\pi}{2^3} + \frac{\text{coth } 3\pi}{3^3} + \dots = \frac{7\pi^3}{180}$$

$$\text{ii. } \frac{\text{coth } \pi}{1^7} + \frac{\text{coth } 2\pi}{2^7} + \frac{\text{coth } 3\pi}{3^7} + \dots = \frac{19\pi^7}{56700}$$

$$\text{iii. } \frac{\tanh \frac{\pi}{2}}{1^3} + \frac{\tanh \frac{3\pi}{2}}{3^3} + \frac{\tanh \frac{5\pi}{2}}{5^3} + \dots = \frac{\pi^3}{32}$$

$$\text{iv. } \frac{\tanh \frac{\pi}{2}}{1^7} + \frac{\tanh \frac{3\pi}{2}}{3^7} + \frac{\tanh \frac{5\pi}{2}}{5^7} + \dots = \frac{7\pi^7}{23040}$$

$$\text{v. } \frac{\text{cosech } \pi}{1^3} - \frac{\text{cosech } 2\pi}{2^3} + \frac{\text{cosech } 3\pi}{3^3} - \dots = \frac{\pi^3}{360}$$

$$\text{vi. } \frac{\text{cosech } \pi}{1^7} - \frac{\text{cosech } 2\pi}{2^7} + \frac{\text{cosech } 3\pi}{3^7} - \dots = \frac{13\pi^7}{453600}$$

$$\text{vii. } \frac{\text{sech } \frac{\pi}{2}}{1} - \frac{\text{sech } \frac{3\pi}{2}}{3} + \frac{\text{sech } \frac{5\pi}{2}}{5} - \dots = \frac{\pi}{8}$$

$$\text{viii. } \frac{\text{sech } \frac{\pi}{2}}{1^5} - \frac{\text{sech } \frac{3\pi}{2}}{3^5} + \frac{\text{sech } \frac{5\pi}{2}}{5^5} - \dots = \frac{\pi^5}{768}$$

$$\text{ix. } \frac{\text{sech } \frac{\pi}{2}}{1^7} - \frac{\text{sech } \frac{3\pi}{2}}{3^7} + \frac{\text{sech } \frac{5\pi}{2}}{5^7} - \dots = \frac{23\pi^7}{1720320}$$

$$\text{x. } \frac{1}{1^2(e^{\pi}-1)} - \frac{1}{3^2(e^{3\pi}-1)} + \frac{1}{5^2(e^{5\pi}-1)} - \dots$$

$$+ \frac{1}{2^2(e^{\pi}+e^{-\pi})} + \frac{1}{4^2(e^{2\pi}+e^{-2\pi})} + \dots = \frac{5\pi^2}{76} - \frac{1}{2} \int_0^1 \frac{\text{Cot } x}{x} dx.$$

$$\text{xi. } \frac{1}{(1^2+2^2)(\sinh 3\pi - \sinh \pi)} + \frac{1}{(2^2+3^2)(\sinh 5\pi - \sinh \pi)} + \dots$$

$$= \left( \frac{1}{\pi} + \text{coth } \pi - \frac{\pi}{2} \text{Cot}^2 \frac{\pi}{2} \right) / (2 \sinh \pi).$$

$$\text{xii. } \frac{1}{25 \cdot 01(e^{\pi}+1)} + \frac{3}{25 \cdot 81(e^{3\pi}+1)} + \frac{5}{31 \cdot 25(e^{5\pi}+1)} = \frac{\pi \text{Cot}^2 \frac{5\pi}{2}}{8} - \frac{4689}{11890}$$

$$1. h \phi(h) + h \phi(2h) + h \phi(3h) + h \phi(4h) + h \phi(5h) + \dots$$

$= \int_0^{\infty} \phi(x) dx + F(h)$ , where  $F(h)$  can be found by expanding the left and writing the constant instead of a series and  $F(0) = 0$ .

ex. If  $\phi(h) = ah^p + bh^q + ch^r + dh^s + \dots$ , then

$$h \phi(h) + h \phi(2h) + h \phi(3h) + h \phi(4h) + \dots$$

$$= \int_0^{\infty} \phi(x) dx + a \frac{B_p}{p} h^p \cos \frac{\pi p}{2} + b \frac{B_q}{q} h^q \cos \frac{\pi q}{2} + \dots$$

N.B. If the expansion of  $\phi(h)$  be an infinite series, then that of  $F(h)$  also will be an infinite series; but if most of the numbers  $p, q, r, s, \dots$  be odd integers  $F(h)$  appears to terminate. In this case the hidden part of  $F(h)$  can't be expanded in ascending powers of  $h$  and is very rapidly diminishing when  $h$  is slowly diminishing and consequently can be neglected for practical purposes when  $h$  is small. e.g. If  $\phi(h) = \frac{1}{1+h}$  then  $F(h) = \frac{2\pi}{e^{\frac{2\pi}{h}} - 1}$  and hence  $F(\frac{1}{10}) = \frac{2\pi}{e^{20\pi} - 1}$ . If  $\phi(h) = e^{-h^2}$  then  $F(\frac{1}{10})$  is very nearly  $10\sqrt{\pi} e^{-100\pi^2}$ .

$$2. \frac{1^{n-1}}{e^x} + \frac{2^{n-1}}{e^{2x}} + \frac{3^{n-1}}{e^{3x}} + \frac{4^{n-1}}{e^{4x}} + \frac{5^{n-1}}{e^{5x}} + \dots$$

$$= \frac{1^{n-1}}{x^n} + \frac{B_n}{x} \cos \frac{\pi n}{2} - \frac{x}{11} \frac{B_{n+1}}{n+1} \cos \frac{\pi(n+1)}{2} + \dots$$

ex i.  $\frac{C_0 + \log x}{x} + \frac{\log 1}{e^x} + \frac{\log 2}{e^{2x}} + \frac{\log 3}{e^{3x}} + \dots = 2 \log(2\pi)$   
when  $x$  vanishes.

ii. The sum of the nos of factors (including unity and the number) of the first  $n$  natural nos divided by  $n$  when  $n$  is very great  $= 2C_0 - 1 + \log n$ .



iii.  $\log n + n^2 \left( \frac{2}{e^{2n}} + \frac{5}{e^{4n}} + \frac{5}{e^{6n}} + \frac{11}{e^{8n}} + \dots \right)$  is finite when  $n$  vanishes, 2, 3, 5, 7, 11 being prime numbers

iv. If  $I(m)$  be the <sup>is of the order</sup> ~~is of the order~~ integer to  $\frac{1}{m}$   $\left\{ \cosh \pi \sqrt{m} - \frac{\sinh \pi \sqrt{m}}{\pi \sqrt{m}} \right.$   
 then  $I(0) + x I(1) + x^2 I(2) + x^3 I(3) + \dots$

$$= \frac{1}{(1-2x+2x^2-2x^3+2x^4-\dots)}$$

$$3. \frac{1^{m-1}}{e^{1^m x}} + \frac{2^{m-1}}{e^{2^m x}} + \frac{3^{m-1}}{e^{3^m x}} + \frac{4^{m-1}}{e^{4^m x}} + \dots$$

$$= \frac{\frac{m}{2}}{m \cdot 2^{\frac{m}{2}}} + \frac{B_m}{2^m} \cos \frac{\pi m}{2} - \frac{x}{1!} \frac{B_{m+2n}}{m+2n} \cos \frac{\pi(m+2n)}{2} + \frac{x^2}{2!} \frac{B_{m+4n}}{m+4n} \cos \frac{\pi(m+4n)}{2} - \dots$$

$$\text{Cor. } \frac{e^{-1^m x}}{1} + \frac{e^{-2^m x}}{2} + \frac{e^{-3^m x}}{3} + \frac{e^{-4^m x}}{4} + \dots$$

$$= \frac{-C_0 - \log x}{n} + C_0 - \frac{x}{1!} \frac{B_{2n}}{2n} \cos \frac{\pi x}{2} + \frac{x^2}{2!} \frac{B_{4n}}{4n} \cos \pi x - \dots$$

$$\text{ex. i. } \frac{e^{-x}}{1} + \frac{e^{-4x}}{2} + \frac{e^{-9x}}{3} + \frac{e^{-16x}}{4} + \dots$$

$$= \frac{C_0 - \log x}{2} + \frac{x}{12} + \frac{x^2}{240} + \frac{x^3}{1512} + \frac{x^4}{5760} + \frac{x^5}{15840} + \dots$$

$$\text{ii. } e^{-x} + 2e^{-16x} + 2e^{-81x} + 4e^{-256x} + 5e^{-625x} + \dots$$

$$= \frac{1}{4} \sqrt{\frac{\pi}{2}} - \frac{1}{12} + \frac{x}{252} - \frac{x^2}{264} + \frac{x^3}{72} - \dots$$

$$\text{iii. } e^{-x} + 2e^{-8x} + 3e^{-27x} + 4e^{-64x} + \dots$$

$$= \frac{1}{3} \sqrt{\frac{\pi}{3}} - \frac{1}{12} + \frac{x^2}{480} - \frac{x^4}{288} + \dots$$

$$\text{iv. } \frac{e^{-x}}{1} + \frac{e^{-4x}}{4} + \frac{e^{-9x}}{9} + \frac{e^{-16x}}{16} + \dots$$

$$= \frac{\pi^2}{6} - \sqrt{\pi x} + \frac{x}{3} \text{ very nearly.}$$

$$\text{v. } 1^2 e^{-1^2 x} + 2^2 e^{-2^2 x} + 3^2 e^{-3^2 x} + \dots = \frac{1}{6} \sqrt{\frac{\pi}{x}} \text{ very nearly.}$$



$$= \left\{ \frac{1^{\frac{m}{p}}}{m x^{\frac{m}{p}}} S_{1 + \frac{m}{p} \nu - n} \right\} + \left\{ \frac{1^{\frac{m}{q}}}{n x^{\frac{m}{q}}} S_{1 + \frac{m}{q} \nu - m} \right\} +$$

$$\frac{B_m}{m} \cdot \frac{B_n}{n} \cos \frac{\pi m}{2} \cos \frac{\pi n}{2} - \frac{x}{u} \cdot \frac{B_{m+p}}{m+p} \frac{B_{n+q}}{n+q} \cos \frac{\pi(m+p)}{2} \cos \frac{\pi(n+q)}{2}$$

$$+ \frac{x^2}{L} \cdot \frac{B_{m+2p}}{m+2p} \frac{B_{n+2q}}{n+2q} \cos \frac{\pi(m+2p)}{2} \cos \frac{\pi(n+2q)}{2} - \&c.$$

v. B. If  $\frac{m}{p} = \frac{n}{q} = \frac{1}{k}$  the right side becomes

$$\frac{1^{\frac{1}{k}}}{m n x^{\frac{1}{k}}} \left\{ k (\approx \frac{1}{k-1} - c_0 - \log x) + C_0(m+n) \right\} + \frac{B_m}{m} \frac{B_n}{n} \cos \frac{\pi m}{2} \cos \frac{\pi n}{2} - \&c.$$

ex. i.  $\frac{1}{1(e^x - 1)} + \frac{1}{2(e^{2x} - 1)} + \frac{1}{3(e^{3x} - 1)} + \&c.$

$$= \frac{S_2}{x} - \frac{c_0 + \log \frac{2\pi}{x}}{4} - \frac{x}{144} + \frac{x^3}{181440} - \frac{x^5}{3991680}$$

$$+ \frac{x^7}{14515200} - \&c.$$

ii.  $\frac{1^2}{e^x - 1} + \frac{2^2}{e^{2x} - 1} + \frac{3^2}{e^{3x} - 1} + \frac{4^2}{e^{4x} - 1} + \&c$

$$= \frac{2S_3}{x^3} - \frac{1}{12x} + \frac{x}{1440} + \frac{x^3}{181440} + \frac{x^5}{7257600}$$

$$+ \frac{x^7}{159667200} + \&c$$

7.  $\frac{1^m \cdot 1^m}{e^x - 1} + \frac{2^m (2^m + 1)}{e^{2x} - 1} + \frac{3^m (3^m + 1)}{e^{3x} - 1} + \frac{4^m (4^m + 1)}{e^{4x} - 1} + \&c$

(the numerator in the  $n$ th term being  $2^m$  x the sum of the  $n$ th powers of the factors of  $2$ .)

$$= \frac{1^m}{x^{m+1}} S_{m+1} S_{m-n+1} + \frac{1^m}{x^{m+1}} S_{m+1} S_{m-n+1} + \frac{1}{2} S_{1-m} S_{1-n}$$

$$\left\{ \frac{1}{2} S_m S_m + \frac{B_2}{L} x S_{-1-m} S_{-1-n} - \frac{B_6}{L} x^3 S_{-3-m} S_{-3-n} + \&c \right.$$

$$\left. \times 1^4 \left( \frac{1^2}{e^x - 1} + \frac{2^2}{e^{2x} - 1} + \frac{3^2}{e^{3x} - 1} + \&c \right) \right.$$

$$+ 2^4 \left( \frac{1^2}{e^{2x} - 1} + \frac{2^2}{e^{4x} - 1} + \frac{3^2}{e^{6x} - 1} + \&c \right)$$

$$+ 3^4 \left( \frac{1^2}{e^{3x} - 1} + \frac{2^2}{e^{6x} - 1} + \frac{3^2}{e^{9x} - 1} + \&c \right) + \&c \&c =$$

$$\left(\frac{2x^2}{19} - 6x^3\right) S_3 - \frac{x}{19} + \frac{x^3}{1210} - \frac{x^5}{2411} + \dots$$

8. If  $f(x)$  in Ex 1, terminated we do not know how far the result is true. But from the following and similar ways we can calculate the error in such cases; let us take

$$\frac{1}{e^{2n}} + \frac{1}{e^{4n}} + \frac{1}{e^{6n}} + \frac{1}{e^{8n}} + \dots$$

$$= \frac{\pi^2}{6n^2} + \frac{1}{2} \sqrt{\frac{\pi}{2n}} S_{\frac{1}{2}} + \frac{1}{4} \text{ very nearly.}$$

But  $\int_0^{\infty} (e^{-2x} + e^{-4x} + e^{-6x} + \dots) \cos ax \, dx$

$$= \frac{1^2}{1^2+a^2} + \frac{2^2}{2^2+a^2} + \frac{3^2}{3^2+a^2} + \frac{4^2}{4^2+a^2} + \dots$$

$$= \frac{\pi}{2\sqrt{2a}} \frac{\sinh \pi\sqrt{2a} - \sin \pi\sqrt{2a}}{\cosh \pi\sqrt{2a} - \cos \pi\sqrt{2a}}$$

Therefore  $\frac{1}{e^{2n}} + \frac{1}{e^{4n}} + \frac{1}{e^{6n}} + \frac{1}{e^{8n}} + \dots$

$$= \frac{\pi^2}{6n^2} + \frac{1}{2} \sqrt{\frac{\pi}{2n}} S_{\frac{1}{2}} + \frac{1}{4} + \sqrt{\frac{\pi}{2n}} \left\{ \frac{\cos(\frac{\pi}{2} + \sqrt{E}) - e^{-\sqrt{E}} \cos \frac{\pi}{4}}{\cosh \sqrt{E} - \cos \sqrt{E}} \right.$$

$$+ \frac{1}{\sqrt{2}} \frac{\cos(\frac{\pi}{2} + \sqrt{2E}) - e^{-\sqrt{2E}} \cos \frac{\pi}{2}}{\cosh \sqrt{2E} - \cos \sqrt{2E}} + \frac{1}{\sqrt{3}} \frac{\cos(\frac{\pi}{2} + \sqrt{3E}) - e^{-\sqrt{3E}} \cos \frac{\pi}{3}}{\cosh \sqrt{3E} - \cos \sqrt{3E}}$$

$$\left. + \frac{1}{\sqrt{4}} \frac{\cos(\frac{\pi}{2} + \sqrt{4E}) - e^{-\sqrt{4E}} \cos \frac{\pi}{4}}{\cosh \sqrt{4E} - \cos \sqrt{4E}} + \dots \text{ad inf.} \right\}$$

where  $t = \frac{4\pi^3}{n}$ .

9.  $1^m \{ 1^m e^{-x} + 2^m e^{-2x} + 3^m e^{-3x} + 4^m e^{-4x} + \dots \}$

$$+ 2^m \{ 1^m e^{-2x} + 2^m e^{-4x} + 3^m e^{-6x} + 4^m e^{-8x} + \dots \}$$

$$+ 3^m \{ 1^m e^{-3x} + 2^m e^{-6x} + 3^m e^{-9x} + 4^m e^{-12x} + \dots \}$$

$$+ 4^m \{ 1^m e^{-4x} + 2^m e^{-8x} + 3^m e^{-12x} + 4^m e^{-16x} + \dots \}$$

$$+ 5^m \{ 1^m e^{-5x} + 2^m e^{-10x} + 3^m e^{-15x} + 4^m e^{-20x} + \dots \}$$

$$+ \dots \dots \dots =$$

$$\frac{1 \cdot 0 \cdot 1}{x^{m+1}} S_{1+m-n} + \frac{1 \cdot 0 \cdot 1}{x^{n+1}} S_{1+n-m} + S_{-m} S_{-n}$$

$$- \frac{x}{11} S_{-m-1} S_{-n-1} + \frac{x^2}{11} S_{-m-2} S_{-n-2} - \&c.$$

N.B. The value of the above series can be exactly found if  $m+n$  be a positive odd integer. For in that case it can always be expressed in terms of three primary series viz

- i.  $1 - 24 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^2} + \frac{9x^3}{1-x^3} + \frac{4x^4}{1-x^4} + \&c \right) = L.$
- ii.  $1 + 240 \left( \frac{x^2}{1-x^2} + \frac{2^3 x^4}{1-x^4} + \frac{3^3 x^6}{1-x^6} + \frac{4^3 x^8}{1-x^8} + \&c \right) = M.$
- iii.  $1 - 504 \left( \frac{x^3}{1-x^3} + \frac{2^5 x^5}{1-x^5} + \frac{3^5 x^7}{1-x^7} + \frac{4^5 x^9}{1-x^9} + \&c \right) = N.$

10. i.  $\frac{B_{2n}}{4^n} \cos \pi n + \frac{1^{2n-1} x}{1-x} + \frac{2^{2n-1} x^2}{1-x^2} + \frac{3^{2n-1} x^3}{1-x^3} + \&c$  Can be expressed in terms of  $M$  and  $N$  only and the series

$$\frac{1^{2n} x}{(1-x)^2} + \frac{2^{2n} x^2}{(1-x^2)^2} + \frac{3^{2n} x^3}{(1-x^3)^2} + \&c \text{ (the diff. of the above series)}$$

$$- \frac{\pi L}{6} \left\{ \frac{B_{2n}}{4^n} \cos \pi n + \frac{1^{2n-1} x}{1-x} + \frac{2^{2n-1} x^2}{1-x^2} + \frac{3^{2n-1} x^3}{1-x^3} + \&c \right\}$$

Can also be expressed in terms of  $M$  &  $N$  only by using indeterminate Coeff<sup>ts</sup>. pay attention to the degree. Thus by successive differentiations the double series in XV can be expressed in terms of  $L, M$  and  $N$ .

ii. The degree of a series is the sum of the highest powers of the  $n$ th terms together with unity if the series contains all the powers of  $x$  or if the powers of  $x$  be in A.P.

If the Coeff<sup>ts</sup> of each  $n$ th term is homogeneous the series is said to be pure and in other cases mixed.

The theory of indices holds good in terms of degrees of series.

If  $F(x)$  in XVI 1. terminates the series is said to be perfect or not it is said to be imperfect.

If  $\phi(x)$  the series is said to be complete in other cases incomplete. 183

A series is said to be absolutely complete when it remains complete when transformed or split up. A linear sc can only be expressed by linear, double by double, treble by treble, pure by pure, perfect by perfect, imperfect, <sup>by imperfect</sup> and absolutely complete by absolutely complete adhering to the laws of indices in all cases. But a mixed series can be split up into a number of pure series of different degrees.

e.g.  $1^n x + 2^n x^2 + 3^n x^3 + \dots$  is an imperfect, incomplete, pure, linear series of the  $(n+1)$ th degree.

$\frac{\sin x}{1} + \frac{\sin 2x}{2} + \frac{\sin 3x}{3} + \dots$  is a perfect, incomplete, pure, linear series of 0 degree.

The series in Art 9 is a perfect, incomplete, pure, double series of  $(m+n+1)$ th degree if  $m+n$  be odd and imperfect if  $m+n$  be even.

The series in Art 7 is a perfect, incomplete, pure, treble series of  $(m+n+2)$ th degree except when both  $m$  &  $n$  be even.

$\frac{1^m}{(e^x + e^{-x})^m} + \frac{2^m}{(e^{2x} + e^{-2x})^m} + \frac{3^m}{(e^{3x} + e^{-3x})^m} + \dots$  is always a mixed, incomplete double series of  $(m+1)$ th degree if  $m = 2, 4, 6, 8, \dots$  is a perfect, complete, pure double series of  $\frac{1}{2}$  a degree.

$L, M$  and  $N$  are perfect, pure double series of  $2m, 4$ th and  $6$ th degree respectively,  $M$  &  $N$  being complete and  $L$  incomplete.

11. If  $d\beta = T^2$ , then

$$\begin{aligned} & \frac{1^2(e^{\alpha} - e^{-\alpha})^2}{1} + \frac{2^2(e^{2\alpha} - e^{-2\alpha})^2}{1} + \frac{3^2(e^{3\alpha} - e^{-3\alpha})^2}{1} + \dots \\ & + \frac{1^2(e^{\beta} - e^{-\beta})^2}{1} + \frac{2^2(e^{2\beta} - e^{-2\beta})^2}{1} + \frac{3^2(e^{3\beta} - e^{-3\beta})^2}{1} + \dots \\ & - 2\alpha \{ 1^2 \log(1 - e^{-2\alpha}) + 2^2 \log(1 - e^{-4\alpha}) + 3^2 \log(1 - e^{-6\alpha}) + \dots \} \\ & - 2\beta \{ 1^2 \log(1 - e^{-2\beta}) + 2^2 \log(1 - e^{-4\beta}) + 3^2 \log(1 - e^{-6\beta}) + \dots \} \\ & = \frac{\alpha^2 + \beta^2}{120} - \frac{\alpha\beta}{72}. \end{aligned}$$

v. b. The theorem in XVIII 24. ii and similar theorems are true only in case of a linear series but approximately in case of other series.

12. i.  $M^3 - N^2 = 1728 x (1-x)^{24} (1-x^2)^{24} (1-x^3)^{24} (1-x^4)^{24} \dots$

ii.  $1 + 480 \left( \frac{1^7 x}{1-x} + \frac{2^7 x^2}{1-x^2} + \frac{3^7 x^3}{1-x^3} + \dots \right) = M^2$

iii.  $1 - 264 \left( \frac{1^5 x}{1-x} + \frac{2^5 x^2}{1-x^2} + \frac{3^5 x^3}{1-x^3} + \dots \right) = MN$

iv.  $1 - 24 \left( \frac{1^3 x}{1-x} + \frac{2^3 x^2}{1-x^2} + \frac{3^3 x^3}{1-x^3} + \dots \right) = M^2 N$

v.  $\frac{1^2 x}{(1-x)^2} + \frac{2^2 x^2}{(1-x^2)^2} + \frac{3^2 x^3}{(1-x^3)^2} + \dots = \frac{M - L^2}{288}$

vi.  $\frac{1^4 x}{(1-x)^4} + \frac{2^4 x^2}{(1-x^2)^4} + \frac{3^4 x^3}{(1-x^3)^4} + \dots = \frac{LM - N}{720}$

vii.  $\frac{1^6 x}{(1-x)^6} + \frac{2^6 x^2}{(1-x^2)^6} + \frac{3^6 x^3}{(1-x^3)^6} + \dots = \frac{M^2 - LN}{1008}$

viii.  $\frac{1^8 x}{(1-x)^8} + \frac{2^8 x^2}{(1-x^2)^8} + \frac{3^8 x^3}{(1-x^3)^8} + \dots = \frac{LM^2 - MN}{720}$

ix.  $L = \frac{1^3 - 3^3 x + 5^3 x^3 - 7^3 x^6 + 9^3 x^{10} - \dots}{1 - 3x + 5x^3 - 7x^6 + 9x^{10} - \dots}$

x.  $M = \left\{ \frac{1^5 x}{1-x} + \frac{3^5 x^3}{1-x^3} + \frac{5^5 x^5}{1-x^5} + \frac{7^5 x^7}{1-x^7} + \dots \right\}$

$= \left\{ \frac{x}{1-x} + \frac{3x^3}{1-x^3} + \frac{5x^5}{1-x^5} + \frac{7x^7}{1-x^7} + \dots \right\}$

i.  $641 + 65520 \left( \frac{1^{11}x}{1-x} + \frac{2^{11}x^2}{1-x^2} + \frac{3^{11}x^3}{1-x^3} + \dots \right)$

$= 241 M^3 + 250 N^2$

ii.  $3617 + 16320 \left( \frac{1^{15}x}{1-x} + \frac{2^{15}x^2}{1-x^2} + \frac{3^{15}x^3}{1-x^3} + \dots \right)$

$= 1617 M^4 + 2000 MN^2$

iii.  $43867 - 28728 \left( \frac{1^{17}x}{1-x} + \frac{2^{17}x^2}{1-x^2} + \frac{3^{17}x^3}{1-x^3} + \dots \right)$

$= 38367 M^3 N + 5500 N^3$

iv.  $174611 + 13200 \left( \frac{1^{19}x}{1-x} + \frac{2^{19}x^2}{1-x^2} + \frac{3^{19}x^3}{1-x^3} + \dots \right)$

$= 53361 M^5 + 121250 M^2 N^2$

v.  $77683 - 552 \left( \frac{1^{21}x}{1-x} + \frac{2^{21}x^2}{1-x^2} + \frac{3^{21}x^3}{1-x^3} + \dots \right)$

$= 57183 M^4 N + 20500 MN^3$

vi.  $236364091 + 131040 \left( \frac{1^{23}x}{1-x} + \frac{2^{23}x^2}{1-x^2} + \frac{3^{23}x^3}{1-x^3} + \dots \right)$

$= 49679091 M^6 + 176400000 M^3 N^2 + 10385000 N^4$

vii.  $657931 - 24 \left( \frac{1^{25}x}{1-x} + \frac{2^{25}x^2}{1-x^2} + \frac{3^{25}x^3}{1-x^3} + \dots \right)$

$= 392931 M^5 N + 265000 M^2 N^3$

viii.  $3392780147 + 6960 \left( \frac{1^{27}x}{1-x} + \frac{2^{27}x^2}{1-x^2} + \frac{3^{27}x^3}{1-x^3} + \dots \right)$

$= 489693897 M^7 + 2507636250 M^4 N^2 + 395450000 MN^4$

ix.  $1723168255201 - 171864 \left( \frac{1^{29}x}{1-x} + \frac{2^{29}x^2}{1-x^2} + \frac{3^{29}x^3}{1-x^3} + \dots \right)$

~~$= 6742902481 M^6 N + 1716211002720 M^3 N^3 + 215050000 N^5$~~

$= 815806500201 M^6 N + 881340705000 M^2 N^3$

$+ 26021050000 N^5$



$$\begin{aligned} & 2. 7709321041217 + 32640 \left( \frac{1^{11}x}{1-x} + \frac{2^{11}x^2}{1-x^2} + \frac{3^{11}x^3}{1-x^3} + \dots \right) \\ & = 764412173217 M^8 + 5323905468000 M^6 N^2 \\ & \quad + 1621003400000 M^2 N^4. \end{aligned}$$

$$N.B. \quad x \frac{dL}{dx} = \frac{L^2 - M}{12}; \quad x \frac{dM}{dx} = \frac{LM - N}{8} \text{ and } x \frac{dN}{dx} = \frac{LN - M^2}{2}.$$

$$\begin{aligned} \text{ex. i. } & 1^5 (1^6 x + 2^6 x^2 + 3^6 x^3 + 4^6 x^4 + \dots) \\ & + 2^6 (1^6 x^2 + 2^6 x^4 + 3^6 x^6 + 4^6 x^8 + \dots) \\ & + 3^6 (1^6 x^3 + 2^6 x^6 + 3^6 x^9 + 4^6 x^{12} + \dots) \\ & + 4^6 (1^6 x^4 + 2^6 x^8 + 3^6 x^{12} + 4^6 x^{16} + \dots) \\ & + \dots \quad \dots \quad \dots \quad \dots \end{aligned}$$

$$= (15LM^2 + 10L^3M - 20L^2N - 4MN - L^5) / 12^4$$

$$\begin{aligned} \text{ii. } & 1^7 (1^7 x + 2^7 x^2 + 3^7 x^3 + 4^7 x^4 + \dots) \\ & + 2^7 (1^7 x^2 + 2^7 x^4 + 3^7 x^6 + 4^7 x^8 + \dots) \\ & + 3^7 (1^7 x^3 + 2^7 x^6 + 3^7 x^9 + 4^7 x^{12} + \dots) \\ & + 4^7 (1^7 x^4 + 2^7 x^8 + 3^7 x^{12} + 4^7 x^{16} + \dots) \\ & + \dots \quad \dots \quad \dots \quad \dots \end{aligned}$$

$$= \frac{2LM^2 - MN - L^2N}{12^3}$$

$$\begin{aligned} \text{iii. } & 1^8 (1^8 x + 2^8 x^2 + 3^8 x^3 + 4^8 x^4 + \dots) \\ & + 2^8 (1^8 x^2 + 2^8 x^4 + 3^8 x^6 + 4^8 x^8 + \dots) \\ & + 3^8 (1^8 x^3 + 2^8 x^6 + 3^8 x^9 + 4^8 x^{12} + \dots) \\ & + 4^8 (1^8 x^4 + 2^8 x^8 + 3^8 x^{12} + 4^8 x^{16} + \dots) \\ & + \dots \quad \dots \quad \dots \quad \dots \end{aligned}$$

$$= (L^3M - 3L^2N + 3LM^2 - MN) / 3456.$$

14.  $S_n = \frac{B_n}{2^n} +$

$$(-1)^n \left\{ \frac{x^1}{1-x} + \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} + \dots \right\} \text{ such that } S_8 = 120 S_4^2$$

then  $\frac{(n+1)(n+3)}{2} S_{n+2} + \frac{n(n-1)(n-2)(n-7)}{12} (n-2)(n-13) S_4 S_{n-2}$

$$+ \frac{n(n-1)(n-2)(n-3)(n-4)(n-5)}{12} (n-7)(n-19) S_6 S_{n-4}$$

$$+ \frac{n(n-1)\dots(n-7)}{16} \left\{ (n-12)(n-23) - 5 \cdot 6 \right\} S_8 S_{n-6}$$

$$+ \frac{n(n-1)\dots(n-9)}{18} \left\{ (n-17)(n-28) - 10 \cdot 7 \right\} S_{10} S_{n-8}$$

$$+ \frac{n(n-1)\dots(n-11)}{110} \left\{ (n-22)(n-32) - 15 \cdot 8 \right\} S_{12} S_{n-10} + \dots$$

N.B. If the last term be a perfect square then half the term must be taken.

15.  $1 + \frac{1}{2} \cdot \frac{2t}{1+t} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \left( \frac{2t}{1+t} \right)^2 + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \left( \frac{2t}{1+t} \right)^3 + \dots$

$$= (1+t) \left\{ 1 + \frac{1}{2} t^2 + \frac{1 \cdot 3}{2 \cdot 2} t^4 + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} t^6 + \dots \right\}; \text{ thus we see,}$$

if  $d = \frac{2t}{1+t}$  and  $\beta = t^2$ ,  $\beta$  is in the 2nd degree of  $d$ .

By supposing  $t^2 = \frac{2t}{1+t}$ ,  $\frac{2t}{1+t} = d$  and  $t^2 = \beta$  we see that

$\beta$  is in the 4th degree of  $d$ , and so on. The relation between  $d$  and  $\beta$  is the modular equation of the degree 4, and the ratio between the two series is denoted by  $M$ . Thus for the

2nd —  $M = 1 + \sqrt{\beta} = \sqrt{1 - \frac{\beta}{\alpha}} = \sqrt{(1-\alpha)(1-\alpha) + 2\sqrt{\beta}}$

3rd —  $M = 1 + 2\sqrt{\frac{\beta}{\alpha}} = \sqrt{\frac{1-\beta}{1-\alpha}}$

$n$ th —  $\beta = \frac{4\alpha^n}{\left\{ (1+\sqrt{1-\alpha})^n + (1-\sqrt{1-\alpha})^n \right\}^2}$

Cor. If end be  $\alpha^2 + 2\alpha = \beta$ , then the  $n$ th is  $\beta = (\alpha+1)^n - 1$ .

ii. If  $p$ th and  $q$ th be  $\phi(x)$  and  $\psi(x)$  and  $r$ th be  $f(x)$ , then if  $p$ th and  $q$ th be  $\phi F(x)$  and  $\psi F(x)$  then  $r$ th is  $f F(x)$ , and also if  $p$ th and  $q$ th be  $F\phi(x)$  and  $F\psi(x)$  then  $r$ th is  $Ff(x)$ .

Cor. Thus we may add or subtract any constant and multiply or divide by any constant to  $x$  in each function or to each function.

Cor. i. If 1st is  $x$  and 2nd  $x^2 + 4x$  then  $n$ th =  $\left\{ \left( \frac{\sqrt{x+4} + \sqrt{x}}{2} \right)^n - \left( \frac{\sqrt{x+4} - \sqrt{x}}{2} \right)^n \right\}$   
 ii.  $\dots \dots \dots x^2 - 2 \dots \dots \dots = \left( \frac{x + \sqrt{x^2 - 4}}{2} \right)^n + \left( \frac{x - \sqrt{x^2 - 4}}{2} \right)^n$

iii. If  $f(x)$  and  $F(x)$  be of the  $p$ th and  $q$ th degree, find  $\phi(x)$  such that  $\sqrt[p]{\phi f(x)} = \sqrt[q]{\phi F(x)} = \chi(x)$  suppose, then the function for the  $n$ th degree =  $\phi^{-1} \{ \chi(x) \}^n$  and the self-repeating series is  $\sqrt[n]{\frac{\phi(x)}{\psi(x) \phi(x)}}$  where  $n$  is any quantity and  $\psi(x)$  any suitable function. Supposing the series to be  $S(x)$  we have  $\frac{S F(x)}{S f(x)} = \sqrt[n]{\frac{\psi}{\phi} \cdot \frac{\psi f(x)}{\psi F(x)} \cdot \frac{F'(x)}{f'(x)}}$

ex. If  $I = x$  and  $II = x^2 + 2nx$ , then if  $x$  is great

$$\frac{III}{II} = x^3 + 3nx^2 + \frac{2n(n+1)}{2}x - \frac{n(n-1)(n-2)x}{2x + \frac{2n(n+1)}{2}} \text{ nearly.}$$

16. If the modular equation for the  $(n-1)$ th degree be

$$\sqrt[n]{a\alpha} + \sqrt[n]{(1-\alpha)(1-\alpha)} = 1$$

$$\text{then that of the } (n-1)\text{th is } \left\{ \sqrt[n]{a(1-\alpha)} - \sqrt[n]{\alpha(1-\alpha)} \right\}^n =$$

$$(\sqrt[n]{a} - \sqrt[n]{\alpha})^n + (\sqrt[n]{1-\alpha} - \sqrt[n]{1-\alpha})^n = 1$$

17. B. The above result is got by eliminating  $\alpha$  from the equations  $\sqrt[n]{\alpha\alpha} + \sqrt[n]{(1-\alpha)(1-\alpha)} = 1$  &  $\sqrt[n]{a\alpha} + \sqrt[n]{(1-\alpha)(1-\alpha)} = 1$

$$1. \quad \Pi(a, x) = (1+a)(1+ax)(1+ax^2)(1+ax^3)(1+ax^4) \&c.$$

$$i. \quad \frac{\Pi(a, x)}{\Pi(a, x)} = (1+a)^n \text{ when } x=1.$$

$$ii. \quad \frac{\Pi(a, x)}{(1-x)^n \Pi(-x^{n+1}, x)} = \sqrt{x} \text{ when } x=1.$$

$$iii. \quad \Pi(a, x) = \Pi(a, x^n) \Pi(ax, x^{2n}) \Pi(ax^2, x^{3n}) \dots \dots \dots \Pi(ax^{n-1}, x^n).$$

$$iv. \quad \Pi(a, x) = \frac{\Pi(a, \sqrt{x})}{\Pi(a/\sqrt{x}, x)}.$$

$$2. \quad \frac{\Pi(b, x)}{\Pi(-a, x)} = 1 + \frac{a+b}{1-x} + \frac{(a+b)(a+bx)}{(1-x)(1-x^2)} + \frac{(a+b)(a+bx)(a+bx^2)}{(1-x)(1-x^2)(1-x^3)} + \dots$$

$$3. \quad \frac{\Pi(ax, x)}{\Pi(-a, x)} = 1 + \frac{ax}{(1-x)(1-ax)} + \frac{a^2x^2}{(1-x)(1-x^2)(1-ax)(1-ax^2)} + \dots$$

$$4. \quad \frac{\Pi(-ab, x) \Pi(-ac, x)}{\Pi(-a, x) \Pi(-abc, x)} = 1 + a \cdot \frac{(1-b)(1-c)}{(1-a)(1-x)} + a^2 \frac{(1-b)(x-b)(1-c)(x-c)}{(1-a)(1-ax)(1-x)(1-x^2)} + \dots$$

$$5. \quad \frac{\Pi(a, x) \Pi(abc, x) \Pi(-abd, x) \Pi(-acd, x)}{\Pi(-ab, x) \Pi(-ac, x) \Pi(-ad, x) \Pi(-abcd, x)} = 1 - a \frac{(1-b)(1-c)(1-d)}{(1-ab)(1-ac)(1-ad)} \cdot \frac{1-ax}{1-x} + a^2 \frac{(1-b)(x-b)(1-c)(x-c)}{(1-ab)(1-abx)(1-ac)(1-acx)} \times \frac{(1-d)(x-d)}{(1-ad)(1-adx)} \cdot \frac{(1-ax^2)(1-a)}{(1-x)(1-x^2)} - a^3 \frac{(1-b)(x-b)(x^2-b)}{(1-ab)(1-abx^2)(1-ac)(1-acx)} \cdot \frac{(1-e)(x-c)}{(1-ae)(1-aex)} \times \frac{(x^2-c)(1-d)(x-d)(x^2-d)}{(1-aex^2)(1-ad)(1-adx)(1-adx^2)} \cdot \frac{(1-ax^3)(1-a)(1-ax)}{(1-x)(1-x^3)(1-x^2)} + \dots$$

$$6. 1 + \frac{a-b}{1-x} \cdot \frac{1-c}{1-d} + \frac{(a-b)(a-bx)}{(1-x)(1-x^2)} \cdot \frac{(1-c)(1-cx)}{(1-d)(1-dx)} + \&c$$

$$= \frac{\Pi(b, x) \Pi(c, x)}{\Pi(a, x) \Pi(d, x)} \left\{ 1 + \frac{c-d}{1-x} \cdot \frac{1-a}{1-b} + \frac{(c-d)(c-dx)}{(1-x)(1-x^2)} \cdot \frac{(1-a)(1-ax)}{(1-b)(1-bx)} + \&c \right\}$$

$$7. \frac{\Pi(-a, x) \Pi(-d, x)}{\Pi(-b, x) \Pi(-c, x)} \left\{ 1 + \frac{a-b}{1-x} \cdot \frac{a-c}{a-d} + \frac{(a-b)(a-bx)(a-c)(a-cx)}{(1-x)(1-x^2)(a-d)(a-dx)} + \&c \right\}$$

$$= 1 + \frac{1-dx}{1-x} \cdot \frac{1-a}{a-d} \cdot \frac{b-d}{1-b} \cdot \frac{c-d}{1-c} + x^2 \frac{(1-dx^2)(1-d)}{(1-x)(1-x^2)} \cdot \frac{(1-a)(1-ax)}{(a-d)(a-dx)} +$$

$$\frac{(b-d)(b-dx)(c-d)(c-dx)}{(1-b)(1-bx)(1-c)(1-cx)} + x^4 \frac{(1-dx^4)(1-d)(1-dx)(1-a)(1-ax)(1-ax^2)}{(1-x)(1-x^2)(1-x^4)(a-d)(a-dx)(a-dx^2)} +$$

$$\times \frac{(b-d)(b-dx)(b-dx^2)(c-d)(c-dx)(c-dx^2)}{(1-b)(1-bx)(1-bx^2)(1-c)(1-cx)(1-cx^2)} + \&c$$

$$8. \frac{\Pi(a, x)}{\Pi(-b, x)} \left\{ 1 + \frac{a-b}{1-x} \cdot \frac{1-c}{1-d} + \frac{(a-b)(a-bx)}{(1-x)(1-x^2)} \cdot \frac{(1-c)(1-cx)}{(1-d)(1-dx)} + \&c \right\}$$

$$= 1 + \frac{1}{1-b} \cdot \frac{a-b}{1-x} \cdot \frac{d-c}{1-d} + \frac{x}{(1-b)(1-bx)} \cdot \frac{(a-b)(a-bx)}{(1-x)(1-x^2)} \cdot \frac{(d-c)(d-cx)}{(1-d)(1-dx)} +$$

$$\frac{x^3}{(1-b)(1-bx)(1-bx^2)} \cdot \frac{(a-b)(a-bx)(a-bx^2)}{(1-x)(1-x^2)(1-x^4)} \cdot \frac{(d-c)(d-cx)(d-cx^2)}{(1-d)(1-dx)(1-dx^2)} + \&c$$

$$9. \Pi(ax, x) \left\{ 1 + \frac{bx}{(1-x)(1-ax)} + \frac{b^2x^2}{(1-x)(1-x^2)(1-ax)(1-ax^2)} + \&c \right\}$$

$$= 1 - x \cdot \frac{a-b}{1-x} + x^3 \frac{(a-b)(a-bx)}{(1-x)(1-x^2)} - x^5 \frac{(a-b)(a-b)(a-bx^2)}{(1-x)(1-x^2)(1-x^4)} + \&c$$

$$\text{Coroll. } 1 + \frac{x^2}{(1-x)^2} + \frac{x^6}{(1-x)^2(1-x^4)^2} + \frac{x^{12}}{(1-x)^2(1-x^4)^2(1-x^8)^2} + \&c$$

$$= \frac{1-x+x^2+x^6+x^{10}-x^{14}+x^{24}-\&c}{(1-x)(1-x^4)(1-x^2)(1-x^4)(1-x^8)\&c}$$

$$\text{ii. } 1 + \frac{x^3}{(1-x)(1-x^2)} + \frac{x^{10}}{(1-x)(1-x^2)(1-x^3)(1-x^6)} + \&c$$

$$= \frac{1-x+x^3-x^9+x^{16}-\&c}{(1-x)(1-x^3)(1-x^6)(1-x^9)\&c}$$

$$10. \frac{1}{x^2} = \frac{\frac{x+l+n-m-1}{2}}{\frac{x+l+n-m-1}{2}} \cdot \frac{\frac{x+l-n-m-1}{2}}{\frac{x+l-n-m-1}{2}} \cdot \frac{\frac{x-l+n+m-1}{2}}{\frac{x-l+n+m-1}{2}} \cdot \frac{\frac{x-l-n+m-1}{2}}{\frac{x-l-n+m-1}{2}}$$

then  $\frac{1}{x^2} = \frac{2lmx}{x^2+l^2+m^2-n^2-1} + \frac{4(x^2-1)(l^2-1)(m^2-1)}{3(x^2+l^2+m^2-n^2-5)} + \dots$   
 11. B. Here the expansion in ascending powers of  $\frac{1}{x}$  is true.  
 But if  $x$  be removed from the numerators, then the results will be true always.

$$11. \frac{\Gamma(a, x) \Gamma(-b, x) - \Gamma(-a, x) \Gamma(b, x)}{\Gamma(a, x) \Gamma(b, x) + \Gamma(-a, x) \Gamma(-b, x)}$$

$$= \frac{a-b}{1-x} + \frac{(a-bx)(ax-b)}{1-x^3} + \frac{x(a-bx^2)(ax^2-b)}{1-x^5} + \frac{x^4(a-bx^3)(ax^3-b)}{1-x^7} + \dots$$

$$12. \frac{\Gamma(-a^2x^3, x^4) \Gamma(-b^2x^2, x^4)}{\Gamma(-a^2x, x^4) \Gamma(-b^2x, x^4)}$$

$$= \frac{1}{1-ab} + \frac{(a-bx)(b-ax)}{(1+x^4)(1-ab)} + \frac{(a-bx^2)(b-ax^2)}{(1+x^4)(1-ab)} + \dots$$

$$13. 1 - ax + a^2x^3 - a^3x^6 + a^4x^{10} - \dots$$

$$= \frac{1}{1+} \frac{ax}{1+} \frac{a(x^2-x)}{1+} \frac{ax^2}{1+} \frac{a(x^4-x^4)}{1+} \frac{ax^5}{1+} \dots$$

$$D_{2n} = 1 + ax^n \cdot \frac{1-x^n}{1-x} + a^2x^{2n} \cdot \frac{(1-x^n)(1-x^{n-1})}{(1-x)(1-x^2)} + \dots$$

$$+ a^3x^{3n} \cdot \frac{(1-x^n)(1-x^{n-1})(1-x^{n-2})}{(1-x)(1-x^2)(1-x^3)} + \dots$$

$$D_{2n+1} = 1 + (ax)x^n \cdot \frac{1-x^n}{1-x} + (ax)^2x^{2n} \cdot \frac{(1-x^n)(1-x^{n-1})}{(1-x)(1-x^2)} + \dots$$

$$+ (ax)^3x^{3n} \cdot \frac{(1-x^n)(1-x^{n-1})(1-x^{n-2})}{(1-x)(1-x^2)(1-x^3)} + \dots$$

$$14. \int_0^\infty \frac{\Gamma(ax, x)}{x^n \Gamma(x, x)} dx = \frac{\pi}{\sin \pi n} \cdot \frac{\Gamma(-a, n) \Gamma(-n^2, n)}{\Gamma(-n, n) \Gamma(-an^{-1}, n)}$$

$$15. \frac{1 + \frac{bx}{(1-x)(1-ax)} + \frac{b^2x^2}{(1-x)(1-x^2)(1-ax)(1-ax^2)} + \dots}{1 + \frac{bx^2}{(1-x)(1-ax)} + \frac{b^2x^6}{(1-x)(1-x^2)(1-ax)(1-ax^2)} + \dots}$$

$$= 1 + \frac{bx}{1-ax} + \frac{bx^2}{1-ax^2} + \frac{bx^3}{1-ax^4} + \dots$$

$$16. \frac{1 + \frac{ax^2}{1-ax} + \frac{a^2x^6}{(1-x)(1-x^2)} + \frac{a^3x^{12}}{(1-x)(1-x^2)(1-x^3)} + \dots}{1 + \frac{ax}{1-x} + \frac{a^2x^4}{(1-x)(1-x^2)} + \frac{a^3x^9}{(1-x)(1-x^2)(1-x^3)} + \dots}$$

$$= \frac{1}{1} + \frac{ax}{1} + \frac{ax^2}{1} + \frac{ax^3}{1} + \frac{ax^4}{1} + \dots$$

$$16. \text{ If } a = 1 + ax \cdot \frac{1-x^n}{1-x} + a^2x^2 \cdot \frac{(1-x^{n-1})(1-x^{n-2})}{(1-x)(1-x^2)} + \dots + a^nx^n \cdot \frac{(1-x^{n-1})(1-x^{n-2})\dots(1-x^2)}{(1-x)(1-x^2)(1-x^3)} + \dots$$

$$\& d = 1 + ax \cdot \frac{x-x^n}{1-x} + a^2x^2 \cdot \frac{(x-x^{n-1})(x-x^{n-2})}{(1-x)(1-x^2)} + \dots + \dots$$

$$\frac{d}{a} = 1 + \frac{ax}{1} + \frac{ax^2}{1} + \frac{ax^3}{1} + \dots + \frac{ax^n}{1}$$

$$17. \frac{\prod(x\gamma, x^2) \prod(\frac{x}{\gamma}, x^2) \prod(-x^2, x^2) \prod(-d\beta x^2, x^2)}{\prod(dx\gamma, x^2) \prod(\beta\frac{x}{\gamma}, x^2) \prod(-dx^2, x^2) \prod(-\beta x^2, x^2)}$$

$$= 1 + \left\{ xy \frac{1-d}{1-\beta x^2} + \frac{x}{y} \cdot \frac{1-\beta}{1-dx^2} \right\} +$$

$$\left\{ (xy)^2 \frac{(1-d)(x^2-d)}{(1-\beta x^2)(1-\beta x^4)} + \left(\frac{x}{y}\right)^2 \frac{(1-\beta)(d-x^2)}{(1-dx^2)(1-dx^4)} \right\} +$$

$$\left\{ (xy)^3 \frac{(1-d)(x^2-d)(x^4-d)}{(1-\beta x^2)(1-\beta x^4)(1-\beta x^6)} + \left(\frac{x}{y}\right)^3 \frac{(1-\beta)(x^2-\beta)(x^4-\beta)}{(1-dx^2)(1-dx^4)(1-dx^6)} \right\} + \dots$$

$$18. \frac{\prod(x\gamma, x^2) \prod(\frac{x}{\gamma}, x^2) \prod(-x^2, x^2) \prod(-x^2x^2, x^2)}{\prod(dx\gamma, x^2) \prod(\frac{nx}{\gamma}, x^2) \prod(-nx^2, x^2) \prod(-nx^2, x^2)}$$

$$= 1 + x(\gamma + \frac{1}{\gamma}) \cdot \frac{1-n}{1-nx^2} + x^2(\gamma^2 + \frac{1}{\gamma^2}) \cdot \frac{(1-n)(x^2-n)}{(1-nx^2)(1-nx^4)} +$$

$$x^3(\gamma^3 + \frac{1}{\gamma^3}) \cdot \frac{(1-n)(x^2-n)(x^4-n)}{(1-nx^2)(1-nx^4)(1-nx^6)} + \dots$$

$f(0) = 1 + (a+b) + ab(a^2+b^2) + (ab)^2(a^2+b^2) + (ab)^3(a^2+b^2) + \dots$

$f(a) = f(b)$ ; ii.  $f(0, a) = 2f(a, a^3)$ ; iii.  $f(-1, a) = 0$

$n$  is any integer  $f(a, b) = a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}} f\{a(ab)^n, b(ab)^n\}$   
if  $n$  is not an integer the result is approximately true.

$f(a, b) = \prod(a, ab) \prod(b, ab) \prod(-ab, ab)$

This result can be got like XVI 17 cor or as follows. —  
we see from it that if  $a(ab)^n$  or  $b(ab)^n$  be equal to  $-1$  then  $f(a, b) = 0$   
and also if  $(ab)^n = 1$ ,  $f(a, b) = 1 - (\frac{a}{b})^{\frac{n}{2}} = 0$  & hence  $f(a, b) = 0$   
Therefore  $\prod(a, ab)$ ,  $\prod(b, ab)$  &  $\prod(-ab, ab)$  are the factors of  $f(a, b)$

20. If  $\alpha\beta = \pi$ , then  $\sqrt{\alpha} f(e^{-\alpha^2+nd}, e^{-\alpha^2-nd}) = e^{\frac{\pi n}{2}} \sqrt{\beta} f(e^{-\beta^2+nd}, e^{-\beta^2-nd})$

21.  $\log \prod(a, x) = \frac{a}{1-x} - \frac{a^2}{2(1-x^2)} + \frac{a^3}{3(1-x^3)} - \frac{a^4}{4(1-x^4)} + \dots$

and consequently  $\log f(a, b) = \log \prod(ab, ab) + \frac{a+b}{1-ab} - \frac{a^2+b^2}{2(1-a^2b^2)} + \frac{a^3+b^3}{3(1-a^3b^3)} - \frac{a^4+b^4}{4(1-a^4b^4)} + \dots$

22. Let i.  $\phi(x) = f(x, x^2) = 1 + 2x + 2x^4 + 2x^9 + 2x^{16} + \dots$   
 $= \frac{1+x}{1-x} \cdot \frac{1-x^2}{1+x^2} \cdot \frac{1+x^3}{1-x^3} \cdot \frac{1-x^4}{1+x^4} \dots$

ii.  $\psi(x) = f(x, x^3) = 1 + x + x^4 + x^9 + x^{16} + \dots$   
 $= \frac{1-x^2}{1-x} \cdot \frac{1-x^4}{1-x^2} \cdot \frac{1-x^6}{1-x^3} \cdot \frac{1-x^8}{1-x^4} \dots$

iii.  $f(-x) = f(-x, -x^2) = 1 - x - x^4 + x^9 + x^{16} - x^{25} - x^{36} + \dots$   
 $= (1-x)(1-x^2)(1-x^3)(1-x^4)(1-x^5) \dots$

iv.  $\chi(x) = \prod(x, x^2) = (1+x)(1+x^2)(1+x^4)(1+x^8) \dots$



$$23. i. \log \phi(x) = 2 \left\{ \frac{x}{1+x} + \frac{x^3}{3(1+x^2)} + \frac{x^5}{5(1+x^4)} + \dots \right\}$$

$$ii. \log \psi(x) = \frac{x}{1+x} + \frac{x^4}{2(1+x^2)} + \frac{x^9}{3(1+x^3)} + \dots$$

$$iii. \log f(-x) = - \left\{ \frac{x}{1-x} + \frac{x^4}{2(1-x^2)} + \frac{x^9}{3(1-x^3)} + \dots \right\}$$

$$iv. \log \chi(x) = \frac{x}{1-x^2} - \frac{x^4}{2(1-x^4)} + \frac{x^9}{2(1-x^6)} - \dots$$

$$v. \frac{\psi(x)}{\phi(x)} = \frac{1+x^4}{1+x} \cdot \frac{1+x^8}{1+x^2} \cdot \frac{1+x^{12}}{1+x^3} \cdot \dots$$

$$ex. \frac{11}{10} \cdot \frac{1111}{1110} \cdot \frac{111111}{111110} \dots \&c = 1.1010010001000010000010000001 + \dots$$

$$24. i. \frac{f(x)}{f(-x)} = \frac{\psi(x)}{\psi(-x)} = \frac{\chi(x)}{\chi(-x)} = \sqrt{\frac{\phi(x)}{\phi(-x)}}$$

$$ii. f^2(-x) = \phi^2(-x) \psi(x) = 1 - 3x + 5x^2 - 7x^6 + 9x^{10} - \dots$$

$$iii. \chi(x) = \frac{f(x)}{f(-x)} = \frac{3\sqrt{\phi(x)}}{\psi(-x)} = \frac{\phi(x)}{f(x)} = \frac{f(-x^2)}{\psi(-x)}$$

$$iv. f^2(-x^2) = \phi^2(-x) \psi^2(x) \text{ and } \chi(x)\chi(-x) = \chi(-x^2)$$

$$25. i. \phi(x) + \phi(-x) = 2\phi(x^2)$$

$$ii. \phi(x) - \phi(-x) = 4x\psi(x^2)$$

$$iii. \phi(x)\phi(-x) = \phi^2(-x^2) \text{ and } \psi(x)\psi(-x) = \psi(x^2)\phi(-x^2)$$

$$iv. \phi(x)\psi(x^2) = \psi^2(x)$$

$$v. \phi^2(x) - \phi^2(-x) = 8x\psi^2(x^2)$$

$$vi. \phi^2(x) + \phi^2(-x) = 2\phi^2(x^2)$$

$$vii. \phi^4(x) - \phi^4(-x) = 16x\psi^4(x^2)$$

$$Cor. \text{ If } \left( \frac{1-t^2}{1+t^2} \right)^2 = \left\{ \frac{\phi(-x)}{\phi(x)} \right\}^2, \text{ then } 1-t^2 = \left\{ \frac{\phi(-x^2)}{\phi(x^2)} \right\}^2$$

26.  $x^{\frac{(m-n)}{2(m+n)}}$ .  $f(x^m, x^n)$  is a perfect, complete, pure, double series of  $\frac{1}{2}$  a degree.

Cor. i.  $\phi(x)$ ,  $\sqrt{x}\psi(x)$  &  $\sqrt{x}f(x)$  are complete series of  $\frac{1}{2}$  a degree

ii.  $\frac{\chi(x)}{\sqrt{x}}$  is a complete series of 0 degree.

i.  $\sqrt{\alpha} \psi(e^{-\alpha}) = \sqrt{\beta} \phi(e^{-\beta})$  with  $\alpha\beta = \pi$ .

ii.  $\sqrt{\alpha} \psi(e^{-2\alpha}) = \sqrt{\beta} e^{\frac{\alpha}{\beta}} \phi(e^{-\beta^2})$  with  $\alpha\beta = \pi$ .

iii.  $e^{-\frac{\alpha}{2}} \sqrt{\alpha} f(e^{-2\alpha}) = e^{-\frac{\beta}{2}} \sqrt{\beta} f(e^{-2\beta})$  with  $\alpha\beta = \pi^2$ .

iv.  $e^{-\frac{\alpha}{2}} \sqrt{\alpha} f(e^{-\alpha}) = e^{-\frac{\beta}{2}} \sqrt{\beta} f(e^{-\beta})$  with  $\alpha\beta = \pi^2$ .

v.  $e^{\frac{\alpha}{2\beta}} \chi(e^{-\alpha}) = e^{\frac{\beta}{2\alpha}} \chi(e^{-\beta})$  with  $\alpha\beta = \pi^2$ .

28.  $f(a, b^{n-1}c) f(a, b^{n-2}c) f(a, b^{n-3}c) \dots f(a, b, c)$   
 $= f(a, c) \cdot \left\{ \frac{f(b^n)}{f(b)} \right\}^n$  where  $b = ab$ .

con.  $f(-x^2, -x^3) f(-x, -x^4) = f(x) f(x^5)$   
 $f(x, -x^4) f(-x^2, -x^5) f(-x^3, -x^6) = f(x) f(-x^7)$   
 and so on.

29. If  $ab = cd$ , then

i.  $f(a, b) f(c, d) + f(a, -b) f(c, -d) = 2 f(c, cd) f(ad, cd)$

ii.  $f(a, b) f(c, d) - f(a, -b) f(c, -d) = 2a f(\frac{b}{c}, \frac{c}{b} abcd) f(\frac{b}{a}, \frac{a}{b} abcd)$

30. i.  $f(a, ab^2) f(b, a^2b) = f(a, b) \psi(ab)$

ii.  $f(a, b) + f(a, -b) = 2 f(a^3b, ab^2)$

iii.  $f(a, b) - f(a, -b) = 2a f(\frac{b}{a}, \frac{a}{b} a^2b^2)$

iv.  $f(a, b) f(a, -b) = f(-a^2, -b^2) \phi(-ab)$

v.  $f^2(a, b) + f^2(a, -b) = 2 f^2(a^2, b^2) \phi(ab)$

vi.  $f^2(a, b) - f^2(a, -b) = 4a f(\frac{b}{a}, \frac{a}{b} a^2b^2) \psi(a^2b^2)$

con. If  $ab = cd$ , then

$$f(a, b) f(c, d) f(an, \frac{b}{n}) f(cn, \frac{d}{n}) - f(-a, -b) f(-c, -d) f(-an, -\frac{b}{n}) f(-cn, -\frac{d}{n}) = 2a f(\frac{c}{a}, ad) f(\frac{d}{an}, acn) f(n, \frac{ab}{n}) \psi(ab)$$

31. If  $u_n = a \frac{n(n+1)}{2} b \frac{n(n-1)}{2}$  and  $v_n = a \frac{n(n-1)}{2} b \frac{n(n+1)}{2}$ , such that  
 $f(u_1, v_1) = 1 + (u_1 + v_1) + (u_2 + v_2) + (u_3 + v_3) + \dots$ , then  
 $f(u, v) = f(u_n, v_n) + u_1 f\left(\frac{u_{n-1}}{u_1}, \frac{u_{n+1}}{u_1}\right) + v_1 f\left(\frac{v_{n-1}}{v_1}, \frac{v_{n+1}}{v_1}\right)$   
 $+ u_2 f\left(\frac{u_{n-2}}{u_2}, \frac{u_{n+2}}{u_2}\right) + v_2 f\left(\frac{v_{n-2}}{v_2}, \frac{v_{n+2}}{v_2}\right)$   
 $+ u_3 f\left(\frac{u_{n-3}}{u_3}, \frac{u_{n+3}}{u_3}\right) + v_3 f\left(\frac{v_{n-3}}{v_3}, \frac{v_{n+3}}{v_3}\right)$   
 $+ \dots \dots \dots + \dots \dots \dots$

e.g. i.  $\phi(x) = \phi(x^2) + 2x f(x^3, x^{11}) = \phi(x^{2^5}) + 2x f(x^{15}, x^{11})$   
 $+ 2x^4 f(x^5, x^{61}) = \dots$

ii.  $\psi(x) = f(x^2, x^6) + x \psi(x^3) = f(x^6, x^{10}) + x f(x^2, x^{14})$   
 $= f(x^{10}, x^{15}) + x f(x^5, x^{20}) + x^3 \psi(x^{2^5})$   
 $= f(x^{15}, x^{24}) + x \psi(x^3) + x^3 f(x^3, x^{37}) = \dots \dots \dots$

ex. i.  $\frac{\phi^2(x)}{\phi^2(x)} + \frac{\phi^2(y)}{\phi^2(y)} + \frac{\phi^2(z)}{\phi^2(z)} + \frac{\phi^2(x)\phi^2(y)\phi^2(z)}{\phi^2(x)\phi^2(y)\phi^2(z)}$   
 $= 4 \cdot \frac{\phi^2(x^4)\phi^2(y^4)\phi^2(z^4)}{\phi^2(x^4)\phi^2(y^4)\phi^2(z^4)} + 256xyz \frac{\psi^2(x^4)\psi^2(y^4)\psi^2(z^4)}{\phi^2(x)\phi^2(y)\phi^2(z)}$

ii.  $\frac{1}{\phi(x^4)} = \frac{1}{\phi(x) \pm \phi(x^4)} + \frac{1}{\phi(x) \pm \phi(x^4)}$  and  
 $\frac{1}{\phi(x^4)} = \frac{1}{\phi(x^4) \pm \phi(x)} + \frac{1}{\phi(x^4) \pm \phi(x)}$

iii. The coeff. of  $x^n$  in the expansion of  $\frac{x}{1-x} \psi(x^2)$  is the nearest integer to  $\sqrt{n}$ .

iv.  $\phi(-x) + \phi(x^2) = 2 \cdot \frac{f^2(x^3, x^5)}{\psi(x)}$  and  
 $\phi(-x) - \phi(x^2) = -2x \cdot \frac{f^2(x, x^7)}{\psi(x)}$

v.  $f(x, x^5) = \psi(-x^2) \chi(x)$

32. i.  $\frac{\phi'(x)}{\psi(x)} - \frac{\psi'(x)}{\psi(x)} = 1 - \frac{\phi'(x)}{\psi(x)}$

ii.  $\frac{\phi'(x)}{\psi(x)} - 2x \frac{\psi'(x)}{\psi(x^2)} = 1 - \frac{\phi'(x)}{\psi(x)}$

iii.  $\frac{\phi'(x)}{\phi(x)} + \frac{\phi'(-x)}{\phi(-x)} = \frac{\phi'(x) - \phi'(-x)}{4x}$

iv.  $\frac{\phi'(x)}{\phi(x)} - \frac{\phi'(-x)}{\phi(-x)} = -4x \frac{\phi'(x^2)}{\phi(-x^2)}$

33. i.  $\log(1 + 2x \cos \theta + 2x^4 \cos 2\theta + 2x^9 \cos 3\theta + \dots)$

$-\log f(x^2) = 2 \left\{ \frac{x}{1-x^2} \cos \theta - \frac{x^2}{2(1-x^4)} \cos 2\theta + \frac{x^3}{3(1-x^6)} \cos 3\theta - \dots \right\}$

ii.  $\frac{1}{2} \log \frac{\sin x - x \sin 3x + x^3 \sin 5x - 2^6 \sin 7x + \dots}{\sin x (1 - 3x + 5x^3 - 7x^6 + 9x^{10} - \dots)}$

$= \frac{x \sin^2 x}{1(1-x)} + \frac{x^2 \sin^2 2x}{2(1-x^2)} + \frac{x^3 \sin^2 3x}{3(1-x^3)} + \dots$

iii.  $1 + \frac{4x \cos x}{1+x^2} + \frac{4x^2 \cos 2x}{1+x^4} + \frac{4x^3 \cos 3x}{1+x^6} + \dots$

$= \phi^2(x^2) \frac{1 + 2x \cos x + 2x^2 \cos 2x + 2x^3 \cos 3x + \dots}{1 - 2x \cos x + 2x^2 \cos 2x - 2x^3 \cos 3x + \dots}$

Cor.  $\frac{f(a, b)}{f(-a, -b)} \phi^2(-ab) = 1 + 2 \left\{ \frac{a+b}{1+ab} + \frac{a^2+b^2}{1+a^2b^2} + \frac{a^3+b^3}{1+a^3b^3} + \dots \right\}$

34. i.  $\log \frac{\phi^2(x)}{1 + 4 \cos x \left( \frac{x \cos x}{1-x} - \frac{x^2 \cos 2x}{1-x^2} + \frac{x^3 \cos 3x}{1-x^3} - \dots \right)}$

$= 4 \left\{ \frac{x \sin^2 x}{1(1+x)} - \frac{x^2 \sin^2 2x}{2(1+x^2)} + \frac{x^3 \sin^2 3x}{3(1+x^3)} - \dots \right\}$

ii.  $\frac{1}{8} \log \frac{\phi^2(x)}{1 + \frac{2x \cos x}{1+x^2} + \frac{4x^2 \cos 2x}{1+x^4} + \dots}$

$= \frac{x \sin^2 x}{1(1-x)} + \frac{x^3 \sin^2 3x}{3(1-x^6)} + \frac{x^5 \sin^2 5x}{5(1-x^{10})} + \dots$

195

$$\text{Ex. i. } \frac{1}{4} \log \frac{\sin 2n - x \sin 4n + x^5 \sin 8n - x^8 \sin 16n + \dots}{\sin n (1 - 2x + 4x^5 - 5x^8 + 7x^{16} - \dots)}$$

$$= \frac{x}{1+x} \sin^2 n + \frac{x^2}{2(1+x^2)} \sin^2 2n + \frac{x^3}{3(1+x^3)} \sin^2 3n + \dots$$

$$+ \frac{x^4}{1-x^4} \sin^2 2n + \frac{x^8}{2(1-x^8)} \sin^2 4n + \frac{x^{12}}{3(1-x^{12})} \sin^2 6n + \dots$$

$$\text{ii. } \frac{1}{4} \log \frac{\sin n - x \sin 5n + x^4 \sin 7n - x^5 \sin 11n + \dots}{\sin n (1 - 5x + 7x^4 - 11x^5 + 13x^7 - \dots)}$$

$$= \frac{x \sin^2 n}{1-x} + \frac{x^2 \sin^2 2n}{2(1-x^2)} + \frac{x^3 \sin^2 3n}{3(1-x^3)} + \dots$$

$$+ \frac{x \sin^2 2n}{1+x} + \frac{x^2 \sin^2 4n}{2(1+x^2)} + \frac{x^3 \sin^2 6n}{3(1+x^3)} + \dots$$

$$35. \text{ i. If } P_n = \frac{B_n}{2^n} \cos \frac{\pi n}{2} + \frac{1^{n+1} x}{1-x} + \frac{2^{n+1} x^2}{1-x^2} + \frac{3^{n+1} x^3}{1-x^3} + \dots$$

$$\text{and } Q_n = \frac{1}{n+1} \cdot \frac{1^{n+1} - 3^{n+1} x + 5^{n+1} x^3 - 7^{n+1} x^6 + \dots}{1 - 3x + 5x^3 - 7x^6 + \dots}$$

$$\text{then } \frac{1}{2} Q_n = -2^n P_n - \frac{(n-1)(n-2)}{1!} 2^{n-2} P_{n-2} Q_2 - \frac{(n-1)(n-2)(n-3)(n-4)}{4!} 2^{n-4} P_{n-4} Q_4 - \dots$$

$$\text{ii. If } P_n = \frac{B_n}{2^n} (2^n - 1) \cos \frac{\pi n}{2} + \frac{1^{n+1} x}{1+x} + \frac{2^{n+1} x^2}{1+x^2} + \frac{3^{n+1} x^3}{1+x^3} - \dots$$

$$\text{and } Q_n = \frac{\frac{1}{2} E_{n+1} \cos \frac{\pi n}{2} + \frac{1^{n+1} x}{1-x} - \frac{3^{n+1} x^3}{1-x^3} + \frac{5^{n+1} x^5}{1-x^5} - \dots}{\frac{1}{2} E_1 + \frac{x}{1-x} - \frac{3x^3}{1-x^3} + \frac{5x^5}{1-x^5} - \dots}$$

$$\text{then } \frac{1}{2} Q_n = 2^n P_n - \frac{(n-1)(n-2)}{1!} 2^{n-2} P_{n-2} Q_2 + \frac{(n-1)(n-2)(n-3)(n-4)}{4!} 2^{n-4} P_{n-4} Q_4 - \dots$$

N.B. Thus the series  $1 - 3^{n+1} x + 5^{n+1} x^3 - 7^{n+1} x^6 + \dots$  can be expressed in terms of  $L$ ,  $M$  and  $N$ .

$$e_2. \frac{1^3 - 3^3x + 5^3x^2 - 7^3x^3 + \dots + \infty}{1x + 5x^3 - 7x^6 + \dots} = L.$$

$$ii. \frac{1^5 - 3^5x + 5^5x^2 - 7^5x^3 + \dots + \infty}{1x + 5x^3 - 7x^6 + \dots} = \frac{5L^2 - 2M}{3}$$

$$iii. \frac{1^7 - 3^7x + 5^7x^2 - 7^7x^3 + \dots + \infty}{1x + 5x^3 - 7x^6 + \dots} = \frac{35L^3 - 42LM + 16N}{9}$$

36. If  $\frac{a^6}{c^2d} = p$ , then

$$i. \frac{1}{2} \{ f(a, b) f(c, d) + f(a, -b) f(c, -d) \}$$

$$= f(ac, bd) + ad f(acp, \frac{bd}{p}) + bc f(bdp, \frac{ac}{p})$$

$$+ (ad)^3 bc f(acp^2, \frac{bd}{p^2}) + (bc)^3 ad f(bdp^2, \frac{ac}{p^2})$$

$$+ (ad)^6 (bc)^3 f(acp^3, \frac{bd}{p^3}) + (bc)^6 (ad)^3 f(bdp^3, \frac{ac}{p^3})$$

$$+ \dots \dots \dots + \dots \dots \dots$$

$$ii. \frac{1}{2} \{ f(a, b) f(c, d) - f(a, -b) f(c, -d) \}$$

$$= a f(\frac{c}{a}, \frac{a}{c} abcd) + d f(\frac{b}{d}, \frac{d}{b} abcd)$$

$$+ a^3 bc f(\frac{c}{ap}, \frac{ap}{c} abcd) + d^3 bc f(\frac{b}{d}, \frac{d}{bp} abcd)$$

$$+ a^5 d (bc)^3 f(\frac{c}{ap^2}, \frac{ap^2}{c} abcd) + ad^5 (bc)^3 f(\frac{b}{d}, \frac{d}{bp^2} abcd)$$

$$+ \dots \dots \dots + \dots \dots \dots$$

$$37. i. \frac{1}{2} \{ \phi(a) \phi(b) + \phi(-a) \phi(-b) \}$$

$$= \phi(ab) + 2ab f(\frac{a^3}{b}, \frac{b^3}{a}) + 2(ab)^2 f(\frac{a^5}{b^2}, \frac{b^5}{a^2}) +$$

$$2(ab)^3 f(\frac{a^7}{b^3}, \frac{b^7}{a^3}) + \dots$$

$$ii. \frac{1}{2} \{ \phi(a) \phi(b) - \phi(-a) \phi(-b) \} = 2ab f(\frac{b}{a}, a^3b) +$$

$$2a^3b f(\frac{b^3}{a^2}, \frac{a^5}{b}) + 2a^5b^2 f(\frac{b^5}{a^3}, \frac{a^7}{b^2}) + \dots$$

$$\text{iii. } \psi(a)\psi(b) = \psi(ab) + a f\left(\frac{b}{a}, a^2\right) + a^2 b f\left(\frac{b^2}{a^2}, \frac{a^3}{b}\right) + a^6 b^3 f\left(\frac{b^3}{a^3}, \frac{a^6}{b^2}\right) + \dots$$

$$\text{Cor. i. } \psi(x^2)\psi(x^{13}) - \psi(x^{26})\psi(x^{13}) = x^{13} \{ \psi(x)\psi(x^{27}) + \psi(x^2)\psi(x^{27}) \}$$

$$\text{ii } \psi(x^5)\psi(x^{11}) - \psi(x^{55})\psi(x^{11}) = x^{55} \{ \psi(x)\psi(x^{55}) + \psi(x^2)\psi(x^{55}) \}$$

$$\text{iii } \psi(x^7)\psi(x^9) - \psi(x^{63})\psi(x^9) = x^{63} \{ \psi(x)\psi(x^{63}) - \psi(x^2)\psi(x^{63}) \}$$

$$\text{ex. } \psi(x)\psi(x^4) - \psi(x^4)\psi(x^4) = 2x \cdot \frac{\phi(x^6)\phi(-x^{12})}{\chi(-x^2)\chi(-x^6)} + \frac{1}{2}x^{15}\psi(x^6)\psi(x^{12})$$

$$38. \text{ i. } \frac{f(x^5)}{f(-x, -x^4)} = 1 + \frac{x}{1-x} + \frac{x^5}{(1-x)(1-x^4)} + \frac{x^9}{(1-x)(1-x^4)(1-x^2)} + \dots$$

$$\text{ii. } \frac{f(-x^5)}{f(x, -x^4)} = 1 + \frac{x^2}{1-x} + \frac{x^6}{(1-x)(1-x^4)} + \frac{x^{12}}{(1-x)(1-x^4)(1-x^2)} + \dots$$

$$\text{iii. } \frac{f(x, -x^4)}{f(-x^5, -x^4)} = \frac{1}{1} + \frac{x}{1} + \frac{x^2}{1} + \frac{x^3}{1} + \frac{x^4}{1} + \frac{x^5}{1} + \dots$$

$$\text{iv. } f^2(x, -x^4) - \sqrt{x^2} f^2(x, -x^4) = f(-x) \{ f(\sqrt{x}) + \sqrt{x} f(x^5) \}$$

$$39. \text{ i. } \left\{ \frac{\sqrt{5}+1}{2} + \frac{e^{-3\alpha}}{1+} \frac{e^{-2\alpha}}{1+} \frac{e^{-4\alpha}}{1+} \frac{e^{-6\alpha}}{1+} \frac{e^{-8\alpha}}{1+} \dots \right\} \times$$

$$\left\{ \frac{\sqrt{5}+1}{2} + \frac{e^{-2\beta}}{1+} \frac{e^{-2\beta}}{1+} \frac{e^{-4\beta}}{1+} \frac{e^{-6\beta}}{1+} \frac{e^{-8\beta}}{1+} \dots \right\} = \frac{5+\sqrt{5}}{2}$$

$$\text{ii } \left\{ \frac{\sqrt{5}-1}{2} + \frac{e^{-\alpha}}{1-} \frac{e^{-\alpha}}{1+} \frac{e^{-2\alpha}}{1-} \frac{e^{-3\alpha}}{1+} \frac{e^{-4\alpha}}{1-} \dots \right\} \times$$

$$\left\{ \frac{\sqrt{5}-1}{2} + \frac{e^{-\beta}}{1-} \frac{e^{-\beta}}{1+} \frac{e^{-2\beta}}{1-} \frac{e^{-2\beta}}{1+} \frac{e^{-4\beta}}{1-} \dots \right\} = \frac{5-\sqrt{5}}{2}$$

with  $\alpha\beta = \pi^2$  in both the cases.

$$\text{Cor. i. } \frac{e^{-\frac{\pi}{5}}}{1-} \frac{e^{-\pi}}{1+} \frac{e^{-2\pi}}{1-} \frac{e^{-3\pi}}{1+} \frac{e^{-4\pi}}{1-} \dots = \sqrt{\frac{5-\sqrt{5}}{2}} - \frac{\sqrt{5}-1}{2}$$

$$\text{ii. } \frac{e^{-\frac{2\pi}{5}}}{1+} \frac{e^{-2\pi}}{1+} \frac{e^{-4\pi}}{1+} \frac{e^{-6\pi}}{1+} \frac{e^{-8\pi}}{1+} \dots = \sqrt{\frac{5+\sqrt{5}}{2}} - \frac{\sqrt{5}+1}{2}$$

$$1. \int_0^{\pi} \frac{\cos\{(1-2n) \operatorname{Sin}^{-1}(\sqrt{x} \operatorname{Sin} \phi)\}}{\sqrt{1-x \operatorname{Sin}^2 \phi}} d\phi$$

$$= \frac{\pi}{2} \left\{ 1 + \frac{n(1-n)x}{(1-x)^2} + \frac{n(n+1)(1-n)(2-n)x^2}{(1-x)^4} + \dots \right\}$$

Case i:  $\int_0^{\frac{\pi}{2}} \frac{\cos\{(1-2n) \operatorname{Sin}^{-1}(\frac{\operatorname{Sin} \phi}{\sqrt{2}})\}}{\sqrt{1-\frac{1}{2} \operatorname{Sin}^2 \phi}} d\phi = \frac{\pi}{2} \cdot \frac{\sqrt{\pi}}{\Gamma(\frac{n-1}{2}) \Gamma(\frac{n+1}{2})}$

ii. If  $\int_0^{\pi} \frac{\cos\{(1-2n) \operatorname{Sin}^{-1}(\sqrt{x} \operatorname{Sin} \phi)\}}{\sqrt{1-x \operatorname{Sin}^2 \phi}} d\phi = U_x$ , then

$$e^{-\pi} \frac{U_{1-x} \cos \pi n}{U_x} = e^{-\frac{1}{2}\pi} \left\{ x + x^2(1-2n-n^2) + x^3(1-7 \cdot \frac{n-n^2}{2} + 13 \cdot \frac{n-n^2}{2} x^2) + \dots \right\}$$

2. Let  $F(x) = e^{-\pi} \frac{1 + (\frac{1}{2})^2(1-x) + (\frac{1}{2})^4(1-x)^2 + \dots}{1 + (\frac{1}{2})^2 x + (\frac{1}{2})^4 x^2 + \dots}$ , then

i.  $F(x) = \frac{x}{10} \cdot e^{-\frac{2}{10} \pi} \frac{(\frac{1}{2})^2(-\frac{1}{2})x + (\frac{1}{2})^4(\frac{1}{2} + \frac{1}{2})x^2 + \dots}{1 + (\frac{1}{2})^2 x + (\frac{1}{2})^4 x^2 + (\frac{1}{2})^6 \frac{1}{2} x^3 + \dots}$

ii.  $F(1-e^{-x}) = \frac{x}{10 + \sqrt{36+x^2}}$  very nearly.

iii.  $\log F(x) \log F(1-x) = \pi^2$ .

iv.  $F(1-x) + F(1-\frac{1}{x}) = 0$

v.  $F\left\{\frac{2x}{(1+x)^2}\right\} = \sqrt{F(x^2)}$

N.B. Suppose we know the expansion of  $F\left(\frac{2x}{1+x}\right)$  to  $n$  terms. changing  $x$  to  $\frac{x^2}{x+1}$  for  $x$  we have the expansion of  $F(x^2)$  to  $n$  terms, i.e. that of  $\left\{F\left(\frac{2x}{1+x}\right)\right\}^2$  to  $2n$  terms. Extracting the square root and expanding the result in ascending



202  
 ing powers of  $\frac{2x}{1+x}$  we can find the expansion of  $F \frac{2x}{1+x}$  to  $2n$  terms.

vi.  $8F\left(\frac{2x}{1+x}\right) = x + \frac{5}{16}x^3 + \frac{369}{2048}x^5 + \frac{4097}{32768}x^7 + \frac{1594875}{16777216}x^9 + \dots$

vii.  $2F(1-e^{-8x}) = x - \frac{x^3}{3} + \frac{31}{120}x^5 - \frac{661}{2520}x^7 + \frac{319677}{725760}x^9 - \dots$

viii.  $F(0) = 0; F\left(\frac{1}{2}\right) = e^{-\pi}; F(1) = 1; F(\sqrt{2}-1) = e^{-\pi\sqrt{2}}; F(\sqrt{2}-1)^2 = e^{-2\pi}$

ix.  $2F\left(1 - e^{-\frac{8x}{1-x^2}}\right) = x + \frac{2}{3}x^3 + \frac{31}{120}x^5 + \frac{37}{1260}x^7 + \frac{5981}{725760}x^9 + \dots$

3.  $\phi^2(x) = 1 + \left(\frac{x}{2}\right)^2 \left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\} + \left(\frac{1.3}{2.4}\right)^2 \left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\}^2 + \dots$

N.B. We know that  $1 + \left(\frac{x}{2}\right)^2 \left\{1 - \left(\frac{x}{1+x}\right)^2\right\} + \left(\frac{1.3}{2.4}\right)^2 \left\{1 - \left(\frac{x}{1+x}\right)^2\right\}^2 + \dots$   
 $= (1+x) \left\{1 + \left(\frac{x}{2}\right)^2 + \left(\frac{1.3}{2.4}\right)^2 x^2 + \dots\right\}$  and also that

$1 + \left(\frac{x}{2}\right)^2 \left(\frac{x}{1+x}\right)^2 + \left(\frac{1.3}{2.4}\right)^2 \left(\frac{x}{1+x}\right)^4 + \dots = \left(\frac{1+x}{2}\right) \left\{1 + \left(\frac{x}{2}\right)^2 + \dots\right\}$

Hence by XVI 75 cor. we have  $1 + \left(\frac{x}{2}\right)^2 \left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\} + \dots$   
 $= \frac{\phi^6(x)}{\phi^6(x)} \left\{1 + \left(\frac{x}{2}\right)^2 \left[1 - \frac{\phi^6(x)}{\phi^6(x)}\right]\right\} + \dots$

Consequently  $1 + \left(\frac{x}{2}\right)^2 \left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\} + \dots$   
 $= \frac{\phi^6(x)}{\phi^6(x)} \left\{1 + \left(\frac{x}{2}\right)^2 \left[1 - \frac{\phi^6(x)}{\phi^6(x)}\right]\right\} + \dots$

By making  $n$  infinite the above result is got.

In a similar manner we can show that

$1 + \left(\frac{x}{2}\right)^2 \left\{\frac{\phi^6(x)}{\phi^6(x)}\right\} + \left(\frac{1.3}{2.4}\right)^2 \left\{\frac{\phi^6(x)}{\phi^6(x)}\right\}^2 + \dots = \frac{\phi^6(x)}{n \phi^6(x)} \left\{1 + \left(\frac{x}{2}\right)^2 \frac{\phi^6(x)}{\phi^6(x)} + \dots\right\}$

from which we have

i.  $F\left\{\frac{\phi^6(x)}{\phi^6(x)}\right\} = \sqrt[n]{F\left\{\frac{\phi^6(x)}{\phi^6(x)}\right\}}$  and similarly

ii.  $F\left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\} = \sqrt[n]{F\left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\}}$  hence we have

5.  $F\left\{1 - \frac{\phi^6(x)}{\phi^6(x)}\right\} = x$

$$6. \sqrt{1+x} = 1 + \left(\frac{1}{2}\right)x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 2 \cdot 2}\right)x^3 + \dots \text{ i.e. } 203$$

$$\text{If } \sqrt{1+x} = 1 + \left(\frac{1}{2}\right)x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)x^2 + \dots \text{ and}$$

$$\int \frac{1 + \left(\frac{1}{2}\right)(1-x) + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)(1-x)^2 + \dots}{1 + \left(\frac{1}{2}\right)x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)x^2 + \dots} \text{ then}$$

$$1 + 2e^{-y} + 2e^{-4y} + 2e^{-9y} + 2e^{-16y} + \dots = \sqrt{2}$$

$$\text{Gen. If } \int d\beta = \pi, \text{ then } \left\{ \sqrt{\alpha} \left\{ \frac{1}{2} + e^{-\alpha^2} + e^{-4\alpha^2} + \dots \right\} \right.$$

$$\text{ex. i. } 1 + 2e^{-\pi} + 2e^{-4\pi} + 2e^{-9\pi} + \dots = \frac{\sqrt{\pi}}{\sqrt{\frac{1}{2}}}$$

$$\text{ii. } 1 + 2e^{-\pi/2} + 2e^{-4\pi/2} + 2e^{-9\pi/2} + \dots = \frac{\sqrt{\frac{1}{2}}}{\sqrt{\pi \frac{1}{2}}}$$

$$\text{iii. } 1 + 2e^{-2\pi} + 2e^{-8\pi} + 2e^{-18\pi} + \dots = \frac{\sqrt{\pi}}{2 \sqrt{\frac{1}{2}}} \sqrt{2 + \sqrt{2}}$$

$$\text{iv. } \frac{\pi - \frac{1}{2}}{e^{\pi}} + \frac{4\pi - \frac{1}{2}}{e^{4\pi}} + \frac{9\pi - \frac{1}{2}}{e^{9\pi}} + \dots = \frac{1}{8}$$

$$7. \text{ i. If } \frac{\sin d}{\sin \beta} = \sqrt{x}, \text{ then } \int_0^{\alpha} \frac{d\phi}{\sqrt{x - \sin^2 \phi}} = \int_0^{\beta} \frac{d\phi}{\sqrt{1 - x \sin^2 \phi}}$$

$$\text{ii. If } \frac{\cos d}{\cos \beta} = \sqrt{1-x}, \text{ then } \int_0^{\alpha} \frac{d\phi}{\sqrt{1-x \cos^2 \phi}} = \int_0^{\beta} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}$$

$$\text{iii. If } \frac{\tan d}{\tan \beta} = \sqrt{1-a}, \text{ then } \int_0^{\beta} \frac{d\phi}{\sqrt{(1-a \sin^2 \phi)(1-b \sin^2 \phi)}} \\ = \frac{1}{\sqrt{1-b}} \int_0^{\alpha} \frac{d\phi}{\sqrt{1 - \frac{a-b}{1-b} \sin^2 \phi}}$$

$$\text{iv. If } \frac{\tan d}{\tan \beta} = \sqrt{1+x}, \text{ then } \int_0^{\alpha} \frac{d\phi}{\sqrt{1+x \cos^2 \phi}} = \int_0^{\beta} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}$$

$$\text{v. If } \cot d \cot \beta = \sqrt{1-x}, \text{ then } \int_0^{\alpha} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} + \int_0^{\beta} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} \\ = \frac{\pi}{2} \left\{ 1 + \left(\frac{1}{2}\right)x + \left(\frac{1 \cdot 3}{2 \cdot 2}\right)x^2 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 2 \cdot 2}\right)x^3 + \dots \right\}$$

vi. If  $\cot \alpha \tan \frac{\beta}{2} = \sqrt{1-x \sin^2 \alpha}$ , then

$$2 \int_0^{\alpha} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} = \int_0^{\beta} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}$$

vii. If  $\alpha = \log \tan \left( \frac{\pi}{4} + \frac{\beta}{2} \right)$ , then

$$\int_0^{\alpha} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} = i \int_0^{\beta} \frac{d\phi}{\sqrt{1-(1-x) \sin^2 \phi}}$$

viii. If  $\int_0^{\alpha} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} + \int_0^{\beta} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} = \int_0^{\gamma} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}$ , then

$$\tan \frac{\gamma}{2} = \frac{\sin \alpha \sqrt{1-x \sin^2 \beta} + \sin \beta \sqrt{1-x \sin^2 \alpha}}{\cos \alpha + \cos \beta}$$

$$\tan^{-1}(\tan \alpha \sqrt{1-x \sin^2 \beta}) + \tan^{-1}(\tan \beta \sqrt{1-x \sin^2 \alpha}) = \gamma$$

$$\text{or } \cot \alpha \cot \beta = \frac{\cos \gamma}{\sin \alpha \sin \beta} + \sqrt{1-x \sin^2 \gamma} \text{ or}$$

$$\frac{\sqrt{x}}{2} = \frac{\sqrt{\sin \alpha \sin(\alpha-\beta) \sin(\alpha-\gamma) \sin \beta}}{\sin \alpha \sin \beta \sin \gamma}, \text{ where } \gamma = \alpha + \beta + \dots$$

ix.  $\frac{\pi}{2} \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1+x \sin \phi}} = \int_0^{\frac{\pi}{2}} \frac{\cos^{-1}(x \sin^2 \phi)}{\sqrt{1-x^2 \sin^4 \phi}} d\phi$

x.  $\int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{d\theta d\phi}{\sqrt{(1-x \sin^2 \theta)(1-x \sin^2 \theta \sin^2 \phi)}} = \left\{ \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} \right\}^2$

xi.  $\int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \frac{x \sin \phi d\theta d\phi}{\sqrt{1-x^2 \sin^2 \phi} \sqrt{1-x^2 \sin^2 \theta \sin^2 \phi}}$   
 $= \int_0^{\frac{\pi}{2}} \int_0^{\sin^2 \alpha} \frac{d\theta}{\sqrt{1-x^2 \sin^2 \phi - \sin^2 \theta \cos^2 \phi}} d\phi$   
 $= \frac{1}{2} \left\{ \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-\frac{1+x}{2} \sin^2 \phi}} \right\}^2 - \frac{1}{2} \left\{ \int_0^{\frac{\pi}{2}} \frac{d\phi}{\sqrt{1-\frac{1-x}{2} \sin^2 \phi}} \right\}^2$

If  $\frac{\sin \beta}{\sin \alpha} = \frac{1+x}{1+x \sin^2 \alpha}$ , then

(1+x)  $\int_0^\alpha \frac{d\phi}{\sqrt{1-x^2 \sin^2 \phi}} = \int_0^\beta \frac{d\phi}{\sqrt{1-\frac{4x}{(1+x)} \sin^2 \phi}}$

xiii. If  $x \sin \alpha = \sin(2\beta - \alpha)$ , then

(1+x)  $\int_0^\alpha \frac{d\phi}{\sqrt{1-x^2 \sin^2 \phi}} = 2 \int_0^\beta \frac{d\phi}{\sqrt{1-\frac{4x}{(1+x)} \sin^2 \phi}}$

- i.  $\phi^2(x) = 1 + 4(\frac{x}{1-x} - \frac{x^3}{1-x^3} + \frac{x^5}{1-x^5} - \dots)$
- ii.  $\phi^4(x) = 1 + 8(\frac{x}{1-x} + \frac{2x^2}{1+x^2} + \frac{3x^3}{1-x^3} + \frac{4x^4}{1+x^4} + \dots)$
- iii.  $\phi(x)\phi(x^2) = 1 + \frac{2x}{1-x} + \frac{2x^2}{1+x^2} - \frac{2x^5}{1-x^5} - \frac{2x^7}{1-x^7} + \dots$
- iv.  $\phi(x)\phi(x^3) = 1 + \frac{2x}{1-x} - \frac{2x^6}{1+x^2} + \frac{2x^6}{1+x^4} - \frac{2x^5}{1-x^5} + \frac{2x^7}{1-x^7} - \dots$
- v.  $\phi^2(x) = 1 - \frac{4x}{1+x} + \frac{4x^2}{1+x^2} - \frac{4x^6}{1+x^2} + \frac{4x^{10}}{1+x^4} - \dots$
- vi.  $\psi(x)\phi(x^2) = \frac{1+x}{1-x} - x \frac{1+x^2}{1-x^2} + x^2 \frac{1+x^4}{1-x^4} - x^6 \frac{1+x^7}{1-x^7} + \dots$
- vii.  $\psi^2(x) = \frac{1+x}{1-x} - x^2 \frac{1+x^2}{1-x^2} + x^6 \frac{1+x^4}{1-x^4} - x^{12} \frac{1+x^7}{1-x^7} + \dots$
- viii.  $\frac{x}{1-x} + \frac{2x^2}{1-x^2} + \frac{3x^3}{1-x^3} + \frac{4x^4}{1-x^4} + \dots$   
 $= x \frac{1+x}{(1-x)^2} - x^3 \frac{1+x^2}{(1-x^2)^2} + x^6 \frac{1+x^3}{(1-x^3)^2} - x^{10} \frac{1+x^4}{(1-x^4)^2} + \dots$
- ix.  $\phi^4(-x)f(-x) = 1 - 6x + 7x^2 - 11x^5 + 13x^7 - \dots$
- x.  $\psi(x^2)f^2(x) = 1 - 2x + 4x^5 - 5x^8 + 7x^{11} - \dots$
- xi.  $f(x)f(-x^2) = \phi(x)\psi(x)$
- xii.  $\frac{f(x)}{f(-x^2)} = \frac{\phi(-x^2)}{\psi(x)}$

ex.  $\psi(x^2)f^2(x) + 2x\psi(x^2)f^2(-x^2) = \phi^2(-x^2)f^2(-x^2)$

9. Let  $y = \pi \frac{1 + (1-x)^2 + (4-x)^2(1-x)^2 + \dots}{1 + (1-x)^2 + (1-x)^2 x^2 + \dots}$

and  $z = 1 + (1-x)^2 + (1-x)^2 x^2 + \dots$  such that  $e^{-y} = F(x)$

i.  $\frac{dy}{dx} = -\frac{1}{x(1-x)z^2}$     ii.  $\frac{dz}{dx} = \frac{\int z dx}{4x(1-x)}$

iii.  $z \int \int x^n (1-x) z^3 (dy)^2 = \frac{x^n}{n!} \left\{ 1 + \left(\frac{n+1}{n+1}\right)^2 x + \left(\frac{n+1}{n+1} \cdot \frac{n+1}{n+1}\right)^2 x^2 + \dots \right\}$

iv.  $1 - 24 \left( \frac{1}{e^{12}} + \frac{2}{e^{42}} + \frac{3}{e^{62}} + \frac{4}{e^{82}} + \dots \right)$

$= (1-2x)z^2 + 6x(1-x)z \frac{dz}{dx}$

ex.  $16'' e^{-11y} = x^{11} + \frac{11}{2} x^{12} + \frac{111}{64} x^{13} + \frac{11111}{2688} x^{14} + \dots$

10. i.  $\phi(e^{-y}) = \sqrt{z}$     ii.  $\phi(-e^{-y}) = \sqrt{z} \sqrt{1-x}$

iii.  $\phi(-e^{-2y}) = \sqrt{z} \sqrt{1-x}$     iv.  $\phi(e^{-2y}) = \sqrt{z} \sqrt{\frac{1+\sqrt{1-x}}{2}}$

v.  $\phi(e^{-4y}) = \sqrt{z} \cdot \frac{1+\sqrt{1-x}}{2}$

vi.  $\phi(e^{-\frac{y}{2}}) = \sqrt{z} \sqrt{1+\sqrt{x}}$     vii.  $\phi(-e^{-\frac{y}{2}}) = \sqrt{z} \sqrt{1-x}$

viii.  $\phi(e^{-\frac{y}{2}}) = \sqrt{z} (1+\sqrt{x})$     ix.  $\phi(e^{-\frac{y}{2}}) = \sqrt{z} (1-\sqrt{x})$

11. i.  $\psi(e^{-y}) = \sqrt{\frac{z}{x}} \sqrt{x e^y}$     ii.  $\psi(-e^{-y}) = \sqrt{\frac{z}{x}} \sqrt{x(1-x)e^y}$

iii.  $\psi(e^{-2y}) = \frac{1}{2} \sqrt{z} \sqrt{x e^y}$     iv.  $\psi(e^{-4y}) = \frac{1}{2} \sqrt{\frac{z}{x}} \sqrt{(1-\sqrt{1-x})e^y}$

v.  $\psi(e^{-8y}) = \frac{\sqrt{z}}{4} (1-\sqrt{1-x})e^y$

vi.  $\psi(e^{-\frac{y}{2}}) = \sqrt{z} \sqrt{\frac{1+\sqrt{x}}{2}} \sqrt{x e^y}$

vii.  $\psi(-e^{-\frac{y}{2}}) = \sqrt{z} \sqrt{\frac{1-\sqrt{x}}{2}} \sqrt{x e^y}$

viii.  $\psi(e^{-\frac{y}{2}}) = \sqrt{z} \sqrt{1+\sqrt{x}} \sqrt{\frac{1+\sqrt{x}}{2}} \sqrt{x e^y}$

ix.  $\psi(-e^{-\frac{y}{2}}) = \sqrt{z} \sqrt{1-\sqrt{x}} \sqrt{\frac{1-\sqrt{x}}{2}} \sqrt{x e^y}$

12. i.  $f(e^{-y}) = \frac{\sqrt{z}}{\sqrt{2}} \sqrt{x(1-x)e^y}$     ii.  $f(-e^{-y}) =$

$\frac{\sqrt{z}}{\sqrt{2}} \sqrt{1-x} \sqrt{x e^y}$

20)

$$\text{iii } f(x, y) = \frac{\sqrt{z}}{\sqrt{z}} \sqrt{x a - x} e^y \cdot \text{iv } f(x, e^{-4y}) = \frac{\sqrt{z}}{\sqrt{z}} \sqrt{1-x} \sqrt{x e^y}$$

$$\text{v. } \chi(x, y) = \frac{\sqrt{z}}{\sqrt{x(1-x)} e^y} \quad \text{vi } \chi(x, e^{-y}) = \frac{\sqrt{z} \sqrt{1-x}}{\sqrt{x e^y}}$$

$$\text{vii } \chi(x, e^{-2y}) = \frac{\sqrt{z} \sqrt{1-x}}{\sqrt{x e^y}}$$

$$\text{13. i. } 1 + 240 \left( \frac{z^3}{e^{12y}-1} + \frac{z^3}{e^{24y}-1} + \frac{z^3}{e^{36y}-1} + \frac{z^3}{e^{48y}-1} + \dots \right)$$

$$= z^4 (1-x+x^2)$$

$$\text{ii } 1 - 504 \left( \frac{z^5}{e^{12y}-1} + \frac{z^5}{e^{24y}-1} + \frac{z^5}{e^{36y}-1} + \frac{z^5}{e^{48y}-1} + \dots \right)$$

$$= z^6 (1+x)(1-\frac{x}{2})(1-2x)$$

$$\text{iii } 1 + 240 \left( \frac{z^7}{e^{12y}-1} + \frac{z^7}{e^{24y}-1} + \frac{z^7}{e^{36y}-1} + \dots \right)$$

$$= z^2 (1+14x+x^2)$$

$$\text{iv. } 1 - 504 \left( \frac{z^5}{e^{12y}-1} + \frac{z^5}{e^{24y}-1} + \frac{z^5}{e^{36y}-1} + \dots \right)$$

$$= z^6 (1+x)(1-34x+x^2)$$

$$\text{v. } 1 + 240 \left( \frac{z^3}{e^{12y}-1} + \frac{z^3}{e^{24y}-1} + \frac{z^3}{e^{36y}-1} + \dots \right)$$

$$= z^4 (1-x + \frac{x^2}{16})$$

$$\text{vi. } 1 - 504 \left( \frac{z^5}{e^{12y}-1} + \frac{z^5}{e^{24y}-1} + \frac{z^5}{e^{36y}-1} + \dots \right)$$

$$= z^6 (1-\frac{x}{2})(1-x-\frac{x^2}{32})$$

vii. If  $x$  is changed to  $\left(\frac{1-\sqrt{1-x}}{1+\sqrt{1-x}}\right)^2$  then  $y$  is changed to  $2y$

$$\text{viii } 1 + 24 \left( \frac{z}{e^{2y}+1} + \frac{z}{e^{4y}+1} + \frac{z}{e^{6y}+1} + \frac{z}{e^{8y}+1} + \dots \right)$$

$$= z^2 (1+x)$$

$$\text{ix. } 1 + 24 \left( \frac{z}{e^{2y}+1} + \frac{z}{e^{4y}+1} + \frac{z}{e^{6y}+1} + \dots \right)$$

$$= z^2 (1-\frac{x}{2})$$

$$x. 1 - 240 \left( \frac{1^3}{e^9+1} + \frac{2^3}{e^{19}+1} + \frac{3^3}{e^{29}+1} + \dots \right) \\ = 2^4 (1 - 16x + x^2)$$

$$xi. 1 + 504 \left( \frac{1^5}{e^9+1} + \frac{2^5}{e^{19}+1} + \frac{3^5}{e^{29}+1} + \dots \right) \\ = 2^6 (1+x)(1+29x+x^2)$$

$$xii. 1 - 240 \left( \frac{1^2}{e^{15}+1} + \frac{2^2}{e^{45}+1} + \frac{3^2}{e^{65}+1} + \dots \right) \\ = 2^4 \left( 1 - x - \frac{7}{8} x^2 \right)$$

$$xiii. 1 + 504 \left( \frac{1^5}{e^{15}+1} + \frac{2^5}{e^{45}+1} + \frac{3^5}{e^{65}+1} + \dots \right) \\ = 2^6 \left( 1 - \frac{x}{2} \right) \left( 1 - x + \frac{31}{16} x^2 \right)$$

$$14. i. 1 - 8 \left( \frac{1}{e^9+1} - \frac{2}{e^{19}+1} + \frac{3}{e^{29}+1} - \dots \right) = 2^2 (1-x)$$

$$ii. 1 + 16 \left( \frac{1^3}{e^9+1} - \frac{2^3}{e^{19}+1} + \frac{3^3}{e^{29}+1} - \dots \right) = 2^4 (1-x^4)$$

$$iii. 1 - 8 \left( \frac{1^5}{e^9+1} - \frac{2^5}{e^{19}+1} + \frac{3^5}{e^{29}+1} - \dots \right) = 2^6 (1-x)(1-x+x^2)$$

$$iv. 17 + 32 \left( \frac{1^7}{e^9+1} - \frac{2^7}{e^{19}+1} + \frac{3^7}{e^{29}+1} - \dots \right) = 2^8 (1-x^4)(17-32x+x^4)$$

$$v. 1 - 16 \left( \frac{1^2}{e^9-1} - \frac{2^2}{e^{19}-1} + \frac{3^2}{e^{29}-1} - \dots \right) = 2^4 (1-x)^4$$

$$vi. 1 + 8 \left( \frac{1^5}{e^9-1} - \frac{2^5}{e^{19}-1} + \frac{3^5}{e^{29}-1} - \dots \right) = 2^6 (1-x)(1-x^4)$$

$$vii. 17 - 32 \left( \frac{1^7}{e^9-1} - \frac{2^7}{e^{19}-1} + \frac{3^7}{e^{29}-1} - \dots \right) =$$

$$2^8 (1-x)^2 (17 - 2x + 17x^4)$$

$$viii. 31 + 8 \left( \frac{1^9}{e^9-1} - \frac{2^9}{e^{19}-1} + \frac{3^9}{e^{29}-1} - \dots \right) =$$

$$2^{10} (1-x)(1-x^4)(31 - 46x + 21x^4)$$

$$ix. 1 - 16 \left( \frac{1^2}{e^{15}-1} - \frac{2^2}{e^{45}-1} + \dots \right) = 2^4 (1-x)$$

x.  $\frac{15}{e^{2y}-e^{2y-1}} - \frac{15}{e^{4y}-e^{4y-1}} + \dots = 2^6(1-x)(1-\frac{x}{2})$

xi.  $\frac{17}{e^{2y}-e^{2y-1}} - \frac{17}{e^{4y}-e^{4y-1}} + \frac{17}{e^{6y}-e^{6y-1}} - \dots = 2^8(1-x)(17-17x+2x^4)$

xii. If x is changed to  $-\frac{x}{1-x}$  then y is changed to  $-e^{-y}$ .

15. i.  $\frac{13}{e^y-e^{-y}} + \frac{2^3}{e^{2y}-e^{-2y}} + \frac{3^3}{e^{3y}-e^{-3y}} + \dots = 2^4 \frac{x}{16}$

ii.  $\frac{15}{e^y-e^{-y}} + \frac{2^5}{e^{2y}-e^{-2y}} + \frac{3^5}{e^{3y}-e^{-3y}} + \dots = 2^6 \frac{x(1+x)}{16}$

iii.  $\frac{17}{e^y-e^{-y}} + \frac{2^7}{e^{2y}-e^{-2y}} + \frac{3^7}{e^{3y}-e^{-3y}} + \dots = 2^8 \frac{x(1+6\frac{1}{2}x+x^4)}{16}$

iv.  $\frac{19}{e^y-e^{-y}} + \frac{2^9}{e^{2y}-e^{-2y}} + \frac{3^9}{e^{3y}-e^{-3y}} + \dots = 2^{10} \frac{x(1+x)(1+29x+x^4)}{16}$

v.  $\frac{13}{e^{1y}-e^{-1y}} + \frac{2^3}{e^{2y}-e^{-2y}} + \frac{3^3}{e^{3y}-e^{-3y}} + \dots = 2^4 \frac{x^2}{256}$

vi.  $\frac{15}{e^{2y}-e^{-2y}} + \frac{2^5}{e^{4y}-e^{-4y}} + \frac{3^5}{e^{6y}-e^{-6y}} + \dots = 2^6 \frac{x^2}{256} (1-\frac{x}{2})$

vii.  $\frac{17}{e^{2y}-e^{-2y}} + \frac{2^7}{e^{4y}-e^{-4y}} + \frac{3^7}{e^{6y}-e^{-6y}} + \dots = 2^8 \frac{x^2(1-x+\frac{17}{32}x^4)}{256}$

viii.  $\frac{19}{e^{2y}-e^{-2y}} + \frac{2^9}{e^{4y}-e^{-4y}} + \frac{3^9}{e^{6y}-e^{-6y}} + \dots = 2^{10} \frac{x^2(1-x+\frac{31}{16}x^4)}{256}$

ix.  $\frac{1}{e^y-e^{-y}} + \frac{2}{e^{2y}-e^{-2y}} + \frac{3}{e^{3y}-e^{-3y}} + \dots = 2^4 \frac{x}{16}$

x.  $\frac{13}{e^y-e^{-y}} + \frac{2^3}{e^{2y}-e^{-2y}} + \frac{3^3}{e^{3y}-e^{-3y}} + \dots = 2^4 \frac{x}{16} (1-\frac{x}{2})$



xi.  $\frac{1^5}{e^2 - e^{-2}} + \frac{3^5}{e^{12} - e^{-12}} + \frac{5^5}{e^{32} - e^{-32}} + \dots = 2^6 \frac{x}{16} (1-x+x^4)$

xii.  $\frac{1^7}{e^2 - e^{-2}} + \frac{3^7}{e^{12} - e^{-12}} + \frac{5^7}{e^{32} - e^{-32}} + \dots = 2^7 \frac{x}{16} (1 - \frac{x}{2})(1-x + \frac{17}{2}x^4)$

xiii.  $\frac{1}{e^{\frac{1}{2}} - e^{-\frac{1}{2}}} + \frac{3}{e^{\frac{9}{2}} - e^{-\frac{9}{2}}} + \frac{5}{e^{\frac{25}{2}} - e^{-\frac{25}{2}}} + \dots = \frac{2}{4} \sqrt{x}$

xiv.  $\frac{1^3}{e^{\frac{1}{2}} - e^{-\frac{1}{2}}} + \frac{3^3}{e^{\frac{9}{2}} - e^{-\frac{9}{2}}} + \frac{5^3}{e^{\frac{25}{2}} - e^{-\frac{25}{2}}} + \dots = 2^4 \frac{\sqrt{x}}{4} (1+x)$

xv.  $\frac{1^5}{e^{\frac{1}{2}} - e^{-\frac{1}{2}}} + \frac{3^5}{e^{\frac{9}{2}} - e^{-\frac{9}{2}}} + \frac{5^5}{e^{\frac{25}{2}} - e^{-\frac{25}{2}}} + \dots = 2^6 \frac{\sqrt{x}}{4} (1+14x+x^4)$

xvi.  $\frac{1^7}{e^{\frac{1}{2}} - e^{-\frac{1}{2}}} + \frac{3^7}{e^{\frac{9}{2}} - e^{-\frac{9}{2}}} + \frac{5^7}{e^{\frac{25}{2}} - e^{-\frac{25}{2}}} + \dots = 2^8 \frac{\sqrt{x}}{4} (1+x)(1+134x+x^4)$

16. i.  $\frac{1}{e^{\frac{1}{2}} + e^{-\frac{1}{2}}} - \frac{3}{e^{\frac{9}{2}} + e^{-\frac{9}{2}}} + \frac{5}{e^{\frac{25}{2}} + e^{-\frac{25}{2}}} - \dots = \frac{2^2}{4} \sqrt{x(1-x)}$

ii.  $\frac{1^3}{e^{\frac{1}{2}} + e^{-\frac{1}{2}}} - \frac{3^3}{e^{\frac{9}{2}} + e^{-\frac{9}{2}}} + \frac{5^3}{e^{\frac{25}{2}} + e^{-\frac{25}{2}}} - \dots = \frac{2^4}{4} \sqrt{x(1-x)} (1-2x)$

iii.  $\frac{1^5}{e^{\frac{1}{2}} + e^{-\frac{1}{2}}} - \frac{3^5}{e^{\frac{9}{2}} + e^{-\frac{9}{2}}} + \frac{5^5}{e^{\frac{25}{2}} + e^{-\frac{25}{2}}} - \dots = \frac{2^6}{4} \sqrt{x(1-x)} \{1-16x(1-x)\}$

iv.  $\frac{1^7}{e^{\frac{1}{2}} + e^{-\frac{1}{2}}} - \frac{3^7}{e^{\frac{9}{2}} + e^{-\frac{9}{2}}} + \frac{5^7}{e^{\frac{25}{2}} + e^{-\frac{25}{2}}} - \dots = \frac{2^8}{4} \sqrt{x(1-x)} (1-2x) \{1-136x(1-x)\}$

v.  $\frac{1^9}{e^{\frac{1}{2}} + e^{-\frac{1}{2}}} - \frac{3^9}{e^{\frac{9}{2}} + e^{-\frac{9}{2}}} + \frac{5^9}{e^{\frac{25}{2}} + e^{-\frac{25}{2}}} - \dots = \frac{2^{10}}{4} \sqrt{x(1-x)} \{1-1232x(1-x)+7936x^2(1-x)^2\}$

vi.  $\frac{1^{11}}{e^{\frac{1}{2}} + e^{-\frac{1}{2}}} - \frac{3^{11}}{e^{\frac{9}{2}} + e^{-\frac{9}{2}}} + \frac{5^{11}}{e^{\frac{25}{2}} + e^{-\frac{25}{2}}} - \dots = \frac{2^{12}}{4} \sqrt{x(1-x)} (1-2x) \{1-11072x(1-x)+176896x^2(1-x)^2\}$

$$\text{vii } \tan^{-1} e^{-x/2} - \tan^{-1} e^{-3x/2} + \tan^{-1} e^{-5x/2} - \dots = \frac{1}{2} \sin^{-1} \sqrt{x}$$

$$\text{viii } \tan^{-1} e^{-x/4} - \tan^{-1} e^{-3x/4} + \tan^{-1} e^{-5x/4} - \dots = \frac{1}{2} \tan^{-1} \sqrt{x}$$

$$\text{ix } \frac{1}{e^x + e^{-x}} + \frac{1}{e^{3x/2} + e^{-3x/2}} + \frac{1}{e^{5x/2} + e^{-5x/2}} = 2 \frac{\sqrt{x}}{4}$$

$$\text{x } \frac{1^2}{e^{3x/2} + e^{-3x/2}} + \frac{3^2}{e^{5x/2} + e^{-5x/2}} + \frac{5^2}{e^{7x/2} + e^{-7x/2}} = 2^3 \frac{\sqrt{x}}{4}$$

$$\text{xi } \frac{1^4}{e^{3x/2} + e^{-3x/2}} + \frac{3^4}{e^{5x/2} + e^{-5x/2}} + \frac{5^4}{e^{7x/2} + e^{-7x/2}} = 2^5 \frac{\sqrt{x}}{4} (1+4x)$$

$$\text{xii } \frac{1^6}{e^{3x/2} + e^{-3x/2}} + \frac{3^6}{e^{5x/2} + e^{-5x/2}} + \frac{5^6}{e^{7x/2} + e^{-7x/2}} + \dots = 2^7 \frac{\sqrt{x}}{4} \{1 + 11(4x) + 8(x)^2\}$$

$$\text{xiii } \frac{1^8}{e^{3x/2} + e^{-3x/2}} + \frac{3^8}{e^{5x/2} + e^{-5x/2}} + \frac{5^8}{e^{7x/2} + e^{-7x/2}} + \dots = 2^9 \frac{\sqrt{x}}{4} \{1 + 57(4x) + 109(4x)^2 + (4x)^3\}$$

$$17. \text{ i. } 1 + 4 \left( \frac{1}{e^x + e^{-x}} + \frac{1}{e^{2x} + e^{-2x}} + \frac{1}{e^{4x} + e^{-4x}} + \dots \right) = 2$$

$$\text{ii. } 4 \left( \frac{1^2}{e^x + e^{-x}} + \frac{2^2}{e^{2x} + e^{-2x}} + \frac{3^2}{e^{4x} + e^{-4x}} + \dots \right) = 2^3 \frac{x}{4}$$

$$\text{iii. } 4 \left( \frac{1^4}{e^x + e^{-x}} + \frac{2^4}{e^{2x} + e^{-2x}} + \frac{3^4}{e^{4x} + e^{-4x}} + \dots \right) = 2^5 \left\{ \frac{x}{4} + \left(\frac{x}{2}\right)^2 \right\}$$

$$\text{iv. } 4 \left( \frac{1^6}{e^x + e^{-x}} + \frac{2^6}{e^{2x} + e^{-2x}} + \frac{3^6}{e^{4x} + e^{-4x}} + \dots \right) = 2^7 \left\{ \frac{x}{4} + 11 \left(\frac{x}{2}\right)^2 + \left(\frac{x}{2}\right)^3 \right\}$$

$$\text{v. } 4 \left( \frac{1^8}{e^x + e^{-x}} + \frac{2^8}{e^{2x} + e^{-2x}} + \frac{3^8}{e^{4x} + e^{-4x}} + \dots \right) = 2^9 \left\{ \frac{x}{4} + 57 \left(\frac{x}{2}\right)^2 + 102 \left(\frac{x}{2}\right)^3 + \left(\frac{x}{2}\right)^4 \right\}$$

$$\text{vi. } 1 + 4 \left( \frac{1}{e^{2x}-1} - \frac{1}{e^{4x}-1} + \frac{1}{e^{8x}-1} - \dots \right) = 2$$

$$\text{vii } 1 - 4 \left( \frac{1^2}{e^{2x}-1} - \frac{3^2}{e^{4x}-1} + \frac{5^2}{e^{8x}-1} - \dots \right) = 2^3 (1-x)$$

$$\text{viii } 5 + 4 \left( \frac{1^4}{e^{2x}-1} - \frac{3^4}{e^{4x}-1} + \frac{5^4}{e^{8x}-1} - \dots \right) = 2^5 (5-x)(1-x)$$

ix.  $61 - 4 \left( \frac{1^6}{e^6 - 1} - \frac{3^6}{e^{12} - 1} + \frac{5^6}{e^{18} - 1} - \dots \right) = 2^7 (1-x)(61 - 46x + x^6)$

ex. i.  $\phi^8(x) = 1 + 16 \left( \frac{1^2 x}{1+x} + \frac{2^2 x^2}{1-x^2} + \frac{3^2 x^3}{1+x^3} + \frac{4^2 x^4}{1-x^4} + \dots \right)$

ii.  $x \psi^8(x) = \frac{1^3 x}{1-x^2} + \frac{2^3 x^2}{1-x^4} + \frac{3^3 x^3}{1-x^6} + \frac{4^3 x^4}{1-x^8} + \dots$

iii.  $x \psi^4(x^2) = \frac{x}{1-x^2} + \frac{3x^3}{1-x^6} + \frac{5x^5}{1-x^{10}} + \frac{7x^7}{1-x^{14}} + \dots$

iv.  $\psi^2(x^2) = \frac{1}{1+x} + \frac{x}{1+x^3} + \frac{x^2}{1+x^5} + \frac{x^3}{1+x^7} + \dots$

v.  $\phi^2(x) \psi^4(x) = \frac{1^2}{1+x} + \frac{2^2 x}{1+x^3} + \frac{3^2 x^2}{1+x^5} + \frac{4^2 x^3}{1+x^7} + \dots$

vi.  $\frac{1^9 x}{1-x^2} + \frac{2^9 x^2}{1-x^4} + \frac{3^9 x^3}{1-x^6} + \dots = x \psi^9(x) \left\{ 1 + 504 \left( \frac{1^5 x}{1+x} + \frac{2^5 x^2}{1+x^3} + \dots \right) \right\}$

18. i.  $\frac{1}{\cosh \frac{\pi}{2}} - \frac{1}{5 \cosh \frac{3\pi}{2}} + \frac{1}{5 \cosh \frac{5\pi}{2}} - \dots = \frac{\pi}{24}$

ii.  $\frac{1}{\cosh \frac{\pi}{2\sqrt{3}}} - \frac{1}{5 \cosh \frac{3\pi}{2\sqrt{3}}} + \frac{1}{5 \cosh \frac{5\pi}{2\sqrt{3}}} - \dots = \frac{5\pi}{24}$

iii.  $\frac{1^{6n-1}}{\cosh \frac{\pi}{2}} - \frac{3^{6n-1}}{\cosh \frac{3\pi}{2}} + \frac{5^{6n-1}}{\cosh \frac{5\pi}{2}} - \dots =$

$\frac{1^{6n-1}}{\cosh \frac{\pi}{2\sqrt{3}}} - \frac{3^{6n-1}}{\cosh \frac{3\pi}{2\sqrt{3}}} + \frac{5^{6n-1}}{\cosh \frac{5\pi}{2\sqrt{3}}} - \dots = 0$

*n* being any positive integer excluding 0.

ex. i. If  $\frac{1^7}{1+x} - \frac{3^7 x}{1+x^3} + \frac{5^7 x^2}{1+x^5} - \frac{7^7 x^3}{1+x^7} + \dots = 0$ , then

$X(x) = \sqrt[7]{2} \sqrt[7]{x}$  or  $\sqrt[7]{2} \cdot \sqrt[7]{34x}$

ii. If  $\frac{1^9}{1+x} - \frac{3^9 x}{1+x^3} + \frac{5^9 x^2}{1+x^5} - \frac{7^9 x^3}{1+x^7} + \dots = 0$ , then

$X(x) = \sqrt[9]{2} \cdot \sqrt[9]{(154 \pm 6\sqrt{645})x}$

iii. If  $\frac{1^{11}}{1+x} - \frac{3^{11} x}{1+x^3} + \frac{5^{11} x^2}{1+x^5} - \frac{7^{11} x^3}{1+x^7} + \dots = 0$ , then

$(1+x)(1+x^3)(1+x^5)(1+x^7)(1+x^9) \dots \propto X(x) =$

$\sqrt[11]{2} \sqrt[11]{x}$  or  $\sqrt[11]{2} \sqrt[11]{4x}$  or  $\sqrt[11]{2} \sqrt[11]{3764x}$

# CHAPTER XVIII

$$1. 1 - \left(\frac{1}{2}\right)x + \left(\frac{1 \cdot 1}{2 \cdot 4}\right)^2 x^2 + \left(\frac{1 \cdot 1 \cdot 3}{2 \cdot 4 \cdot 6}\right)^2 x^3 + \left(\frac{1 \cdot 1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 8}\right)^2 x^4 + \dots$$

$$= 2(0-x) + \int 2 dx = \frac{2}{3}(1+x) + \frac{2}{32} \left\{ 1 - 24 \left( \frac{1}{e^{18}} + \frac{2}{e^{42}} + \dots \right) \right\}$$

$$2. 1 - \frac{1}{2}x - \frac{1 \cdot 3}{2 \cdot 4} x^2 - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} x^3 - \frac{1 \cdot 3 \cdot 5 \cdot 7}{2 \cdot 4 \cdot 6 \cdot 8} x^4 - \dots$$

$$= 2(1-x) + \frac{1}{2} \int 2 dx = \frac{2}{3}(2-x) + \frac{1}{32} \left\{ 1 - 24 \left( \frac{1}{e^{18}} + \frac{2}{e^{42}} + \dots \right) \right\}$$

3. The perimeter of an ellipse whose eccentricity is  $h$ , is

$$2a\pi \left\{ 1 - \frac{1}{2} h^2 - \frac{1 \cdot 3}{2 \cdot 4} h^4 - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} h^6 - \dots \right\}$$

$$= \pi(a+b) \left\{ 1 + \left(\frac{1}{2}\right)^2 \left(\frac{a-b}{a+b}\right)^2 + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \left(\frac{a-b}{a+b}\right)^4 + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 \left(\frac{a-b}{a+b}\right)^6 + \dots \right\}$$

$$= \pi \left\{ 3(a+b) - \sqrt{(a+3b)(3a+b)} \right\} \text{ nearly}$$

$$= \pi(a+b) \left\{ 1 + \frac{3x}{10 + \sqrt{4-3x}} \right\} \text{ very nearly where } x = \left(\frac{a-b}{a+b}\right)^2.$$

N.B. i.  $\pi = 3.1415926535897932384626434$

ii.  $\log 10 = 2.302585092994045684018$

iii.  $e^{-\pi} = .04321391826377225$

iv.  $e^{\pi} = 4.810477380965351655473$

Con.  $\pi = \frac{355}{113} \left( 1 - \frac{.0003}{36.99} \right)$  very nearly

$$= \sqrt[4]{97\frac{1}{2} - \frac{1}{11}} \text{ nearly}$$

$$4. \frac{\sqrt{x}}{2} \left\{ 1 + \left(\frac{1}{2}\right)^2 \frac{x}{3} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \frac{x^2}{5} + \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 \frac{x^3}{7} + \dots \right\}$$

$$= \log \frac{1+e^{-x/2}}{1-e^{-x/2}} - 3 \log \frac{1+e^{-3x/2}}{1-e^{-3x/2}} + 5 \log \frac{1+e^{-5x/2}}{1-e^{-5x/2}} - \dots$$

$$5. \log \frac{1}{x} - \left(\frac{1}{2}\right)^2 \frac{x}{1} - \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 \frac{x^2}{2} - \left(\frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}\right)^2 \frac{x^3}{3} - \dots$$

$$= 9 - 4 \left\{ \log(1-e^{-x}) - 3 \log(1-e^{-3x}) + 5 \log(1-e^{-5x}) - \dots \right\}$$

$$5. \frac{1}{1^2(e^{\pi/2} + e^{-\pi/2})} + \frac{1}{3^2(e^{3\pi/2} + e^{-3\pi/2})} + \frac{1}{5^2(e^{5\pi/2} + e^{-5\pi/2})} + \dots$$

$$= \frac{\sqrt{x}}{4 \cdot 2} \left\{ 1 + \left(\frac{2}{3}\right)^2 x + \left(\frac{2 \cdot 4}{3 \cdot 5}\right)^2 x^2 + \left(\frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7}\right)^2 x^3 + \dots \right\}$$

$$7. \frac{1}{1^2(e^{\pi} + 1)} - \frac{1}{3^2(e^{3\pi} + 1)} + \frac{1}{5^2(e^{5\pi} + 1)} - \dots$$

$$= \frac{1}{2} \left( \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots \right) - \frac{\pi}{16} y$$

$$+ \frac{\sqrt{1-x}}{4 \cdot 2} \left\{ 1 + \left(\frac{2}{3}\right)^2 (1-x) + \left(\frac{2 \cdot 4}{3 \cdot 5}\right)^2 (1-x)^2 + \left(\frac{2 \cdot 4 \cdot 6}{3 \cdot 5 \cdot 7}\right)^2 (1-x)^3 + \dots \right\}$$

N.B.  $\frac{1}{1(e^{\pi/2} - e^{-\pi/2})} + \frac{1}{3(e^{3\pi/2} - e^{-3\pi/2})} + \frac{1}{5(e^{5\pi/2} - e^{-5\pi/2})} + \dots = \frac{1}{8} \log \frac{1+\sqrt{x}}{1-\sqrt{x}}$

$$8. i. \frac{\cos \theta + 2 \cos \frac{\theta}{2} \cosh \frac{\theta \sqrt{3}}{2}}{\cosh \frac{\pi \sqrt{3}}{2}} - \frac{\cos 3\theta + 2 \cos \frac{3\theta}{2} \cosh \frac{3\theta \sqrt{3}}{2}}{3 \cosh \frac{3\pi \sqrt{3}}{2}}$$

$$+ \frac{\cos 5\theta + 2 \cos \frac{5\theta}{2} \cosh \frac{5\theta \sqrt{3}}{2}}{5 \cosh \frac{5\pi \sqrt{3}}{2}} - \dots = \frac{\pi}{8}$$

$$ii. \frac{\cos \theta}{\cosh \frac{\pi \sqrt{3}}{2}} (\cos \theta + \cosh \theta \sqrt{3}) - \frac{\cos 3\theta}{3 \cosh \frac{3\pi \sqrt{3}}{2}} (\cos 3\theta + \cosh 3\theta \sqrt{3})$$

$$+ \frac{\cos 5\theta}{5 \cosh \frac{5\pi \sqrt{3}}{2}} (\cos 5\theta + \cosh 5\theta \sqrt{3}) - \dots = \frac{\pi}{12}$$

$$iii. \frac{\sin \theta}{1^4 \cosh \frac{\pi \sqrt{3}}{2}} (\cos \theta - \cosh \theta \sqrt{3}) - \frac{\sin 3\theta}{3^4 \cosh \frac{3\pi \sqrt{3}}{2}} (\cos 3\theta - \cosh 3\theta \sqrt{3})$$

$$+ \frac{\sin 5\theta}{5^4 \cosh \frac{5\pi \sqrt{3}}{2}} (\cos 5\theta - \cosh 5\theta \sqrt{3}) - \dots = \frac{\pi}{12} \theta^3$$

$$iv. \frac{\cos \theta}{17 \cosh \frac{\pi \sqrt{3}}{2}} (\cos \theta + \cosh \theta \sqrt{3}) - \frac{\cos 3\theta}{37 \cosh \frac{3\pi \sqrt{3}}{2}} (\cos 3\theta + \cosh 3\theta \sqrt{3})$$

$$+ \frac{\cos 5\theta}{57 \cosh \frac{5\pi \sqrt{3}}{2}} (\cos 5\theta + \cosh 5\theta \sqrt{3}) - \dots$$

$$= \frac{\pi 7}{11520} - \frac{\pi \theta^6}{180}$$

$$9. \frac{x^{15}}{1-x^6} \cdot \frac{1}{\cosh \frac{\pi\sqrt{3}}{2}} - \frac{x^5}{3^6-x^6} \cdot \frac{1}{\cosh \frac{3\pi\sqrt{3}}{2}} + \dots$$

$$= \frac{\pi}{12} \frac{1}{\cos \frac{\pi x}{2} \{ \cos \frac{\pi x}{2} + \cosh \frac{\pi x \sqrt{3}}{2} \}}$$

i.  $\frac{1}{2} \cos \frac{x}{2} \{ \cos \frac{x}{2} + \cosh \frac{x\sqrt{3}}{2} \}$

$$= 1 - \frac{x^6}{4} \left\{ \frac{x^6}{12} - \frac{x^{12}}{112} + \frac{x^{18}}{112} - \frac{x^{24}}{1344} + \dots \right\}$$

$$= \left(1 - \frac{x^6}{116}\right) \left(1 - \frac{x^6}{3^2 \pi^6}\right) \left(1 - \frac{x^6}{5^2 \pi^6}\right) \left(1 - \frac{x^6}{7^2 \pi^6}\right) \dots$$

ii.  $\frac{1}{2} \sin \frac{x}{2} \{ \cos \frac{x}{2} - \cosh \frac{x\sqrt{3}}{2} \}$

$$= -\frac{x^3}{2} \left\{ \frac{x^3}{12} - \frac{x^9}{112} + \frac{x^{15}}{112} - \frac{x^{21}}{121} + \dots \right\}$$

$$= -\frac{x^3}{8} \left(1 - \frac{x^6}{2^2 \pi^6}\right) \left(1 - \frac{x^6}{4^2 \pi^6}\right) \left(1 - \frac{x^6}{6^2 \pi^6}\right) \dots$$

10.  $\frac{x^{15}}{1-x^6} \cdot \frac{1}{\cosh \frac{\pi}{2\sqrt{3}}} - \frac{x^5}{3^6-x^6} \cdot \frac{1}{\cosh \frac{3\pi}{2\sqrt{3}}} + \dots$

$$= \frac{\pi}{12} \frac{4 \cosh \frac{\pi x}{2\sqrt{3}} (\cos \frac{\pi x}{2} + \cosh \frac{\pi x}{2\sqrt{3}}) - 3}{\cos \frac{\pi x}{2} \{ \cos \frac{\pi x}{2} + \cosh \frac{\pi x \sqrt{3}}{2} \}}$$

i.  $\frac{x^3}{1-x^6} + \frac{x^9}{3^6-x^6} + \frac{x^{15}}{5^6-x^6} + \frac{x^{21}}{7^6-x^6} + \dots$

$$= \frac{\pi}{12} \frac{\cosh \frac{\pi x \sqrt{3}}{2} - \cos \frac{\pi x}{2}}{\cosh \frac{\pi x \sqrt{3}}{2} + \cos \frac{\pi x}{2}} \tan \frac{\pi x}{2}$$

ex.  $\frac{1}{1^7 \cosh \frac{\pi\sqrt{3}}{2}} - \frac{1}{3^7 \cosh \frac{3\pi\sqrt{3}}{2}} + \frac{1}{5^7 \cosh \frac{5\pi\sqrt{3}}{2}} - \dots$

$$= \frac{\pi^7}{23040}$$

ii.  $\left\{ 1 + 2 \left( \frac{\cos \theta}{\cosh \pi} + \frac{\cos 2\theta}{\cosh 2\pi} + \frac{\cos 3\theta}{\cosh 3\pi} + \dots \right) \right\}^{-2}$

$$+ \left\{ 1 + 2 \left( \frac{\cosh \theta}{\cosh \pi} + \frac{\cosh 2\theta}{\cosh 2\pi} + \frac{\cosh 3\theta}{\cosh 3\pi} + \dots \right) \right\}^{-2} = \frac{2}{(\sqrt{\pi}/\sqrt{2})^2}$$

ii.  $\left\{ \frac{\cos \theta}{\cosh \frac{\theta}{2}} + \frac{\cos 3\theta}{\cosh \frac{3\theta}{2}} + \frac{\cos 5\theta}{\cosh \frac{5\theta}{2}} + \dots \right\} \times$

$$\left\{ \frac{\cosh \theta}{\cosh \frac{\theta}{2}} + \frac{\cosh 3\theta}{\cosh \frac{3\theta}{2}} + \frac{\cosh 5\theta}{\cosh \frac{5\theta}{2}} + \dots \right\} = \frac{2\sqrt{2}}{4} \sqrt{x(1-x)}$$

$$12. i. \frac{1}{2} + \frac{\operatorname{sech} y}{1+n^2} + \frac{\operatorname{sech} 2y}{1+(2n)^2} + \frac{\operatorname{sech} 3y}{1+(3n)^2} + \dots$$

$$= \frac{x}{2} + \frac{(n^2)^{-x}}{2} + \frac{(2n^2)^{-x}}{2} + \frac{(3n^2)^{-x}}{2} + \frac{(4n^2)^{-x}}{2} + \dots$$

$$ii. \frac{\operatorname{sech} y/2}{1+n^2} + \frac{\operatorname{sech} 3y/2}{1+(3n)^2} + \frac{\operatorname{sech} 5y/2}{1+(5n)^2} + \dots$$

$$= \frac{1}{2} \cdot \frac{2\sqrt{x}}{1+x} \frac{(n^2)^{-x}}{1+x} + \frac{(2n^2)^{-x}}{1+x} + \frac{(3n^2)^{-x}}{1+x} + \frac{(4n^2)^{-x}}{1+x} + \dots$$

Cor. If A and G be the A.M and G.M between  $\alpha$  and  $\beta$

and  $F(\alpha, \beta) = \frac{\alpha}{n} + \frac{(\alpha)^L}{n} + \frac{(2\alpha)^L}{n} + \frac{(3\alpha)^L}{n} + \frac{(4\alpha)^L}{n} + \dots$  then

$F(A, G)$  is the A.M between  $F(\alpha, \beta)$  and  $F(\beta, \alpha)$ .

$$13. i. \frac{\operatorname{cosech} y/2}{1+n^2} - \frac{\operatorname{cosech} 3y/2}{1+(3n)^2} + \frac{\operatorname{cosech} 5y/2}{1+(5n)^2} - \dots$$

$$= \frac{1}{2} \cdot \frac{2\sqrt{x}}{1-x} \frac{(1-x)(n^2)^{-x}}{1-x} - \frac{x(2n^2)^{-x}}{1-x} + \frac{(1-x)(3n^2)^{-x}}{1-x} - \dots$$

$$ii. \frac{\operatorname{sech} y/2}{1+n^2} - \frac{3 \operatorname{sech} 3y/2}{1+(3n)^2} + \frac{5 \operatorname{sech} 5y/2}{1+(5n)^2} - \dots$$

$$= \frac{1}{2} \cdot \frac{2^x \sqrt{x(1-x)}}{1+(n^2)^x(1-2x)} + \frac{2^x(2^2-1)x(1-x)(n^2)^x}{1+(3n^2)^x(1-2x)} + \frac{4^x(4^2-1)x(1-x)(n^2)^x}{1+(5n^2)^x(1-2x)} + \dots$$

$$iii. \frac{\operatorname{cosech} y/2}{1+n^2} + \frac{3 \operatorname{cosech} 3y/2}{1+(3n)^2} + \frac{5 \operatorname{cosech} 5y/2}{1+(5n)^2} + \dots$$

$$= \frac{1}{2} \cdot \frac{2^x \sqrt{x}}{1+(n^2)^x(1+x)} - \frac{2^x(2^2-1)x(n^2)^x}{1+(3n^2)^x(1+x)} + \frac{4^x(4^2-1)x(n^2)^x}{1+(5n^2)^x(1+x)} - \dots$$

Cor.  $\frac{\operatorname{sech} \pi/2}{1+n^2} - \frac{3 \operatorname{sech} 3\pi/2}{1+(3n)^2} + \frac{5 \operatorname{sech} 5\pi/2}{1+(5n)^2} - \frac{7 \operatorname{sech} 7\pi/2}{1+(7n)^2} + \dots$

$$= \frac{1}{2} + \frac{1.3 (n\pi)^4}{1 + \frac{6.10 (n\pi)^4}{1 + \frac{15.31 (n\pi)^4}{1 + \dots}} \quad \text{where } n = \frac{\sqrt{x}}{(\sqrt{1-x})^2}$$

14. Let  $S = \frac{\sin \theta}{\sinh \frac{\theta}{2}} + \frac{\sin 3\theta}{\sinh \frac{3\theta}{2}} + \frac{\sin 5\theta}{\sinh \frac{5\theta}{2}} + \dots$

$$C = \frac{\cos \theta}{\cosh \frac{\theta}{2}} + \frac{\cos 3\theta}{\cosh \frac{3\theta}{2}} + \frac{\cos 5\theta}{\cosh \frac{5\theta}{2}} + \dots$$

and  $C_1 = \frac{1}{2} + \frac{\cos \theta}{\cosh \theta} + \frac{\cos 2\theta}{\cosh 2\theta} + \frac{\cos 3\theta}{\cosh 3\theta} + \dots$ , then

we see that  $C^2 + S^2 = \frac{x}{1-x} z^2$  and  $C_1^2 + S_1^2 = \frac{z^2}{1-x}$ .

and  $CS = \frac{\sin \theta}{\cosh \theta} + \frac{2 \sin 2\theta}{\cosh 2\theta} + \frac{3 \sin 3\theta}{\cosh 3\theta} + \dots$

$\therefore CS + \frac{dC_1}{d\theta} = 0$ ;  $C_1 S + \frac{dC}{d\theta} = 0$  and  $CC_1 = \frac{dS}{d\theta}$ .

Let  $C = \frac{\sqrt{x}}{2} z \cos \phi$  and  $S = \frac{\sqrt{x}}{2} z \sin \phi$

$\therefore C_1 = \frac{z}{2} \sqrt{1-x} \sin \phi$ .

$\therefore \frac{z}{2} \cos \phi \sqrt{1-x} \sin \phi = -\frac{d \sin \phi}{d\theta} = \cos \phi \frac{d\phi}{d\theta}$

$\therefore \theta = \frac{z}{2} \int_0^\phi \frac{d\phi}{\sqrt{1-x} \sin \phi}$

15. Let  $z\theta = \int_0^\phi \frac{d\phi}{\sqrt{1-x} \sin \phi}$ ;  $y = \pi \cdot \frac{z'}{z}$ ;  $y' = \pi \cdot \frac{z''}{z} - \pi \cdot \frac{z'^2}{z^2}$

$z' = 1 + (\frac{1}{2})^x (1-x) + (\frac{1-3}{2})^x (1-x) + \dots$  and  $z = 1 + (\frac{1}{2})^x + (\frac{1-3}{2})^x + \dots$

i.  $1 + z \left( \frac{\cos 3\theta}{\cosh 3\theta} + \frac{\cos \theta}{\cosh \theta} + \frac{\cos 5\theta}{\cosh 5\theta} + \dots \right) = 2 \sqrt{1-x} \sin \phi$

ii.  $\frac{\cos \theta}{\cosh \frac{\theta}{2}} + \frac{\cos 3\theta}{\cosh \frac{3\theta}{2}} + \frac{\cos 5\theta}{\cosh \frac{5\theta}{2}} + \dots = \frac{\sqrt{x}}{2} z \cos \phi$

iii.  $\frac{\sin \theta}{\sinh \frac{\theta}{2}} + \frac{\sin 3\theta}{\sinh \frac{3\theta}{2}} + \frac{\sin 5\theta}{\sinh \frac{5\theta}{2}} + \dots = \frac{\sqrt{x}}{2} z \sin \phi$



$$IV. \theta + \frac{\sin 2\theta}{\cosh \frac{\theta}{2}} + \frac{\sin 4\theta}{2 \cosh \frac{\theta}{2}} + \frac{\sin 6\theta}{3 \cosh \frac{\theta}{2}} + \dots = \phi$$

$$V. \frac{\sin \theta}{\cosh \frac{\theta}{2}} + \frac{\sin 3\theta}{3 \sinh \frac{\theta}{2}} + \frac{\sin 5\theta}{5 \sinh \frac{\theta}{2}} + \dots = \frac{1}{2} \sin^{-1}(\sqrt{x} \sin \phi)$$

$$VI. \frac{\cos \theta}{\sinh \frac{\theta}{2}} + \frac{\cos 3\theta}{3 \sinh \frac{\theta}{2}} + \frac{\cos 5\theta}{5 \sinh \frac{\theta}{2}} + \dots = \frac{1}{2} \log \frac{\sqrt{1-x} \sin \phi - \sqrt{1-x}}{\sqrt{1-x}}$$

16. If  $\theta$  is changed to  $\frac{\pi}{2} - \theta$ , then  $\cot \phi$  to  $\sqrt{1-x} \tan \phi$ ;

$\sin \phi$  to  $\frac{\cos \phi}{\sqrt{1-x \sin^2 \phi}}$ ;  $\cos \phi$  to  $\frac{\sin \phi}{\sqrt{1-x \sin^2 \phi}}$

$\sqrt{1-x \sin^2 \phi}$  to  $\frac{\sqrt{1-x}}{\sqrt{1-x \sin^2 \phi}}$

$$i. \frac{\cos \theta}{\sinh \frac{\theta}{2}} - \frac{\cos 3\theta}{\sinh \frac{3\theta}{2}} + \frac{\cos 5\theta}{\sinh \frac{5\theta}{2}} - \dots = \frac{\sqrt{x}}{2} \frac{\cos \phi}{\sqrt{1-x \sin^2 \phi}}$$

$$ii. \frac{\sin \theta}{\cosh \frac{\theta}{2}} - \frac{\sin 3\theta}{\cosh \frac{3\theta}{2}} + \frac{\sin 5\theta}{\cosh \frac{5\theta}{2}} - \dots = \frac{\sqrt{x(1-x)}}{2} \frac{\sin \phi}{\sqrt{1-x \sin^2 \phi}}$$

$$iii. \operatorname{cosec} \theta + 4 \left( \frac{\sin \theta}{e^{\theta} - 1} + \frac{\sin 3\theta}{e^{3\theta} - 1} + \frac{\sin 5\theta}{e^{5\theta} - 1} + \dots \right) = 2 \operatorname{cosec} \phi$$

$$iv. \sec \theta + 4 \left( \frac{\cos \theta}{e^{\theta} - 1} - \frac{\cos 3\theta}{e^{3\theta} - 1} + \frac{\cos 5\theta}{e^{5\theta} - 1} - \dots \right) = 2 \sec \phi \sqrt{1-x \sin^2 \phi}$$

$$v. \log \tan \left( \frac{\pi}{4} + \frac{\theta}{2} \right) + 4 \left\{ \frac{\sin \theta}{e^{\theta} - 1} - \frac{\sin 3\theta}{3(e^{3\theta} - 1)} + \frac{\sin 5\theta}{5(e^{5\theta} - 1)} - \dots \right\} = \log \tan \left( \frac{\pi}{4} + \frac{\phi}{2} \right)$$

$$17. i. \frac{\cos \theta}{\sin^3 \theta} = 8 \left( \frac{1 - \sin 2\theta}{e^{2\theta} - 1} + \frac{2^2 \sin 4\theta}{e^{4\theta} - 1} + \frac{3^2 \sin 6\theta}{e^{6\theta} - 1} + \dots \right) = 2^3 \frac{\cos \phi}{\sin^3 \phi} \sqrt{1-x \sin^2 \phi}$$

$$ii. \frac{1}{\sin^2 \theta} = 8 \left( \frac{\cos 2\theta}{e^{2\theta} - 1} + \frac{2 \cos 4\theta}{e^{4\theta} - 1} + \frac{3 \cos 6\theta}{e^{6\theta} - 1} + \dots \right)$$

$$= \frac{1}{3} \left( 1 - 24 \left( \frac{1}{e^{2\theta}} + \frac{2}{e^{4\theta}} + \frac{1}{e^{6\theta}} + \dots \right) \right)$$

$$\text{iii. } \frac{\sin 2\theta}{e^{2\theta}} + \frac{\sin 4\theta}{e^{4\theta}} + \frac{\sin 6\theta}{e^{6\theta}} + \dots$$

$$= 2 \int_0^\phi \sqrt{x \sin^2 \phi} d\phi + 2 \int_0^\phi \sqrt{x \sin^2 \phi} d\phi - \frac{2\theta z}{\pi} \int_0^\pi \sqrt{x \sin^2 \phi} d\phi$$

$$\text{iv. } \frac{\sin 2\theta}{\sinh y} + \frac{\sin 4\theta}{\sinh 2y} + \frac{\sin 6\theta}{\sinh 3y} + \dots$$

$$= \frac{2}{\pi} \int_0^\phi \sqrt{x \sin^2 \phi} d\phi - \frac{\theta z}{\pi} \int_0^\pi \sqrt{x \sin^2 \phi} d\phi$$

18. i. If  $\theta$  is changed to  $\frac{\theta}{2}$  and  $y$  to  $\frac{y}{2}$ , then  $x$  must be changed to  $\frac{4\sqrt{x}}{(1+\sqrt{x})^2}$  and  $z\phi$  to  $\phi + \sin^{-1}(\sqrt{x} \sin \phi)$  and  $z$  to  $(1+\sqrt{x})z$ .

ii. If  $\theta$  is changed to  $\frac{\pi}{2} - \theta$  and  $e^{-y}$  to  $-e^{-y}$ , then  $x$  must be changed to  $-\frac{x}{1-x}$ ;  $\phi$  to  $\frac{\pi}{2} - \phi$ ; &  $z$  to  $z\sqrt{1-x}$ .

iii. If  $e^{-y}$  is changed to  $-e^{-y}$ , then change  $x$  to  $-\frac{x}{1-x}$ ;  $z$  to  $z\sqrt{1-x}$  and  $\cot \phi$  to  $\cot \phi \sqrt{1-x}$ .

iv. If  $\theta$  is changed to  $i\theta$  and  $y$  to  $y'$ , then change  $x$  to  $1-x$ ;  $z$  to  $z'$ ;  $\sin \phi$  to  $i \tan \phi$ ;  $\cos \phi$  to  $\sec \phi$ ; and  $\phi$  to  $i \log \tan(\frac{\pi}{4} + \frac{\phi}{2})$ .


19. i. The length of the arc AP

$$\text{in an ellipse} = a \int_0^\phi \sqrt{1 - e^2 \cos^2 \phi} d\phi$$

where  $e$  is the eccentricity.

ii. The length of AP in a hyperbola

$$= a \tan \phi \sqrt{e^2 - \cos^2 \phi} - a \int_0^\phi \sqrt{e^2 - \cos^2 \phi} d\phi$$



$\frac{b^2}{a} \int_0^\phi \frac{d\phi}{\sqrt{e^2 - \cos^2 \phi}}$  where  $x = a \sec \phi$  and  $y = b \tan \phi$

iii. If the perimeter of an ellipse =  $\pi(a+b) \left(1 + \frac{1}{2} e^2 \sin^2 \theta_2\right)$  where  $\sin \theta = \frac{a-b}{a+b} \sin \phi$ . When  $e=1$ ,  $\phi = 30^\circ 18' 6''$  and very gradually diminishes to  $30^\circ$  when  $e$  becomes 0.

iv. If the perimeter of an ellipse =  $\pi(a+b) \left\{1 + \frac{\sin^2 \theta}{2 + \cos^2 \theta}\right\}$  where  $\sin \theta = \frac{a-b}{a+b} \sin \phi$ . When  $e=1$ ,  $\phi = 60^\circ 4' 55''$  and suddenly falls to  $60^\circ$  when  $e$  becomes 0.


Corol. If  $l = (a-b) \cos \phi = (a+b) \tan \theta$  then  $\frac{\pi l}{a+b}$  will be the perimeter of the ellipse when  $\phi$  diminishes from  $30^\circ$  to  $0^\circ$  when  $e$  increases from 0 to 1.  
 $\phi = \frac{2\sqrt{ab}}{a+b} \left\{30^\circ + 6^\circ 18' 8'' \frac{(a-b)^2}{a+b} - 1^\circ 10' 9'' \left(\frac{a-b}{a+b}\right)^2\right\}$

Prob. 7. Draw AN perp to AC.

Make CP & CQ equal to CB.

Draw QM making an L  $\phi$  with AQ & meeting AN at M.

Join PM & make HPN equal to  $\frac{1}{2}$  of APN. With P as centre and PA as radius describe a circle cutting PN at K & PB produced at L.



Then  $\frac{\text{arc AL}}{\text{arc AK}} = \frac{\text{arc AB}}{\text{AN}}$ .  $\phi = 30^\circ$  very nearly

$\phi = 30^\circ + h(a-b) \left\{5^\circ 19' 4'' - 6^\circ 3' 5'' h\right\}$  where  $h = \left(\frac{a-b}{a+b}\right)^2$ .

N.B. i.  $\phi = 30^\circ$  when  $e = 0, 1$  or  $.99948$ .

ii when  $e = .999886$ ,  $\phi$  assumes the minimum value of  $29^\circ 58' \frac{1}{2}$  and when  $e = .9689$ ,  $\phi$  has the maximum value of  $30^\circ 44' \frac{1}{4}$ .

20. i. To Construct a square equal to a given circle.

Let O be the centre and PR any diameter.

Bisect OP at H and trisect OR at T. Draw TQ perp to OR.


Draw  $RS \perp TR$ . Join  $PS$ .

Draw  $CD \parallel TN \parallel$  to  $RS$ .

Draw  $PL = PM$ ; &  $PL = MN$  and  
perp to  $PS$ . Join  $RL, RK$  &  $KL$   
Cut off  $PC = RH$ . Draw  $CD \parallel$  to  $KL$

Then  $\odot D^c = \odot PQR$ .

N.B.  $RD$  is  $\frac{1}{100}$ th of an inch  
greater than the true length  
if the given  $\odot$  is 14 Sq. miles  
in area.



Cor. 1. One of the two mean No  
-portionals between a side of an equilateral  
triangle inscribed in the  $\odot$  and the length  $PS$  is only  
less by  $\frac{1}{30000}$ th part of it than the true length.

Cor. 2. The app. length got by assuming  $\pi = \sqrt[4]{972} = \frac{5}{11}$   
is  $\frac{1}{100}$ th of an inch less than the true length if the  $\odot$   
is a million square miles in area.

ii.  $\{6n^2 + (2n^2 - n)\}^3 + \{6n^2 - (2n^2 - n)\}^3 = \{6n^2(3n^2 + 1)\}^2$

iii.  $\{m^7 - 3m^4(1+p) + m(3+3p^2-1)\}^3$   
 $+ \{2m^6 - 3m^3(1+2p) + (1+3p+3p^2)\}^3$   
 $+ \{m^6 - (1+3p+3p^2)\}^3 = \{m^7 - 3m^4p + m(3p^2-1)\}^3$

ex.  $(11\frac{1}{2})^3 + (1\frac{1}{2})^3 = 39^2$ ;  $(3 - \frac{1}{105})^3 + (\frac{1}{105})^3 = (\frac{57}{35})^2$   
 $(3\frac{1}{2})^3 - (\frac{1}{2})^3 = (5\frac{1}{2})^2$ ;  $(3 - \frac{1}{102})^3 - (\frac{1}{102})^3 = (5\frac{2}{102})^2$

$3^3 + 4^3 + 5^3 = 6^3$ ;  $1^3 - 2^3 + 2^3 = 9^3 + 10^3$ ;  $1^3 + 75^3 = (75\frac{1}{2})^3 + (1\frac{1}{2})^3$   
 $3^3 + 509^3 + 34^6 = 1188^3$ ;  $18^3 + 19^3 + 21^3 = 28^3$   
 $7^3 + 14^3 + 17^3 = 20^3$ ;  $19^3 + 60^3 + 69^3 = 82^3$ ;  $15^3 + 92^3 + 87^3$   
 $= 102^3$ ;  $3^3 + 36^3 + 37^3 = 36^3$ ;  $1^3 + 126^3 + 138^3 = 172^3$

$$23^3 + 134^3 = 95^3 + 116^3; 133^3 + 174^3 = 45^3 + 196^3,$$

$$1^3 + 6^3 + 8^3 = 7^3; 11^3 + 37^3 = 298^2; 71^3 - 23^3 = 588^2$$

$$21. i. \frac{1}{2\pi\sqrt{3}x^4} + \frac{1}{1^4 + 1^2x^2 + x^4} + \frac{2}{2^4 + 2^2x^2 + x^4} + \frac{3}{3^4 + 3^2x^2 + x^4} + \dots$$

$$= \frac{\pi}{3x^2\sqrt{3}} \frac{\cosh \pi x\sqrt{3} + 2\cosh \pi x}{\cosh \pi x\sqrt{3} - \cos \pi x} + 2 \left\{ \frac{1}{e^{\pi\sqrt{3}/2} + 1} \cdot \frac{1}{1^4 + 1^2x^2 + x^4} \right.$$

$$\left. - \frac{2}{e^{2\pi\sqrt{3}} - 1} \cdot \frac{1}{2^4 + 2^2x^2 + x^4} + \frac{3}{e^{3\pi\sqrt{3}} + 1} \cdot \frac{1}{3^4 + 3^2x^2 + x^4} - \dots \right\}$$

$$ii. \frac{\sqrt{3}}{2\pi x^4} + \frac{1}{1^4 + 1^2x^2 + x^4} + \frac{2}{2^4 + 2^2x^2 + x^4} + \frac{3}{3^4 + 3^2x^2 + x^4} + \dots$$

$$= \frac{\pi}{3x^2\sqrt{3}} \frac{\cosh \pi x\sqrt{3} + 2\cosh \pi x + 6\cosh \frac{\pi x}{\sqrt{3}}}{\cosh \pi x\sqrt{3} - \cos \pi x} + 2 \left\{ \frac{1}{e^{\pi/\sqrt{3}} + 1} \cdot \frac{1}{1^4 + 1^2x^2 + x^4} \right.$$

$$\left. - \frac{2}{e^{2\pi/\sqrt{3}} - 1} \cdot \frac{1}{2^4 + 2^2x^2 + x^4} + \frac{3}{e^{3\pi/\sqrt{3}} + 1} \cdot \frac{1}{3^4 + 3^2x^2 + x^4} - \dots \right\}$$

$$iii. \frac{1}{2n^2} + \frac{1}{1^2 + n^2 + n^2} + \frac{1}{2^2 + 2n^2 + n^2} + \frac{1}{3^2 + 3n^2 + n^2} + \dots$$

$$+ 2n \left\{ \frac{1}{e^{\pi n\sqrt{3}} + 1} \cdot \frac{1}{1^4 + 1^2n^2 + n^4} - \frac{2}{e^{2\pi n\sqrt{3}} - 1} \cdot \frac{1}{2^4 + 2^2n^2 + n^4} + \dots \right\}$$

$$= \frac{1}{2\pi n^3\sqrt{3}} + \frac{2\pi}{3n\sqrt{3}} - \frac{2\pi}{n\sqrt{3}} \cdot \frac{1}{e^{2\pi n\sqrt{3}} - 2e^{\pi n\sqrt{3}} \cos \pi n + 1}$$

$$iv. \frac{1}{6n^2} + \frac{1}{1^2 + 3n^2 + 3n^2} + \frac{1}{2^2 + 6n^2 + 3n^2} + \frac{1}{3^2 + 9n^2 + 3n^2} + \dots$$

$$+ 6n \left\{ \frac{1}{e^{\pi n\sqrt{3}} + 1} \cdot \frac{1}{1^4 - 3n^2 + 9n^4} - \frac{2}{e^{2\pi n\sqrt{3}} - 1} \cdot \frac{1}{2^4 - 6n^2 + 9n^4} + \dots \right\}$$

$$= \frac{1}{6\pi n^2\sqrt{3}} + \frac{\pi}{3n\sqrt{3}} - \frac{2\pi}{n\sqrt{3}} \cdot \frac{1}{e^{2\pi n\sqrt{3}} - 2e^{\pi n\sqrt{3}} \cos 2\pi n + 1}$$

$$ex. \frac{1}{7 \cdot 13 (e^{\pi\sqrt{3}} + 1)} - \frac{2}{7 \cdot 19 (e^{2\pi\sqrt{3}} - 1)} + \frac{3}{9 \cdot 27 (e^{3\pi\sqrt{3}} + 1)}$$

$$- \frac{4}{13 \cdot 37 (e^{4\pi\sqrt{3}} - 1)} + \dots = \frac{1}{324\pi\sqrt{3}} + \frac{25}{756} - \frac{\pi}{54\sqrt{3}}$$

$$+ \frac{\pi}{18\sqrt{3}} \cdot \frac{1}{1 + \cosh 3\pi\sqrt{3}}$$

N.B. The series  $\frac{1}{1^2 + 9n^2 + 9n^2} + \frac{1}{2^2 + 9n^2 + 9n^2} + \frac{1}{3^2 + 9n^2 + 9n^2} + \dots$   
 + &c can be exactly found if n is any integer  
 and of any quantity.

$$\int_0^{\pi} \frac{\phi d\theta}{\sqrt{1-x\sin^2\theta}} d\phi = \frac{1}{n} + \frac{x}{n} + \frac{4x}{n} + \frac{9x}{n} + \frac{16}{n} + \dots$$

$$\int_0^{\pi} \frac{\phi d\theta}{\sqrt{1-x\sin^2\theta}} \frac{\cos\phi}{\sqrt{1-x\sin^2\phi}} d\phi = \frac{1}{n} + \frac{1}{n} + \frac{4x}{n} + \frac{9}{n} + \frac{16x}{n} + \dots$$

$$\int_0^{\pi} e^{-n\phi} \frac{d\theta}{\sqrt{1-x\sin^2\theta}} \frac{\cos\phi}{1-x\sin^2\phi} d\phi = \frac{1}{n} + \frac{1-x}{n} - \frac{4x}{n} + \frac{9(1-x)}{n} + \dots$$

23. i  $\sqrt{x} \left\{ \frac{1}{2} + e^{-\frac{\pi x}{x^2+y^2}} \cos \frac{\pi y}{x^2+y^2} + e^{-\frac{4\pi x}{x^2+y^2}} \cos \frac{4\pi y}{x^2+y^2} + e^{-\frac{9\pi x}{x^2+y^2}} \cos \frac{9\pi y}{x^2+y^2} + \dots \right\}$

$$= \sqrt{x^2+y^2+x} \left\{ \frac{1}{2} + e^{-\pi x} \cos \pi y + e^{-4\pi x} \cos 4\pi y + e^{-9\pi x} \cos 9\pi y + \dots \right\}$$

$$+ \sqrt{x^2+y^2-x} \left\{ e^{-\pi x} \sin \pi y + e^{-4\pi x} \sin 4\pi y + e^{-9\pi x} \sin 9\pi y + \dots \right\}$$

ii.  $\sqrt{x} \left\{ e^{-\frac{\pi x}{x^2+y^2}} \sin \frac{\pi y}{x^2+y^2} + e^{-\frac{4\pi x}{x^2+y^2}} \sin \frac{4\pi y}{x^2+y^2} + \dots \right\}$

$$= \sqrt{x^2+y^2+x} \left\{ \frac{1}{2} + e^{-\pi x} \cos \pi y + e^{-4\pi x} \cos 4\pi y + e^{-9\pi x} \cos 9\pi y + \dots \right\}$$


$$- \sqrt{x^2+y^2-x} \left\{ e^{-\pi x} \sin \pi y + e^{-4\pi x} \sin 4\pi y + e^{-9\pi x} \sin 9\pi y + \dots \right\}$$

cos.  $\frac{1}{2} + e^{-\pi x} \cos \pi \sqrt{1-x} + e^{-4\pi x} \cos 4\pi \sqrt{1-x} + \dots$

$$= \frac{\sqrt{2+\sqrt{1+x}}}{\sqrt{1-x}} \left\{ e^{-\pi x} \sin \pi \sqrt{1-x} + e^{-4\pi x} \sin 4\pi \sqrt{1-x} + \dots \right\}$$

ex.  $\phi(e^{-\pi}) = \phi(e^{-5\pi}, \sqrt{5}\sqrt{5}-10); (\sqrt{5}+\sqrt{3}) \phi(e^{-\pi/5}) = (3+\sqrt{5}) \phi(e^{-2\pi})$

24. i. Let  $\angle TMM'$  be any angle. On  $MM'$  desc. a semi-c. Cutting the bisector of  $\angle TMM'$  at R. Draw  $RP$  perp to  $MM'$ . On  $MP$  desc. a semi-c. In it desc. a chord  $PQ$  equal to  $PM$ . Join  $Q$ . Let  $S$  be the middle point of  $MM'$ .



ii. If  $RP$  divides  $MM'$  in medial sec. then the MC coincides with MR.

A pendulum oscillating through  $4A$  takes  $\frac{MM'}{MP}$  to make the time required through  $4B$ . Let  $\sin^2 A = \alpha$  &  $\sin^2 B = \beta$

& Let  $\frac{MM'}{MP} = m$  then  $2PS = m \cos A$  &  $m = \frac{1 + (\frac{1}{2})^2 \alpha + (\frac{1}{2})^4 \alpha^2 + \dots}{1 + (\frac{1}{2})^2 \beta + (\frac{1}{2})^4 \beta^2 + \dots}$

N.B. Here  $\beta$  is in the second degree of  $\alpha$ .

ii. 2nd degree: -  $m\sqrt{1-\alpha} + \sqrt{\beta} = 1$  and  $m^2\sqrt{1-\alpha} + \beta = 1$ ;

$$\frac{m^2}{2} = \frac{1+\sqrt{\beta}}{1+\sqrt{1-\alpha}} = \frac{1+\beta}{1+(\frac{1}{2})\alpha}$$

iii. 4th degree: -  $\sqrt{m}\sqrt[4]{1-\alpha} + \sqrt[4]{\beta} = 1$  and  $m^2\sqrt{1-\alpha} + \sqrt{\beta} = 1$ ;

$$\frac{m}{2} = \frac{1+\sqrt[4]{\beta}}{1+\sqrt[4]{1-\alpha}} = \frac{1+\sqrt{\beta}}{1+\sqrt{1-\alpha}}$$

iv. 8th degree: -  $\sqrt{m}\sqrt[8]{1-\alpha} + \sqrt[8]{\beta} = 1$

$$16\text{th degree: - } \frac{\sqrt{m}}{2} = \frac{1+\sqrt[8]{\beta}}{1+\sqrt[8]{1-\alpha}}$$

v. If any equation  $\alpha$  may be changed to  $1-\beta$ ,  $\beta$  to  $1-\alpha$  and  $m$  to  $n/m$  where  $n$  is the degree of  $\beta$ ; thus we see that

2nd degree: -  $\frac{2}{m}\sqrt{\beta} + \sqrt{1-\alpha} = 1$  and  $(1-\sqrt{1-\alpha})(1-\sqrt{\beta}) = 2\sqrt{\beta}(1-\alpha)$

4th degree: -  $\frac{2}{\sqrt{m}}\sqrt[4]{\beta} + \sqrt[4]{1-\alpha} = 1$  and  $(1-\sqrt[4]{1-\alpha})(1-\sqrt[4]{\beta}) = 2\sqrt[4]{\beta}(1-\alpha)$

8th degree: -  $\frac{2\sqrt{2}}{\sqrt{m}}\sqrt[8]{\beta} + \sqrt[8]{1-\alpha} = 1$  and  $(1-\sqrt[8]{1-\alpha})(1-\sqrt[8]{\beta}) = 2\sqrt{2}\sqrt[8]{\beta}(1-\alpha)$

vi.  $n\pi \cdot \frac{1 + (\frac{1}{2})^n (1-\alpha) + (\frac{1}{2})^{2n} (1-\alpha)^2 + \dots}{1 + (\frac{1}{2})^n \alpha + (\frac{1}{2})^{2n} \alpha^2 + \dots} = \pi \cdot \frac{1 + (\frac{1}{2})^n (1-\beta) + (\frac{1}{2})^{2n} (1-\beta)^2 + \dots}{1 + (\frac{1}{2})^n \beta + (\frac{1}{2})^{2n} \beta^2 + \dots}$

Differentiating both sides we have,

$$n \frac{d\alpha}{d\beta} = \frac{\alpha(1-\alpha)}{\beta(1-\beta)} m^2. \text{ Again by differentiating any equa-}$$

-tion we know  $\frac{d\alpha}{d\beta}$  and hence  $m$  is known.

vii. Equations in terms of  $\Psi$  functions can be transformed to those of  $\phi$  functions and vice versa while those of  $f$  and  $X$  functions remains unchanged. e.g. the identity

$$\frac{\Psi(x^2)}{\sqrt{x} \Psi(x^2)} = 1 + \sqrt{\frac{\Psi^2(x)}{x \Psi^2(x^2)} - 1} \text{ becomes } \frac{\phi(x^2)}{\phi(x^2)} = 1 + \sqrt{\frac{\phi^2(x)}{\phi^2(x^2)} - 1}.$$

$$1. \text{ i. } \psi(x) = \frac{x^4}{1 + \frac{x}{1+x} + \frac{x^2}{1+x^2} + \frac{x^3}{1+x^3} + \frac{x^4}{1+x^4} + \dots}$$

$$\text{ii. } \psi(x) = \sqrt{x} \cdot \frac{f(x, -x^7)}{f(x^2, -x^7)}, \text{ then}$$

$$u = \frac{x^4}{1+x} + \frac{x^2}{1+x^2} + \frac{x^6}{1+x^5} + \frac{x^6}{1+x^7} + \frac{x^8}{1+x^9} + \dots$$

$$\frac{1}{u} - u = \frac{\phi(x^2)}{\sqrt{x} \psi(x^4)} \text{ and } \frac{1}{u} + u = \frac{\phi(0)}{\sqrt{x} \psi(6x^6)}$$

$$2. \text{ i. } f(-x, -x^4) f^3(x^{15}) = f(x^5) f(x^6, -x^9) f(x, -x^{14}) f(x^4, -x^{11});$$

$$f(x^4, -x^3) f^3(x^{15}) = f(x^2) f(x^3, -x^{11}) f(-x^7, -x^{13}) f(x^2, -x^8)$$

$$\text{ii. } f(x, -x^6) f^3(x^{21}) = f(x^7) f(x^6, -x^{15}) f(-x, -x^{20}) f(x^8, -x^{13});$$

$$f(-x^2, -x^5) f^3(x^{21}) = f(x^7) f(x^9, -x^{12}) f(x^7, -x^{17}) f(x^5, -x^{16});$$

$$f(x^3, -x^4) f^3(x^{21}) = f(x^7) f(x^3, -x^{18}) f(x^4, -x^{17}) f(x^{10}, -x^{11}).$$

and so on.

$$3. \text{ i. } x \psi(x^2) \psi(x^6) = \frac{x}{1-x^2} - \frac{x^5}{1-x^{10}} + \frac{x^7}{1-x^{14}} - \frac{x^{11}}{1-x^{22}} + \dots$$

$$\text{ii. } \phi(x) \phi(x^3) = 1 + 2 \left( \frac{x}{1-x} - \frac{x^2}{1-x^4} + \frac{x^6}{1-x^6} - \frac{x^5}{1-x^5} + \frac{x^7}{1-x^7} - \dots \right)$$

$$\text{iii. } x^2 \psi^2(x) \psi^2(x^3) = \frac{x}{1-x} + \frac{2x^2}{1-x^4} + \frac{4x^6}{1-x^6} + \frac{2x^5}{1-x^{10}} + \dots$$

$$\text{iv. } \phi^4(x) \phi^4(x^3) = 1 + 4 \left( \frac{x}{1-x} + \frac{4x^6}{1-x^6} + \frac{5x^5}{1-x^5} + \frac{7x^7}{1-x^7} + \frac{5x^8}{1-x^8} + \dots \right)$$

$$4. \text{ i. } x \psi^5(x) \psi(x^3) - 9x^4 \psi(x) \psi^5(x^3)$$

$$= \frac{x}{1-x^2} - \frac{x^2}{1-x^4} + \frac{5^2 x^6}{1-x^6} - \frac{5^2 x^{15}}{1-x^{10}} + \dots$$

$$\text{ii. } 9\phi(x) \phi^5(x^3) - \phi^5(x) \phi(x^2)$$

$$= 8 \left\{ 1 + \frac{x}{1+x} - \frac{4^2 x^2}{1-x^2} + \frac{4^2 x^6}{1-x^6} - \frac{5^2 x^{15}}{1-x^5} + \frac{7x^7}{1+x^7} - \dots \right\}$$

$$\text{iii. } \frac{\psi^3(x)}{\psi(x^3)} = 1 + 3 \left( \frac{x}{1-x} - \frac{x^5}{1-x^5} + \frac{x^7}{1-x^7} - \frac{x^{11}}{1-x^{11}} + \dots \right)$$

$$\text{iv. } \frac{\phi^3(x)}{\phi(x^3)} = 1 + 6 \left( \frac{x}{1-x} + \frac{x^2}{1+x} - \frac{x^6}{1+x^6} - \frac{x^{15}}{1-x^5} + \dots \right)$$

5. From these we get the following results.



If  $\beta$  be of the 3rd degree,

$$i. \sqrt[8]{\frac{\alpha^3}{\beta}} - \sqrt[8]{\frac{(1-\alpha)^3}{1-\beta}} = \sqrt[8]{\frac{(1-\beta)^3}{1-\alpha}} - \sqrt[8]{\frac{\beta^3}{\alpha}} = 1.$$

$$ii. \sqrt[4]{\alpha\beta} + \sqrt[4]{(1-\alpha)(1-\beta)} = 1.$$

$$iii. m = 1 + 2\sqrt[8]{\frac{\beta^3}{\alpha}} \text{ and } \frac{3}{m} = 1 + 2\sqrt[8]{\frac{(1-\alpha)^3}{1-\beta}}$$

$$iv. m^2 (\sqrt[8]{\frac{\alpha^3}{\beta}} - \alpha) = \sqrt[8]{\frac{\alpha^3}{\beta}} - \alpha.$$

$$v. m = \frac{1 - 2\sqrt[8]{\frac{\beta^3(1-\alpha)^3}{\alpha(1-\beta)}}}{1 - 2\sqrt[8]{\alpha\beta}} = \sqrt[4]{1 + 4\sqrt[8]{\frac{\beta^3(1-\alpha)^3}{\alpha(1-\beta)}}} \text{ and}$$

$$\frac{3}{m} = \frac{2\sqrt[8]{\frac{\alpha^3(1-\alpha)^3}{\beta(1-\beta)}} - 1}{1 - 2\sqrt[8]{\alpha\beta}} = \sqrt[4]{1 + 4\sqrt[8]{\frac{\alpha^3(1-\alpha)^3}{\beta(1-\beta)}}}$$

$$vi. \text{ If } \alpha = p \cdot \left(\frac{2+p}{1+3p}\right)^3 \text{ then } \beta = p^3 \cdot \frac{2+p}{1+3p}. \text{ So that}$$

$$1-\alpha = (1+p) \left(\frac{1-p}{1+3p}\right)^3 \text{ \& } 1-\beta = (1+p)^3 \cdot \frac{1-p}{1+3p}$$

$$vii. m^2 = \sqrt{\frac{\beta}{\alpha}} + \sqrt{\frac{1-\beta}{1-\alpha}} - \sqrt{\frac{\beta(1-\beta)}{\alpha(1-\alpha)}} \text{ and hence}$$

$$9/m^2 = \sqrt{\frac{\alpha}{\beta}} + \sqrt{\frac{1-\alpha}{1-\beta}} - \sqrt{\frac{\alpha(1-\alpha)}{\beta(1-\beta)}}$$

$$viii. \sqrt[8]{\alpha\beta^5} + \sqrt[8]{(1-\alpha)(1-\beta)^5} = 1 - \sqrt[8]{\frac{\beta^3(1-\alpha)^3}{\alpha(1-\beta)}} \\ = \sqrt[8]{\alpha^5\beta} + \sqrt[8]{(1-\alpha)^5(1-\beta)} = \frac{\sqrt{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}}{2}$$

$$ix. \sqrt{\alpha(1-\beta)} + \sqrt{\beta(1-\alpha)} = 2\sqrt[8]{\alpha\beta(1-\alpha)(1-\beta)}.$$

$$x. m^2 \sqrt{\alpha(1-\alpha)} + \sqrt{\beta(1-\beta)} = 9/m^2 \cdot \sqrt{\beta(1-\beta)} + \sqrt{\alpha(1-\alpha)}.$$

$$xi. m \sqrt{1-\alpha} + \sqrt{1-\beta} = \frac{3}{m} \sqrt{1-\beta} - \sqrt{1-\alpha} = 2\sqrt[8]{(1-\alpha)(1-\beta)} \text{ and} \\ m \sqrt{\alpha} - \sqrt{\beta} = \frac{3}{m} \sqrt{\beta} + \sqrt{\alpha} = 2\sqrt[8]{\alpha\beta}.$$

$$xii. m - \frac{3}{m} = 2 \left\{ \sqrt[8]{\alpha\beta} - \sqrt[8]{(1-\alpha)(1-\beta)} \right\} \text{ and}$$

$$m + \frac{3}{m} = 4 \frac{\sqrt{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}}{2}$$

$$xiii. \text{ If } P = \sqrt[8]{16\alpha\beta(1-\alpha)(1-\beta)} \text{ and } Q = \sqrt[4]{\frac{\beta(1-\beta)}{\alpha(1-\alpha)}}, \text{ then}$$

$$Q + \frac{1}{Q} + 2\sqrt{2} \left(P - \frac{1}{P}\right) = 0$$

xiii. If  $p = \frac{2 \sin \alpha}{1 + \cos \alpha}$  and  $Q = \sqrt{\frac{2}{1 + \cos \alpha}}$  then

$$Q - \frac{1}{Q} = 2 \left( P - \frac{1}{P} \right)$$

xiv. If  $\alpha = \sin^2(u+v)$  and  $\beta = \sin^2(u-v)$ , then  $\sin 2u = 2 \sin \alpha$

xv. If  $\alpha(1-\alpha) = p \cdot \left( \frac{1-p}{1+4p} \right)^3$  then  $\beta(1-\beta) = p^3 \cdot \frac{2-p}{1+4p}$

6. i.  $1 + \left(\frac{1}{2}\right)^2 p \cdot \left(\frac{1+p}{1+4p}\right)^3 + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 p^2 \cdot \left(\frac{1+p}{1+4p}\right)^6 + \dots$

$$= (1+2p) \left\{ 1 + \left(\frac{1}{2}\right)^2 p^2 \cdot \frac{1+p}{1+4p} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 p^4 \cdot \left(\frac{1+p}{1+4p}\right)^2 + \dots \right\}$$

ii  $1 + \left(\frac{1}{2}\right)^2 4p \cdot \left(\frac{1+p}{1+4p}\right)^3 + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 16p^2 \cdot \left(\frac{1+p}{1+4p}\right)^6 + \dots$

$$= \sqrt{1+4p} \left\{ 1 + \left(\frac{1}{2}\right)^2 4p^3 \cdot \frac{1+p}{1+4p} + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 16p^6 \cdot \left(\frac{1+p}{1+4p}\right)^2 + \dots \right\}$$

iii. If  $\tan \frac{A+B}{2} = (1+p) \tan A$ , then

$$(1+2p) \int_0^A \frac{d\phi}{\sqrt{1-p^2 \cdot \frac{1+p}{1+4p} \sin^2 \phi}} = \int_0^B \frac{d\phi}{\sqrt{1-p \cdot \left(\frac{1+p}{1+4p}\right)^2 \sin^2 \phi}}$$

iv. If  $\tan \frac{A-B}{2} = \frac{1-p}{1+2p} \tan B$ , then

$$(1+2p) \int_0^A \frac{d\phi}{\sqrt{1-p^2 \cdot \frac{2+p}{1+4p} \sin^2 \phi}} = 3 \int_0^B \frac{d\phi}{\sqrt{1-p \cdot \left(\frac{1+p}{1+2p}\right)^2 \sin^2 \phi}}$$

v. If  $\tan \frac{A+B}{2} = \frac{2 \tan B + 2 \tan^3 B (1-x)}{1 - \tan^4 B (1-x)}$  then

$$\int_0^A \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} = 3 \int_0^B \frac{d\phi}{\sqrt{1-x \sin^2 \phi}}$$

7. If  $\alpha = p \cdot \left(\frac{1+p}{1+2p}\right)^3$  and  $\gamma = 1 + \left(\frac{1}{2}\right)^2 x + \left(\frac{1 \cdot 3}{2 \cdot 4}\right)^2 x^2 + \dots$

i. If  $\cos A = \frac{1-p}{2+p}$  then  $\int_0^A \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} = \frac{\pi}{3} \gamma$

ii. If  $\sin A = \frac{1+2p}{2+p}$ , then  $\int_0^A \frac{d\phi}{\sqrt{1-x \sin^2 \phi}} = \frac{\pi}{6} \gamma$

iii. PA is any diameter of a circle whose center is O. Draw TB any perp to AP and PR & PR, equal to TB. Join AB, AR & AR. Then a pendulum A oscillating through B, AR, takes  $\frac{AR + OT}{AO}$  or  $\frac{1+3AO}{AR+OT}$  times the time required to oscillate through B, AR.



Cor. If T coincides with O,  $\angle BAR = 15^\circ$  &  $\angle BAR_1 = 75^\circ$  and so that  $\frac{AR}{AO}$  or  $\frac{3AO}{AR+OT} = \sqrt{3}$  that is  
 A pendulum oscillating through  $60^\circ$  takes  $\sqrt{3}$  times the time required to oscillate through  $60^\circ$ .

IV. Let AQP be any  $\odot$ .

Let AP & PQ be a diameter and a chord.

Let B be the middle point of the arc PA.

Join AB & PB

Draw  $AB_1$  &  $PB_1$ , equal to AB & PB respectively.

Draw  $PR$  &  $PR_1$  equal to  $\frac{1}{2}PQ$ . Join AR &

$AR_1$ , cutting PB &  $PB_1$  at  $C$  &  $C_1$ , respectively.

Produce AB &  $AB_1$  to meet the tangent at P at  $M$  &  $M_1$ , respectively. Produce BP &

$BR_1$  to meet at  $C_2$  and produce  $B_1P$  &  $AR$  at  $C_3$ .

Then a  $\odot$  will pass through  $M, C, C_1, M_1, C_2$  and  $C_3$  and this  $\odot$  will be orthogonal to the  $\odot$  APB


and touch the st. lines AB &  $AB_1$  at  $M$  &  $M_1$ .

Let O be the centre of the new  $\odot$ . Join  $OM$  &  $OM_1$ , and  $QR$  &  $QR_1$ .

The  $\odot$   $MC_2M_1$ , passes also through the intersections of the  $\odot$  whose centres are A & P and radii AB & PA respectively.

The distances of any pt. on the  $\odot$  of  $MC_2M_1$ , from A & P bear a constant ratio:  $QR \cdot QR_1 = 3RP^2$ .

A pendulum oscillating through 4 times  $PAR$ , takes  $\frac{QR}{RP}$  or  $\frac{3RP}{QR}$  times the time required to oscillate through 4 times  $PAR$ .



v. The result on page 236 can be proved geometrically as follows:  
 $\sqrt{a} = \frac{BC}{AC_2}$ ;  $\sqrt{b} = \frac{BC}{AC_1}$ ;  $\sqrt{1-a} = \frac{AB}{AC_2}$ ;  $\sqrt{1-b} = \frac{AB}{AC_1}$   
 $\sqrt{ab} = \frac{BC \cdot AB}{AC_1 \cdot AC_2} = \sqrt{\frac{BM}{AM}} = \frac{BP}{AP}$ , similarly  $\sqrt{(1-a)(1-b)} = \sqrt{\frac{AP}{BP}}$   
 $\therefore m = \frac{AB}{BP}$  and  $\frac{3}{m} = \frac{AP}{BP}$ .

(i)  $\sqrt{a} + \sqrt{(1-a)(1-b)} = \frac{BM}{AM} + \frac{AB}{AM} = 1$ .

(ii)  $\sqrt{\frac{a}{1-a}} - \sqrt{\frac{(1-a)^2}{1-a}} = \frac{\sqrt{a}}{\sqrt{1-a}} - \frac{\sqrt{1-a}}{\sqrt{(1-a)(1-b)}} = \frac{BC_2}{BP} \cdot \frac{AP}{AC_2} - \frac{AP}{AC_2}$   
 $= \frac{PC_2}{BP} \cdot \frac{AP}{AC_2} = \frac{PC_2}{AC_2} \cdot \frac{AM}{PM} = 1$ .

(iii)  $\sqrt{\frac{1-a}{1-a}} - \sqrt{\frac{a^2}{a}} = \frac{\sqrt{1-a}}{\sqrt{(1-a)(1-b)}} - \frac{\sqrt{a}}{\sqrt{1-a}} = \frac{AP}{AC_2} - \frac{BC_2}{AP} \cdot \frac{AP}{BP}$   
 $= \frac{AP}{AC_2} \cdot \frac{CP}{BP} - \frac{CP}{AC_2} \cdot \frac{AM}{PM} = 1$ .

and so on.

8. i.  $x\psi^3(x) - 5x^2\psi(x)\psi'(x)$

$$= \frac{x}{1-x} - \frac{2x^2}{1-x^2} - \frac{3x^3}{1-x^3} + \frac{4x^4}{1-x^4} + \frac{6x^6}{1-x^6} - \dots$$

ii.  $5\phi(x)\phi^3(x^5) - \phi^5(x)$

$$= 4 \left\{ 1 + \frac{x}{1+x} - \frac{2x^2}{1-x^2} - \frac{3x^3}{1+x^3} + \frac{4x^4}{1-x^4} + \frac{6x^6}{1-x^6} - \dots \right\}$$

iii.  $25\phi(x)\phi^3(x^5) - \phi^5(x)$

$$= 24 + 40 \left( \frac{x}{1+x} - \frac{3x^2}{1+x^3} - \frac{7x^7}{1+x^7} + \frac{9x^9}{1+x^9} + \dots \right)$$

iv.  $\frac{\psi^5(x)}{\psi(x^5)} - 25x^2\psi(x)\psi'(x)$

$$= 1 + 5 \left( \frac{x}{1+x} - \frac{2x^2}{1+x^2} - \frac{3x^3}{1+x^3} + \frac{4x^4}{1+x^4} + \dots \right)$$

9. i.  $\frac{f^5(x)}{f(x^5)} = 1 - 5 \left( \frac{x}{1+x} - \frac{3x^2}{1+x^3} + \frac{4x^4}{1+x^4} - \frac{7x^7}{1+x^7} + \frac{9x^9}{1+x^9} \right.$   
 $\left. + \frac{11x^{11}}{1+x^{11}} - \frac{12x^{12}}{1+x^{12}} - \dots \right)$

ii.  $4x \frac{f^5(x^5)}{f(x)} \neq \frac{f^5(x^5)}{f(x)} = \phi(x)\phi^3(x^5)$ .

iii.  $\phi^4(x) - \phi^4(x^5) = 4x\chi(x)f(x^5)f(x^{20})$ .

$$iv. \left\{ \phi(x^5) + 2x^{\frac{1}{2}} f(x^2, x^7) \right\}^2 + \left\{ \phi(x^5) + 2x^{\frac{1}{2}} f(x, x^9) \right\}^2$$

$$= \phi^2(x^5) - 4\phi^2(x) + 3\phi^2(x^2)$$

$$v. 1 - \frac{f^5(x)}{f(x^2)} = 5x \frac{d \log \frac{f(x^2, x^3)}{f(x, x^4)}}{dx}$$

$$vi. \frac{\psi^5(x)}{\psi(x^5)} - 25x^4 \psi(x) \psi^3(x^5) = 1 - 5x \frac{d \log \frac{f(x^2, x^9)}{f(x, x^4)}}{dx}$$

$$vii. f(x, x^4) f(x^5, x^3) = \frac{\phi(x^5) f(x^5)}{x(x^2)} \quad \& \quad f(x, x^4) f(x^5, x^3) = f(x) f(x^5) \text{ and } f(x, x^4) f(x^5, x^3) = \chi(x) f(x^5) f(x^5)$$

$$10. i. \psi(x^5) = x^{\frac{1}{2}} \psi(x^5) = f(x^5, x^3) + x^{\frac{1}{2}} f(x, x^4)$$

$$ii. \phi(x^5) - \phi(x^5) = 2x^{\frac{1}{2}} f(x^2, x^7) + 2x^{\frac{1}{2}} f(x, x^9)$$

$$iii. f(x) \{ f(x^{\frac{1}{2}}) + x^{\frac{1}{2}} f(x^5) \} = f(x^{\frac{1}{2}}, x^5) - \sqrt{x} f(x^{\frac{1}{2}}, x^9)$$

$$iv. \phi^2(x) - \phi^2(x^5) = 4x f(x, x^4) f(x^5, x^3)$$

$$v. \psi^2(x) - x \psi^2(x^5) = f(x, x^4) f(x^5, x^3)$$

$$vi. f^5(x^2, x^3) + x f^5(x, x^4) = \left\{ \frac{\psi^5(x)}{\psi(x^5)} - x \psi(x^5) \right\} x$$

$$vii. 32x f^5(x^2, x^7) + 32x^2 f^5(x, x^9) = \left\{ \frac{\phi^5(x)}{\phi(x^5)} - \phi(x^5) \right\} x$$

& hence

$$\left\{ \phi^6(x) - 4\phi^2(x) \phi^4(x^5) + 11\phi^6(x^5) \right\}$$

$$viii. f^{10}(x^2, x^3) - x f^{10}(x, x^4) = \frac{f^6(x)}{f(x^5)} + 11x f^5(x) f^5(x^5)$$

$$11. i. \phi(x^5) = \phi(x^5) + \sqrt{u} + \sqrt{v} \quad \text{where}$$

$$u + v = \frac{\phi^2(x) - \phi^2(x^5)}{\phi(x^5)} \left\{ \phi^2(x) - 4\phi^2(x) \phi^2(x^5) + 11\phi^6(x^5) \right\}$$

$$u - v = \frac{\phi^2(x) - \phi^2(x^5)}{\phi(x^5)} \left\{ 5\phi^2(x^5) - \phi^2(x) \right\} \sqrt{\phi^6(x) - 2\phi^2(x) \phi^4(x^5) + \phi^6(x^5)}$$

$$\sqrt{uv} = \phi^2(x) - \phi^2(x^5)$$

$$ii. x^{\frac{1}{2}} \psi(x^5) = x^{\frac{1}{2}} \psi(x^5) + \sqrt{u} + \sqrt{v} \quad \text{where}$$

$$u + v = x^{\frac{1}{2}} \frac{\psi^2(x) - x\psi^2(x^5)}{\psi(x^5)} \left\{ \psi^2(x) - 4x\psi^2(x)\psi^2(x^5) + 11x^2\psi^4(x^5) \right\}$$

$$u - v = x^{\frac{1}{2}} \frac{\psi^2(x) - x\psi^2(x^5)}{\psi(x^5)} \left\{ \psi^2(x) - 5x\psi^2(x^5) \right\} x$$

$$\sqrt{\psi^2(x) - 2x\psi^2(x)\psi^2(x^5) + 5x^2\psi^4(x^5)}$$

$$\sqrt[5]{uv} = x^{\frac{1}{2}} \left\{ \psi^2(x) - x\psi^2(x^5) \right\}$$

$$\text{iii. } \sqrt[5]{2u} = 11 + \frac{f^6(x)}{xf^6(x^5)} \quad \text{and } uv = 1 + \frac{f(x^5)}{x^5 f(x^5)}, \text{ then}$$

$$\sqrt[5]{\sqrt{u^2+1}-u} = \sqrt{v^2+1}-v = \frac{\sqrt{x}}{1 + \frac{x}{1} + \frac{x^2}{1} + \frac{x^3}{1} + \frac{x^4}{1} + \dots}$$

$$= x^{\frac{1}{2}} \frac{f(x, -x)}{f(x^5, -x^5)}$$

$$\text{iv. } \frac{f(-x^5)}{x^5 f(x^5)} = \sqrt[3]{5 + \sqrt{u} - \sqrt{v}} \quad \text{where } \sqrt[5]{uv} = 25 + 3 \frac{f^6(x)}{xf^6(x^5)}$$

$$\text{and } uv - u = 5^5 \cdot 11 + 75^2 \cdot \frac{f^6(x)}{xf^6(x^5)} + 15^2 \frac{f^{12}(x)}{x^2 f^{12}(x^5)} - \frac{f^{18}(x)}{x^3 f^{18}(x^5)}$$

$$\text{2. i. } 1 + 5x \frac{f(-x^5)}{f(-x)} = \sqrt{u} - \sqrt{v} \quad \text{where } uv = 1 \text{ and}$$

$$u - v = 11 + 125x \frac{f^6(x^5)}{f^6(x)}$$

$$\text{ii. } x \frac{f(-x^5)}{f(-x)} = \sqrt[3]{\frac{1 + \sqrt{u} - \sqrt{v}}{25}} \quad \text{where } \sqrt[5]{uv} = 1 + 15x \frac{f^6(x^5)}{f^6(x)}$$

$$\text{and } v - u = 11 + 15^2 x \frac{f^6(x^5)}{f^6(x)} + 5 \cdot 15^4 x^2 \frac{f^{12}(x^5)}{f^{12}(x)} - 25^2 x^3 \frac{f^{18}(x^5)}{f^{18}(x)}$$

$$\text{iii. } 5 \frac{\phi(x^5)}{\phi(x)} = 1 + \sqrt{u} + \sqrt{v} \quad \text{where } \sqrt[5]{uv} = 5 \frac{\phi^2(x^5)}{\phi^2(x)} - 1$$

$$\& u + v = \left\{ 5 \frac{\phi^2(x^5)}{\phi^2(x)} - 1 \right\} \left\{ 11 - 20 \frac{\phi^2(x^5)}{\phi^2(x)} + 25 \frac{\phi^4(x^5)}{\phi^4(x)} \right\}$$

$$\text{iv. } 5x^3 \frac{\psi(x^5)}{\psi(x)} = 1 - \sqrt{u} + \sqrt{v} \quad \text{where } \sqrt[5]{uv} = 1 - 5x \frac{\psi^2(x^5)}{\psi^2(x)}$$

$$\& u - v = \left\{ 1 - 5x \frac{\psi^2(x^5)}{\psi^2(x)} \right\} \left\{ 11 - 20x \frac{\psi^2(x^5)}{\psi^2(x)} + 25x^2 \frac{\psi^4(x^5)}{\psi^4(x)} \right\}$$

$$\text{v. } \frac{f(-x^5)}{f(x^5)} = \frac{f(x^5, -x^5)}{f(-x, -x^5)} = x^{\frac{1}{2}} - x^{\frac{3}{2}} \frac{f(x^5, -x^5)}{f(-x, -x^5)}$$

$$\text{vi. } \frac{\phi(-x^5)\phi(x^5)}{\phi^2(x)} + x^{\frac{5}{2}} \left\{ \frac{\psi(x^5)\psi(x^5)}{\psi^2(x)} + \frac{\psi(-x^5)\psi(x^5)}{\psi^2(-x)} \right\} = 1$$

13. If  $\beta$  be of the fifth degree,

$$i. \sqrt{a\beta} + \sqrt{(1-a)(1-\beta)} + 2\sqrt[5]{16a\beta(1-a)(1-\beta)} = 1$$

$$ii. \sqrt[5]{\frac{a^5}{\beta}} - \sqrt[5]{\frac{(1-a)^5}{1-\beta}} = 1 + \sqrt[5]{2} \sqrt[5]{\frac{a^5(1-a)^5}{\beta(1-\beta)}}$$

$$iii. \sqrt[5]{\frac{(1-a)^5}{1-a}} - \sqrt[5]{\frac{a^5}{a}} = 1 + \sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-a)}}$$

$$iv. m = 1 + 2\sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-a)}} \quad \& \quad \frac{5}{m} = 1 + 2\sqrt[5]{2} \sqrt[5]{\frac{a^5(1-a)^5}{\beta(1-\beta)}}$$

$$v. m = \frac{1 + \sqrt[5]{\frac{a^5(1-a)^5}{1-a}}}{1 + \sqrt[5]{\frac{(1-a)^5(1-\beta)}{1-a}}} = \frac{1 - \sqrt[5]{\frac{a^5}{a}}}{1 - \sqrt[5]{\frac{a^5}{\beta}}}$$

$$vi. \frac{5}{m} = \frac{1 + \sqrt[5]{\frac{a^5}{\beta}}}{1 + \sqrt[5]{\frac{(1-a)^5}{a\beta}}} = \frac{1 - \sqrt[5]{\frac{(1-a)^5}{1-\beta}}}{1 - \sqrt[5]{\frac{(1-a)^5(1-\beta)^3}{\beta(1-\beta)^3}}}$$

$$vii. \sqrt[5]{a\beta^2} + \sqrt[5]{(1-a)(1-\beta)^3} = 1 - \sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-\beta)}} =$$

$$\sqrt[5]{a^3\beta} + \sqrt[5]{(1-a)^3(1-\beta)} = \frac{\sqrt{1 + \sqrt{2a} + \sqrt{(1-a)(1-\beta)}}}{2}$$

viii. For all values of  $a$  and  $b$

$$m = \frac{a + 2(a-b) \sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-a)}} + 6 \sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-a)}}}{a - 6 \sqrt[5]{16a\beta(1-a)(1-\beta)}}$$

$$= \frac{1 - \sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-a)}} - \sqrt[5]{2} \sqrt[5]{\frac{a^5(1-a)^5}{a(1-a)}}}{\sqrt{1 - 3\sqrt[5]{16a\beta(1-a)(1-\beta)}} + \sqrt[5]{16a\beta(1-a)(1-\beta)}}$$

$$ix. 1 + \sqrt[5]{2} \sqrt[5]{\frac{\beta^5(1-a)^5}{a(1-a)}} = m \cdot \frac{1 + \sqrt{a\beta} + \sqrt{(1-a)(1-\beta)}}{2} \quad \&$$

$$1 + \sqrt[5]{2} \sqrt[5]{\frac{a^5(1-a)^5}{a(1-a)}} = \frac{5}{m} \cdot \frac{1 + \sqrt{a\beta} + \sqrt{(1-a)(1-\beta)}}{2}$$

$$x. \sqrt[5]{a(1-a)} + \sqrt[5]{\beta(1-a)} = \sqrt[5]{2} \sqrt[5]{a\beta(1-a)(1-\beta)} \quad \times$$

$$= m \sqrt[5]{a(1-a)} + \frac{5}{m} \sqrt[5]{\beta(1-a)}$$

$$xi. \sqrt[5]{\frac{(1-a)^5}{1-a}} + \sqrt[5]{\frac{a^5}{a}} = m \frac{\sqrt{1 + \sqrt{a\beta} + \sqrt{(1-a)(1-\beta)}}}{2} \quad \text{and}$$

$$\sqrt[5]{\frac{(1-a)^5}{1-\beta}} + \sqrt[5]{\frac{a^5}{\beta}} = \frac{5}{m} \frac{\sqrt{1 + \sqrt{a\beta} + \sqrt{(1-a)(1-\beta)}}}{2}$$

xii.  $m = \sqrt[3]{\frac{1-a}{1-a}} + \sqrt[3]{\frac{1-a}{1-a}} - \sqrt[3]{\frac{\beta(1-a)}{\alpha(1-a)}}$  and hence

$$\frac{5}{m} = \sqrt[3]{\frac{1-a}{1-a}} + \sqrt[3]{\frac{1-a}{1-a}} - \sqrt[3]{\frac{\alpha(1-a)}{\beta(1-a)}}$$

xiii.  $m - \frac{5}{m} = 4 \left\{ \sqrt{4\beta} - \sqrt{\alpha(1-a)\beta} \right\} / \sqrt{1 + \sqrt{4\beta} + \sqrt{\alpha(1-a)\beta}}$

and  $m + \frac{5}{m} = 2 \left\{ 2 + \sqrt{4\beta} + \sqrt{\alpha(1-a)\beta} \right\}$

xiv. If  $P = \sqrt[3]{16\alpha\beta\alpha(1-a)(1-a)}$  and  $Q = \sqrt[3]{\frac{\beta(1-a)}{\alpha(1-a)}}$ , then

$$Q + \frac{1}{Q} + 2(P - \frac{1}{P}) = 0.$$

xv. If  $P = \sqrt[3]{3\alpha}$  and  $Q = \sqrt[3]{\frac{\beta}{\alpha}}$ , then

$$(Q - \frac{1}{Q})^3 + 8(Q - \frac{1}{Q}) = 4(P - \frac{1}{P}).$$

14. i. If  $\alpha = \sin^2(u+v)$  and  $\beta = \sin^2(u-v)$ , then  $\sin 2u = \sin v(1 + \cos v)$ .

ii. If  $2\alpha(1-\alpha) = p \left( \frac{2-p}{1+p} \right)^5$  then  $\sqrt[5]{\beta(1-\beta)} = p^5 \frac{2-p}{1+p}$

iii. If  $1-2\alpha = \frac{1-11p-p^2}{(1+p)^2} \sqrt{\frac{1+p^2}{1+2p}}$  then  $1-2\beta = \frac{(1+p-p^2)}{(1+p)^2} \sqrt{\frac{1+p^2}{1+2p}}$

iv.  $1 + (\frac{1}{2})^2 \frac{1-11p-p^2}{(1+p)^2} \sqrt{\frac{1+p^2}{1+2p}} + \dots$   
 $= (1+2p) \left\{ 1 + (\frac{1}{2})^2 \frac{1-(1+p-p^2)}{2} \sqrt{\frac{1+p^2}{1+2p}} + \dots \right\}$

v.  $1 + (\frac{1}{2})^2 p \left( \frac{2-p}{1+p} \right)^5 + (\frac{1}{2} \frac{5}{8})^2 p^2 \left( \frac{2-p}{1+p} \right)^{10} + \dots$   
 $= (1+2p) \left\{ 1 + (\frac{1}{2})^2 p^5 \frac{2-p}{1+p} + (\frac{1}{2} \frac{5}{8})^2 p^{10} \left( \frac{2-p}{1+p} \right)^2 + \dots \right\}$

15. If  $\gamma$  be of the 25th degree,

i.  $\sqrt[5]{\frac{\gamma}{\alpha}} + \sqrt[5]{\frac{1-\gamma}{1-\alpha}} - \sqrt[5]{\frac{\gamma(1-\gamma)}{\alpha(1-\alpha)}} - 2 \sqrt[5]{\frac{\gamma(1-\gamma)}{\alpha(1-\alpha)}} = \sqrt{\frac{1+(\frac{1}{2})^2 \alpha + \dots}{1+(\frac{1}{2})^2 \gamma + \dots}}$

ii.  $\sqrt[5]{\frac{\alpha}{\gamma}} + \sqrt[5]{\frac{1-\alpha}{1-\gamma}} - \sqrt[5]{\frac{\alpha(1-\alpha)}{\gamma(1-\gamma)}} - 2 \sqrt[5]{\frac{\alpha(1-\alpha)}{\gamma(1-\gamma)}} = 5 \sqrt{\frac{1+(\frac{1}{2})^2 \gamma + \dots}{1+(\frac{1}{2})^2 \alpha + \dots}}$

iii.  $\sqrt[5]{\frac{\alpha\gamma}{\beta^2}} + \sqrt[5]{\frac{\beta(1-\alpha)(1-\gamma)}{(1-\beta)^2}} + \sqrt[5]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta^2(1-\beta)^2}} = \frac{1+(\frac{1}{2})^2 \beta + (\frac{1}{2} \frac{3}{2})^2 \beta^2 + \dots}{\sqrt{1+(\frac{1}{2})^2 \alpha + \dots} \sqrt{1+(\frac{1}{2})^2 \gamma + \dots}}$



$$iv. \sqrt{\frac{\beta^2}{\alpha^2}} + \sqrt{\frac{0-\beta^2}{(1-\alpha)(1-\beta)}} + \sqrt{\frac{\beta^2(1-\beta)^2}{\alpha^2(1-\alpha)(1-\beta)}} - \left\{ \sqrt{\frac{\beta^2(1-\beta)^2}{\alpha^2(1-\alpha)(1-\beta)}} \right\} + \sqrt{\frac{\beta^2}{\alpha^2}} + \frac{1}{\sqrt{\beta}}$$

$$= 5 \cdot \frac{1 + (\frac{\beta}{\alpha})^2 + \dots}{1 + (\frac{\beta}{\alpha})\beta + \dots} \cdot \frac{1 + (\frac{\beta}{\alpha})^2 + \dots}{1 + (\frac{\beta}{\alpha})\beta + \dots} \quad ?$$

$$v. \frac{1 + \sqrt{\frac{\beta^2}{\alpha^2}} \sqrt{\frac{\beta^{10}(1-\beta)^0}{\alpha^7(1-\alpha)(1-\beta)}}}{1 + \sqrt{\frac{\beta^2}{\alpha^2}} \sqrt{\frac{\alpha^5 \beta^5 (1-\alpha)^5 (1-\beta)^5}{\beta^5 (1-\beta)^5}}} = \frac{1 + (\frac{\beta}{\alpha})^2 + \dots}{1 + (\frac{\beta}{\alpha})\beta + \dots} \cdot \frac{1 + (\frac{\beta}{\alpha})^2 + \dots}{1 + (\frac{\beta}{\alpha})\beta + \dots}$$


16. i. If  $\int_0^A \frac{d\phi}{\sqrt{1-\alpha \sin^2 \phi}} = m \int_0^B \frac{d\phi}{\sqrt{1-\beta \sin^2 \phi}}$ , then

$$\tan \frac{A-\beta}{2} = \frac{p \tan B}{1 + 1 + p + \sqrt{(1+p)(1+p')}} \tan^2 B$$

ii. Let  $O$  be the centre and  $AP$  be a diameter of the  $\odot PAQ$ . Take any point  $T$  between the point of medial section & the point  $P$ .

Through  $T$  draw a perp.  $RR_1$  to  $AP$  and join  $PR, RA$  &  $AR_1$ .

Through  $N$  draw  $MN \parallel^l$  to  $PR$ ,  $M$  being the middle pt. of  $TP$ . Draw  $BC$  perp. to  $AP$  and equal  $OC = MN$ . Cut off the arc  $BQ = BP$ . Join  $AB, QR$  &  $QR_1$ .



Then if the time required by a pendulum to oscillate through  $n$  times the  $\angle BAR_1$  be  $m$  times the time required to oscillate through  $l$  times the  $\angle BAR$ , then

$$1+m = 2 \frac{QR}{RT} \text{ and } 1 + \frac{l}{m} = 2 \frac{QR_1}{R_1T} \text{ and } \frac{l}{m} - m = 8 \cdot \frac{OC}{AR}$$

NB. i. Taking  $AP=1$  we see that  $TP = \sqrt{\frac{\beta(1-\alpha)(1-\beta)}{\alpha}}$  &  $CT = \sqrt{\beta\alpha}$

and  $OC + OT = \sqrt{(1-\alpha)(1-\beta)}$  so that  $\sqrt{\beta\alpha} + \sqrt{(1-\alpha)(1-\beta)} + 2\sqrt{\beta\alpha(1-\alpha)(1-\beta)} = 1$

ii. If  $T$  be the point of medial section of  $AP$ , then  $C$  will coincide with the centre  $O$  and the ratio between the times to oscillate through  $\angle BAR$  &  $\angle BAR_1$  is  $1:\sqrt{5}$ .

$$17. i. x \psi(x) \psi(x^{-1}) = \frac{x}{1-x} - \frac{x^2}{1-x^2} + \frac{x^5}{1-x^5} + \frac{x^9}{1-x^9} + \frac{x^{11}}{1-x^{11}} - \frac{x^{13}}{1-x^{13}} + \frac{x^{17}}{1-x^{17}} - \frac{x^{17}}{1-x^{17}} + \frac{x^{23}}{1-x^{23}} + \dots$$

$$ii. \phi(x) \phi(x^{-1}) = 1 + 2 \left( \frac{x}{1-x} - \frac{x^2}{1-x^2} - \frac{x^3}{1-x^3} + \frac{x^4}{1-x^4} - \frac{x^5}{1-x^5} + \frac{x^6}{1-x^6} + \frac{x^8}{1-x^8} + \frac{x^9}{1-x^9} + \frac{x^{10}}{1-x^{10}} + \dots \right)$$

$$iii. \phi(x^4) - \phi(x^{-1}) = 2x^{1/2} f(x^5, x^9) + 2x^{3/4} f(x^3, x^{11}) + 2x^{5/4} f(x, x^{13})$$

$$iv. \psi(x^{1/2}) - x^{1/4} \psi(x^{-1}) = f(x^3, x^6) + x^{1/2} f(x^2, x^5) + x^{3/4} f(x, x^6)$$

$$v. \frac{f(-x^{1/2})}{f(-x^7)} = \frac{f(-x^4, -x^6)}{f(-x, -x^6)} - x^{1/2} \frac{f(x^3, -x^4)}{f(x^2, -x^5)} - x^{3/4} + x^{5/4} \frac{f(x, -x^6)}{f(x^3, -x^9)}$$

$$18. i. 1 + \frac{f(-x^{1/2})}{x^{1/2} f(-x^7)} = \sqrt{u} - \sqrt{v} + \sqrt{w} \text{ where}$$

$$u - v + w = 57 + 14 \frac{f^4(-x)}{x f^4(x^7)} + \frac{f^8(-x)}{x^2 f^8(x^7)}$$

$$uv - uw + vw = 289 + 126 \frac{f^6(-x)}{x f^6(x^7)} + 19 \frac{f^8(-x)}{x^2 f^8(x^7)}$$

$$uvw = 1 + \frac{f^{12}(-x)}{x^3 f^{12}(x^7)}$$

$$ii. 1 + 7x^2 \frac{f(-x^{1/2})}{f(-x)} = \sqrt{u} - \sqrt{v} + \sqrt{w} \text{ where}$$

$$u - v + w = 57 + 2 \cdot 7^3 x \frac{f^4(-x^7)}{f^4(x)} + 7^4 x^2 \frac{f^8(-x^7)}{f^8(x)}$$

$$uv - uw + vw = 289 + 18 \cdot 7^3 x \frac{f^6(-x^7)}{f^6(x)} + 19 \cdot 7^4 x^2 \frac{f^8(-x^7)}{f^8(x)}$$

$$iii. f(x, x^6) f_0(x^7, x^5) f(x^3, x^6) = \frac{f^2(-x^7)}{x f^2(x)} \phi(x^7) + 7^6 x^3 \frac{f^{12}(-x^7)}{f^{12}(x)}$$

$$iv. f(x, x^6) f(-x^5, -x^7) f(-x^3, -x^6) = f(-x) f^2(-x^7)$$

$$v. f(x, x^{13}) f(x^3, x^{11}) f(x^5, x^9) = \chi(x) \psi(-x^2) f^4(-x^{14})$$

$$vi. \text{ If } u = \frac{f^4(-x)}{x f^4(x^7)} \text{ and } v = \frac{f(-x^{1/2})}{x^{1/2} f(-x^7)}, \text{ then}$$

$$2u = 7(v^3 + 5v^2 + 7v) + (v^2 + 7v + 7) \sqrt{4v^3 + 21v^2 + 28v}$$

19. If  $\beta$  be of the 7th degree,

i.  $\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)} = 1$  so that  $\frac{\sqrt{1+\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}}{2}$

ii.  $m = \frac{1 - \sqrt[7]{\frac{\alpha^7(1-\beta)^7}{\alpha(1-\alpha)}}}{\sqrt[7]{(1-\alpha)(1-\beta)} - \sqrt[7]{\alpha}}$  and  $\frac{7}{m} = \frac{1 - \sqrt[7]{\frac{\alpha^7(1-\beta)^7}{\beta(1-\beta)}}}{\sqrt[7]{\alpha\beta} - \sqrt[7]{(1-\alpha)(1-\beta)}}$

iii.  $\sqrt[7]{\frac{(1-\beta)^7}{1-\alpha}} - \sqrt[7]{\frac{\beta^7}{\alpha}} = m \sqrt{\frac{1+\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}}$  and

$\sqrt[7]{\frac{\alpha^7}{\beta}} - \sqrt[7]{\frac{(1-\alpha)^7}{1-\beta}} = \frac{7}{m} \sqrt{\frac{1+\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}}$

iv.  $\sqrt[7]{\frac{(1-\beta)^7}{1-\alpha}} - 1 = \sqrt[7]{\alpha\beta} \left\{ \sqrt[7]{\frac{(1-\beta)^7}{1-\alpha}} - \sqrt[7]{\frac{\beta^7}{\alpha}} \right\}$  and

$\sqrt[7]{\frac{\alpha^7}{\beta}} - 1 = \sqrt[7]{(1-\alpha)(1-\beta)} \left\{ \sqrt[7]{\frac{\alpha^7}{\beta}} - \sqrt[7]{\frac{(1-\alpha)^7}{1-\beta}} \right\}$

v.  $m^2 = \sqrt{\frac{\beta}{\alpha}} + \sqrt{\frac{1-\beta}{1-\alpha}} - \sqrt{\frac{\beta(1-\beta)}{\alpha(1-\alpha)}} - \sqrt[7]{\frac{\beta(1-\beta)}{\alpha(1-\alpha)}}$  and

$\frac{49}{m^2} = \sqrt{\frac{\alpha}{\beta}} + \sqrt{\frac{1-\alpha}{1-\beta}} - \sqrt{\frac{\alpha(1-\alpha)}{\beta(1-\beta)}} - \sqrt[7]{\frac{\alpha(1-\alpha)}{\beta(1-\beta)}}$

vi.  $\sqrt[7]{\frac{(1-\beta)^3}{1-\alpha}} + \sqrt[7]{\frac{\beta^3}{\alpha}} - \sqrt[7]{\frac{\beta^3(1-\beta)^3}{\alpha(1-\alpha)}} = m^2 \cdot \frac{1+\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}$

$\sqrt[7]{\frac{(1-\alpha)^3}{1-\beta}} + \sqrt[7]{\frac{\alpha^3}{\beta}} - \sqrt[7]{\frac{\alpha^3(1-\alpha)^3}{\beta(1-\beta)}} = \frac{49}{m^2} \cdot \frac{1+\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}$

vii.  $\sqrt[7]{\frac{(1-\beta)^7}{1-\alpha}} + \sqrt[7]{\frac{\beta^7}{\alpha}} + 2\sqrt[7]{\frac{\beta^7(1-\beta)^7}{\alpha(1-\alpha)}} = \frac{3}{4} + \frac{m^2}{4}$  and

$\sqrt[7]{\frac{(1-\alpha)^7}{1-\beta}} + \sqrt[7]{\frac{\alpha^7}{\beta}} + 2\sqrt[7]{\frac{\alpha^7(1-\alpha)^7}{\beta(1-\beta)}} = \frac{3}{4} + \frac{49}{4m^2}$

viii.  $m - \frac{7}{m} = 2(\sqrt{\alpha\beta} - \sqrt{(1-\alpha)(1-\beta)})(2 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)})$

ix. If  $P = \sqrt[7]{\frac{\beta(1-\beta)}{\alpha(1-\alpha)}}$  and  $Q = \sqrt[7]{\frac{\alpha(1-\alpha)}{\beta(1-\beta)}}$ , then

$Q + \frac{1}{Q} + 7 = 2\sqrt{2}(P + \frac{1}{P})$

x. If  $P = \sqrt{\alpha\beta}$  and  $Q = \sqrt{\frac{\beta}{\alpha}}$ , then

$P + \frac{1}{P} = Q + \frac{1}{Q} + (\frac{2}{P} - \frac{1}{Q})^2$

xi. If  $\alpha = \sin^2(u+v)$  &  $\beta = \sin^2(u-v)$ , then  $\cos 2u = (2\cos v - 1)\sqrt{4\cos v - 3}$

- i. Let  $v = \sqrt[3]{x} \cdot \frac{\chi(x)}{\chi^3(x^3)} = \frac{\sqrt[3]{x}}{1 + \frac{x+x^2}{1 + \frac{x^2+x^4}{1 + \frac{x^4+x^6}{1 + \frac{x^6+x^8}{1 + \dots}}}}$  then  
 $1 + \frac{1}{v} = \frac{\psi(x^3)}{x^3 \psi(x^3)}$  &  $1 + \frac{1}{v^3} = \frac{\psi^3(x)}{x \psi^3(x^3)}$
- ii.  $1 + \frac{\psi(-x^3)}{2x \psi(x^3)} = \sqrt[3]{1 + \frac{\psi^3(x)}{2 \psi^3(x^3)}}$  and  $2v = 1 - \frac{\phi(x^3)}{\phi(x)}$
- iii.  $\frac{\phi(x^3)}{\phi(x^9)} = 1 + \frac{\sqrt[3]{\frac{\phi^3(x)}{\phi^3(x^3)} - 1}}{3}$  and  $\frac{1}{\cos 40} + \frac{1}{\cos 80} = \frac{1}{\cos 20} + 6$   
 $\frac{\phi(x^9)}{\phi(x)} = 1 + \frac{\sqrt[3]{9 \frac{\phi^3(x^3)}{\phi^3(x^9)} - 1}}{3}$
- iv.  $3 + \frac{f^3(x^3)}{x^3 f^3(x^3)} = \sqrt[3]{27 + \frac{f^6(x)}{x f^6(x^3)}}$  and  $= \frac{1}{v} + 4v^2$   
 $1 + 9x \frac{f^3(x^3)}{f^3(x^3)} = \sqrt[3]{1 + 27x \frac{f^6(x^3)}{f^6(x^3)}}$
- v.  $f^3(x^3) + 3x^3 f^3(x^3) = f(x) \left\{ 1 + 6 \left( \frac{x^6}{1-x^6} - \frac{x^6}{1-x^6} + \frac{x^6}{1-x^6} - \frac{x^6}{1-x^6} + \dots \right) \right\}$

2 i.  $\phi(x) \phi(x^3) - \phi^2(x^3) = 2x \phi(x^2) \psi(x^3) \chi(x^3)$

ii.  $\psi(x) - 3x \psi(x^3) = \frac{\phi(x)}{\chi(x^3)}$

iii.  $\phi(x) \phi(x^3) + \phi^2(x^3) = 2\psi(x) \phi(x^3) \chi(x^3)$

iv.  $\psi(x^3) - x^3 \psi(x) = f(x^2, x^5) + x^3 f(x^2, x^7) + x^5 f(x, x^8)$

v.  $f(x^3) = f(x^4, -x^5) - x^3 f(x^2, -x^7) - x^5 f(x, -x^8)$

vi.  $f(x, -x^8) f(x^2, -x^7) f(x^3, -x^5) = \frac{f(x) f^3(x^3)}{f(x^3)}$

vii.  $\frac{f(-x^4, -x^5)}{f(x^2, -x^7)} + x \frac{f(-x, -x^8)}{f(x^3, -x^5)} = \frac{f(-x^4, -x^7)}{f(x, -x^8)}$

viii.  $\frac{f(x^4, -x^5)}{f(x, -x^8)} + x \frac{f(x^2, -x^7)}{f(x^3, -x^5)} = x \frac{f(x, -x^8)}{f(x^3, -x^5)} + \frac{f^4(x^3)}{f(x) f^3(x^3)}$

ix.  $\phi(x^3) - x^3 \phi(x) = 2x^3 f(x^2, x^{11}) + 2x^5 f(x^2, x^{12}) + 2x^{11} f(x, x^{18})$

3. If  $\beta$  be of the 3rd degree and  $\gamma$  of the 9th degree then

$$i. 1 + \sqrt[3]{\frac{\alpha^3(1-\alpha)^2}{\beta(1-\beta)}} = 3\sqrt[3]{\frac{1 + (\frac{1}{3})^2\gamma + (\frac{1}{3})^2\gamma^2 + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + (\frac{1}{3})^2\alpha^2 + \beta\epsilon}}$$

$$ii. 1 + \sqrt[3]{\frac{\gamma^3(1-\gamma)^2}{\beta(1-\beta)}} = \sqrt[3]{\frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}}$$

$$iii. 1 - \sqrt[3]{\frac{\alpha^3(1-\alpha)^2}{\beta(1-\beta)}} - \sqrt[3]{\frac{\gamma^3(1-\gamma)^2}{\beta(1-\beta)}} = \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon} \cdot \frac{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}$$

$$iv. 1 - \sqrt[3]{\frac{\alpha^3(1-\alpha)^2}{\beta(1-\beta)}} = \sqrt[3]{\frac{\alpha^3(1-\alpha)^2}{\beta(1-\beta)}} - 1$$

$$= \sqrt{\frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}} \sqrt{\frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}}$$

$$v. \sqrt{\alpha\gamma} + \sqrt{(1-\alpha)(1-\gamma)} + 2\sqrt[3]{\frac{\alpha\beta(1-\beta)}{\beta(1-\beta)}} = 1 + 8\sqrt[3]{\frac{\alpha\beta(1-\beta)}{\beta(1-\beta)}} \sqrt[3]{\alpha\gamma(1-\alpha)(1-\gamma)}$$

$$vi. \sqrt[3]{\frac{\alpha(1-\alpha)}{\beta(1-\beta)}} + \sqrt[3]{\frac{\gamma(1-\gamma)}{\beta(1-\beta)}} = \sqrt[3]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta(1-\beta)}}$$

$$vii. \frac{-1 + \sqrt[3]{\frac{(1-\alpha)^3}{1-\alpha}}}{1 - \sqrt[3]{(1-\alpha)(1-\alpha)}} = \frac{1 - \sqrt[3]{\frac{\beta^3}{\alpha}}}{1 - \sqrt[3]{\alpha\beta}} = \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}$$

$$viii. 1 + \sqrt[3]{\frac{\beta^3(1-\beta)^3}{\alpha(1-\alpha)}} = \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon} \cdot \frac{\sqrt{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}}{2}$$

$$ix. 1 + \sqrt[3]{\frac{\alpha^3(1-\alpha)^3}{\beta(1-\beta)}} = 3 \cdot \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon} \cdot \frac{\sqrt{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}}{2}$$

$$x. \sqrt[3]{\frac{\gamma}{\alpha}} + \sqrt[3]{\frac{1-\gamma}{1-\alpha}} - \sqrt[3]{\frac{\gamma(1-\gamma)}{\alpha(1-\alpha)}} = \sqrt{\frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}}$$

$$xi. \sqrt[3]{\frac{\alpha}{\gamma}} + \sqrt[3]{\frac{1-\alpha}{1-\gamma}} - \sqrt[3]{\frac{\alpha(1-\alpha)}{\gamma(1-\gamma)}} = 3\sqrt{\frac{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}}$$

$$xii. \sqrt[3]{\frac{\beta^2}{\alpha\gamma}} + \sqrt[3]{\frac{(1-\beta)^2}{(1-\alpha)(1-\gamma)}} - \sqrt[3]{\frac{\beta^2(1-\beta)^2}{\alpha\gamma(1-\alpha)(1-\gamma)}}$$

$$= -3 \cdot \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon} \cdot \frac{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}$$

$$xiii. \sqrt[3]{\frac{\alpha\gamma}{\beta^2}} + \sqrt[3]{\frac{(1-\alpha)(1-\gamma)}{(1-\beta)^2}} - \sqrt[3]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta^2(1-\beta)^2}}$$

$$= \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon} \cdot \frac{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}$$

$$xiv. \frac{\sqrt[3]{\frac{\alpha\beta(1-\beta)}}{\beta(1-\beta)}}{\sqrt[3]{\alpha(1-\alpha)} - \sqrt[3]{\gamma(1-\gamma)}} = \sqrt{\frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}} \sqrt{\frac{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}}$$

$$xv. (\sqrt[3]{\alpha} - \sqrt[3]{\gamma})^4 + (\sqrt[3]{1-\alpha} - \sqrt[3]{1-\gamma})^4 = \left\{ \sqrt[3]{\alpha(1-\alpha)} - \sqrt[3]{\gamma(1-\gamma)} \right\}^4$$

$$xvi. 1 = \sqrt{\alpha\gamma} + \sqrt{(1-\alpha)(1-\gamma)} + 2\sqrt[3]{\frac{\alpha\beta(1-\beta)}{\beta(1-\beta)}} \cdot \frac{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon} \cdot \frac{1 + (\frac{1}{3})^2\gamma + \beta\epsilon}{1 + (\frac{1}{3})^2\alpha + \beta\epsilon}$$

i.  $\frac{\phi(x^{18})}{\phi(x^6)} + x \left\{ \frac{\psi(x^9)}{\psi(x^3)} - \frac{\psi(x^{27})}{\psi(x^9)} \right\} = 1$

ii.  $\frac{\phi(x^2)}{\phi(x^{12})} + \frac{1}{x} \left\{ \frac{\psi(x^3)}{\psi(x^9)} - \frac{\psi(x^9)}{\psi(x^{27})} \right\} = 3$

iii.  $\frac{\phi(x^4)\phi(x^{54})}{\phi(x^6)\phi(x^{18})} + \frac{1}{x^3} \left\{ \frac{\psi(x)\psi(x^{27})}{\psi(x^3)\psi(x^9)} + \frac{\psi(x^3)\psi(x^{27})}{\psi(x^9)\psi(x^{27})} \right\} = 1$

iv.  $\phi(x)\phi(x^{27}) - \phi(x^3)\phi(x^9) = 4x^2 f(x^6)f(x^{18}) + 4x^7 \psi(x^3)\psi(x^{54})$

5. i. If  $\alpha, \beta, \gamma, \delta$  be of the 1st, 3rd, 9th and 27th degree respectively,

ii.  $\sqrt[8]{\frac{\alpha\delta}{\beta\gamma}} + \sqrt[8]{\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}} + \sqrt[8]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} = \sqrt{\frac{1+(5)^4\beta+8c}{1+(5)^4\alpha+8c}} \sqrt{\frac{1+(5)^4\gamma+8c}{1+(5)^4\delta+8c}}$

iii.  $\sqrt[4]{\frac{\beta\gamma}{\alpha\delta}} + \sqrt[4]{\frac{(1-\beta)(1-\gamma)}{(1-\alpha)(1-\delta)}} + \sqrt[4]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} - 2\sqrt[4]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} \left\{ 1 + \sqrt[4]{\frac{\beta\gamma}{\alpha\delta}} + \sqrt[4]{\frac{(1-\beta)(1-\gamma)}{(1-\alpha)(1-\delta)}} \right\} = -3 \cdot \frac{1+(5)^4\alpha+8c}{1+(5)^4\beta+8c} \cdot \frac{1+(5)^4\delta+8c}{1+(5)^4\gamma+8c}$

iv.  $\frac{1 - \sqrt[4]{\alpha\delta} - \sqrt[4]{(1-\alpha)(1-\delta)}}{2\sqrt[8]{16\beta\gamma(1-\beta)(1-\gamma)}} = \sqrt{\frac{1+(5)^4\beta+8c}{1+(5)^4\alpha+8c}} \sqrt{\frac{1+(5)^4\gamma+8c}{1+(5)^4\delta+8c}}$

v.  $= \frac{\sqrt[4]{16\beta\gamma(1-\beta)(1-\gamma)}}{\sqrt[8]{16\beta\gamma(1-\beta)(1-\gamma)}} + \frac{\sqrt[4]{16\alpha\delta(1-\alpha)(1-\delta)}}{\sqrt[8]{16\alpha\delta(1-\alpha)(1-\delta)}}$

6. i.  $\psi(x^{11}) - x^{1/11} \psi(x^{17}) = f(x^5, x^6) + x^{1/11} f(x^5, x^7) + x^{2/11} f(x^5, x^8) + x^{3/11} f(x^5, x^9) + x^{4/11} f(x^5, x^{10})$

ii.  $\phi(x^{11}) - \phi(x^{17}) = 2x^{1/11} f(x^5, x^{13}) + 2x^{2/11} f(x^5, x^{14}) + 2x^{3/11} f(x^5, x^{15}) + 2x^{4/11} f(x^5, x^{16}) + 2x^{5/11} f(x^5, x^{17})$

iii.  $\frac{f(x^{11})}{f(x^{17})} = \frac{f(x^6, -x^7)}{f(x^5, -x^8)} - x^{1/11} \frac{f(x^6, -x^9)}{f(x^5, -x^{10})} - x^{2/11} \frac{f(x^6, -x^6)}{f(x^5, -x^8)} + x^{3/11} + x^{4/11} \frac{f(x^6, -x^6)}{f(x^5, -x^7)} - x^{5/11} \frac{f(x^6, -x^{10})}{f(x^5, -x^6)}$

iv. If  $\beta$  be of the 11th degree,

i.  $\sqrt[4]{\alpha\beta} + \sqrt[4]{(1-\alpha)(1-\beta)} + 2\sqrt[8]{16\alpha\beta(1-\alpha)(1-\beta)} = 1$

$$ii. m - \frac{11}{m} = 2 (\sqrt[3]{\alpha\beta} - \sqrt[3]{(1-\alpha)(1-\beta)}) (1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)})$$

$$iii. m + \frac{11}{m} = 4 (2 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}) \sqrt{\frac{1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}}{2}}$$

$$iv. \frac{\sqrt[3]{(1-\alpha)^3}}{\sqrt{1-\alpha}} - \frac{\sqrt[3]{\beta^3}}{\sqrt{\beta}} - \frac{\sqrt[3]{\beta^3(1-\alpha)^3}}{\alpha(1-\alpha)} = m \sqrt{\frac{1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}}{2}}$$

$$v. \frac{\sqrt[3]{\alpha^3}}{\sqrt{\alpha}} - \frac{\sqrt[3]{(1-\alpha)^3}}{\sqrt{1-\alpha}} - \frac{\sqrt[3]{\alpha^3(1-\alpha)^3}}{\beta(1-\alpha)} = \frac{11}{m} \sqrt{\frac{1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}}{2}}$$

$$vi. \frac{1}{m} \left\{ 1 + 8 \sqrt[3]{2} \sqrt[3]{\frac{\beta^{11}(1-\alpha)^{11}}{\alpha(1-\alpha)}} \right\} - \frac{m}{11} \left\{ 1 + 8 \sqrt[3]{2} \sqrt[3]{\frac{\alpha^{11}(1-\alpha)^{11}}{\beta(1-\beta)}} \right\} \\ = 2 (\sqrt[3]{\alpha\beta} - \sqrt[3]{(1-\alpha)(1-\beta)})$$

$$vii. \frac{1}{m} \left\{ 1 + 8 \sqrt[3]{2} \sqrt[3]{\frac{\alpha^{11}(1-\alpha)^{11}}{\alpha(1-\alpha)}} \right\} + \frac{m}{11} \left\{ 1 + 8 \sqrt[3]{2} \sqrt[3]{\frac{\alpha^{11}(1-\alpha)^{11}}{\beta(1-\beta)}} \right\} \\ = 4 (\sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}) \sqrt{\frac{1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}}{2}}$$

~~$$viii. \frac{3}{2} \sqrt[3]{\frac{\alpha^{11}(1-\alpha)^{11}}{\beta(1-\alpha)}} - \frac{3}{2} \sqrt[3]{\frac{\beta^{11}(1-\alpha)^{11}}{\alpha(1-\alpha)}} \\ = (3 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}) \sqrt{\frac{1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}}{2}}$$~~

~~$$ix. 4 - \frac{3}{2} \sqrt[3]{\frac{\alpha^{11}(1-\alpha)^{11}}{\beta(1-\alpha)}} - \frac{3}{2} \sqrt[3]{\frac{\beta^{11}(1-\alpha)^{11}}{\alpha(1-\beta)}} \\ = 2 \sqrt[3]{16\alpha\beta(1-\alpha)(1-\beta)} \left\{ 2 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} \right\}$$~~

$$8. i. \frac{f(-x^{1/2})}{x^{1/3} f(x^{1/2})} = \frac{f(-x^4, -x^8)}{x^8 f(x^4, -x^8)} - \frac{f(-x^6, -x^7)}{x^6 f(x^3, -x^6)} - \frac{f(-x^9, -x^{11})}{x^9 f(-x^3, -x^9)}$$

$$+ \frac{f(-x^5, -x^8)}{x^{1/3} f(x^5, -x^8)} + 1 - x^5 \frac{f(-x^3, -x^6)}{f(-x^5, -x^8)} + x^{1/3} \frac{f(-x, -x^{11})}{f(-x^6, -x^9)}$$

$$= u_1 - u_2 - u_3 + u_4 + 1 - u_5 + u_6, \text{ where}$$

$$u_1 u_2 - u_3 u_5 - u_4 u_6 = 1 + \frac{f^2(-x)}{x f^2(x^{11})}$$

$$\frac{1}{u_1 u_2} - \frac{1}{u_3 u_5} - \frac{1}{u_4 u_6} = -\frac{1}{4} - \frac{f^2(-x)}{x f^2(x^{11})}$$

$$u_2 u_3 u_4 - u_1 u_5 u_6 = 3 + \frac{f^2(-x)}{x f^2(x^{11})} \text{ \& } u_1 u_2 u_3 u_4 u_5 u_6 = 1.$$

ii.  $f(x, -x^1) f(x^2, -x^1) f(x^3, -x^1) f(x^4, -x^2) f(x^5, -x^2) f(x^6, -x^2)$   
 $= f(x) f^5(x^2)$

*f* of order of the 13th degree,

iii.  $m = \sqrt[3]{\frac{a}{2}} + \sqrt[3]{\frac{1-a}{1-a}} - \sqrt[3]{\frac{\beta(1-a)}{\alpha(1-a)}} - \sqrt[3]{\frac{\beta(1-a)}{\beta(1-a)}}$  and

iv.  $\frac{12}{21} = \sqrt[3]{\frac{a}{2}} + \sqrt[3]{\frac{1-a}{1-a}} - \sqrt[3]{\frac{\beta(1-a)}{\alpha(1-a)}} - \sqrt[3]{\frac{\beta(1-a)}{\beta(1-a)}}$

9 i.  $\psi(\alpha^3) \psi(\alpha^5) - \psi(x^2) \psi(x^7) = 2x^2 \psi(x^2) \psi(x^{20})$

ii.  $\phi(x^6) \phi(x^{10}) + 2x \psi(\alpha^2) \psi(\alpha^5) = \phi(0) \phi(15)$

iii.  $\phi(-x^6) \phi(x^{30}) + 2x^5 \psi(\alpha) \psi(\alpha^{11}) = \phi(\alpha^2) \phi(x^7)$

iv.  $\psi(\alpha) \psi(\alpha^{15}) + \psi(0) \psi(x^{15}) = 2 \psi(x^6) \psi(\alpha^{10})$

v.  $\phi(0) \phi(x^{15}) - \phi(\alpha^2) \phi(\alpha^5) = 2x f(x^2) f(x^{20}) \chi(\alpha^3) \chi(\alpha^5)$

vi.  $\phi(x) \phi(\alpha^{15}) + \phi(\alpha^2) \phi(\alpha^5) = 2 f(x^6) f(x^{10}) \chi(x) \chi(\alpha^{12})$

vii.  $\{\psi(x^2) \psi(\alpha^5) - x \psi(0) \psi(\alpha^{15})\} \phi(-x^2) \phi(-x^7)$   
 $= \{\psi(\alpha^2) \psi(\alpha^5) + x \psi(\alpha) \psi(\alpha^{15})\} \phi(x) \phi(x^{15})$   
 $= f(-x) f(x^3) f(x^5) f(-x^{15})$

10 i.  $f(x^7, -x^2) + x f(x^6, -x^{13}) = \frac{f(x^6, -x^2)}{f(-x, -x^4)} f(-x^7)$

ii.  $f(-x^4, -x^{11}) - x f(-x, -x^{14}) = \frac{f(-x, -x^4)}{f(-x^7, -x^2)} f(-x^5)$

iii.  $f(-x^7, -x^8) - x f(-x^6, -x^{13})$   
 $= f(-x^5, -x) + x^{\frac{2}{3}} f(-x^2, -x^{12})$

iv.  $\{f(-x^6, -x^{11}) + x f(-x, -x^4)\} \frac{1}{x} x^{\frac{5}{3}}$   
 $= f(-x^6, -x^7) - f(-x^{\frac{5}{3}}, -x^{\frac{2}{3}})$

v.  $x \psi(\alpha^2) \psi(\alpha^5) + x^2 \psi(\alpha) \psi(\alpha^{11}) = \frac{x}{1-x} - \frac{x^4}{1-x^4} - \frac{x^{11}}{1-x^{11}} - \frac{x^{13}}{1-x^{13}}$   
 $+ \frac{x^{17}}{1-x^{17}} + \frac{x^{19}}{1-x^{19}} + \dots$

vi.  $\phi(\alpha^2) \phi(\alpha^7) + \phi(\alpha) \phi(\alpha^{11}) = 2(1 + \frac{x}{1-x} - \frac{x^6}{1-x^6} + \frac{x^9}{1-x^9} - \frac{x^7}{1-x^7} + \dots)$



11. If  $\alpha, \beta, \gamma, \delta$  be of the 1st, 3rd, 5th & 15th degree

$$i. \sqrt[4]{\alpha\delta} + \sqrt[4]{(1-\alpha)(1-\delta)} = \sqrt{\frac{1+(\frac{1}{2})^2\beta+\alpha c}{1+(\frac{1}{2})^2\alpha+\beta c}} \sqrt{\frac{1+(\frac{1}{2})^2\gamma+\delta c}{1+(\frac{1}{2})^2\delta+\gamma c}}$$

$$ii. \sqrt[4]{\beta\gamma} + \sqrt[4]{(1-\beta)(1-\gamma)} = \sqrt{\frac{1+(\frac{1}{2})^2\alpha+\beta c}{1+(\frac{1}{2})^2\beta+\alpha c}} \sqrt{\frac{1+(\frac{1}{2})^2\delta+\gamma c}{1+(\frac{1}{2})^2\gamma+\delta c}}$$

$$= \frac{\sqrt[4]{\beta\gamma} - \sqrt[4]{\beta\gamma(1-\beta)(1-\gamma)}}{\sqrt[4]{\alpha\delta}} = \frac{\sqrt[4]{(1-\beta)(1-\gamma)} - \sqrt[4]{\beta\gamma(1-\beta)(1-\gamma)}}{\sqrt[4]{(1-\alpha)(1-\delta)}}$$

$$iii. \sqrt[4]{\alpha\delta} - \sqrt[4]{(1-\alpha)(1-\delta)} = \sqrt[4]{\beta\gamma} - \sqrt[4]{(1-\beta)(1-\gamma)}$$

$$iv. 1 + \sqrt[4]{\beta\gamma} + \sqrt[4]{(1-\beta)(1-\gamma)} = \sqrt[4]{4} \sqrt[4]{\frac{\alpha^2\gamma^2(1-\alpha)^2(1-\gamma)^2}{\alpha\delta(1-\alpha)(1-\delta)}}$$

$$v. 1 - \sqrt[4]{\alpha\delta} - \sqrt[4]{(1-\alpha)(1-\delta)} = \sqrt[4]{4} \sqrt[4]{\frac{\alpha^2\delta^2(1-\alpha)^2(1-\delta)^2}{\beta\gamma(1-\beta)(1-\gamma)}}$$

$$vi. \sqrt[4]{\alpha\delta} (\sqrt[4]{1+\sqrt{\alpha}} \sqrt[4]{1+\sqrt{\delta}} + \sqrt[4]{1-\sqrt{\alpha}} \sqrt[4]{1-\sqrt{\delta}})$$

$$+ \sqrt[4]{(1-\alpha)(1-\delta)} (\sqrt[4]{1+\sqrt{1-\alpha}} \sqrt[4]{1+\sqrt{1-\delta}} + \sqrt[4]{1-\sqrt{1-\alpha}} \sqrt[4]{1-\sqrt{1-\delta}}) = \sqrt{2}$$

$$vii. \sqrt[4]{\beta\gamma} (\sqrt[4]{1+\sqrt{\beta}} \sqrt[4]{1+\sqrt{\gamma}} - \sqrt[4]{1-\sqrt{\beta}} \sqrt[4]{1-\sqrt{\gamma}})$$

$$+ \sqrt[4]{(1-\beta)(1-\gamma)} (\sqrt[4]{1+\sqrt{1-\beta}} \sqrt[4]{1+\sqrt{1-\gamma}} + \sqrt[4]{1-\sqrt{1-\beta}} \sqrt[4]{1-\sqrt{1-\gamma}}) = \sqrt{2}$$

$$viii. \sqrt[4]{\frac{\alpha\delta}{\beta\gamma}} + \sqrt[4]{\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}} - \sqrt[4]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} = \sqrt{\frac{1+(\frac{1}{2})^2\beta+\alpha c}{1+(\frac{1}{2})^2\alpha+\beta c}} \sqrt{\frac{1+(\frac{1}{2})^2\gamma+\delta c}{1+(\frac{1}{2})^2\delta+\gamma c}}$$

$$ix. \sqrt[4]{\frac{\beta\gamma}{\alpha\delta}} + \sqrt[4]{\frac{(1-\beta)(1-\gamma)}{(1-\alpha)(1-\delta)}} - \sqrt[4]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} = -\sqrt{\frac{1+(\frac{1}{2})^2\alpha+\beta c}{1+(\frac{1}{2})^2\beta+\alpha c}} \sqrt{\frac{1+(\frac{1}{2})^2\delta+\gamma c}{1+(\frac{1}{2})^2\gamma+\delta c}}$$

$$x. \sqrt[4]{\frac{\beta\delta}{\alpha\gamma}} + \sqrt[4]{\frac{(1-\beta)(1-\delta)}{(1-\alpha)(1-\gamma)}} = \sqrt[4]{\frac{\beta\delta(1-\beta)(1-\delta)}{\alpha\gamma(1-\alpha)(1-\gamma)}} - 4 \sqrt[4]{\frac{\delta/\beta\delta(1-\beta)(1-\delta)}{\alpha\gamma(1-\alpha)(1-\gamma)}}$$

$$= \frac{1+(\frac{1}{2})^2\alpha+\beta c}{1+(\frac{1}{2})^2\beta+\alpha c} \cdot \frac{1+(\frac{1}{2})^2\gamma+\delta c}{1+(\frac{1}{2})^2\delta+\gamma c}$$

$$xi. \sqrt[4]{\frac{\alpha\gamma}{\beta\delta}} + \sqrt[4]{\frac{(1-\alpha)(1-\gamma)}{(1-\beta)(1-\delta)}} - \sqrt[4]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta\delta(1-\beta)(1-\delta)}} = 4 \sqrt[4]{\frac{\delta/\alpha\gamma(1-\alpha)(1-\gamma)}{\beta\delta(1-\beta)(1-\delta)}}$$

$$= 9 \cdot \frac{1+(\frac{1}{2})^2\beta+\alpha c}{1+(\frac{1}{2})^2\alpha+\beta c} \cdot \frac{1+(\frac{1}{2})^2\delta+\gamma c}{1+(\frac{1}{2})^2\gamma+\delta c}$$

$$xii. \sqrt[4]{\frac{\gamma\delta}{\alpha\beta}} + \sqrt[4]{\frac{(1-\gamma)(1-\delta)}{(1-\alpha)(1-\beta)}} + \sqrt[4]{\frac{\gamma\delta(1-\gamma)(1-\delta)}{\alpha\beta(1-\alpha)(1-\beta)}}$$

$$- 2 \sqrt[4]{\frac{\gamma\delta(1-\gamma)(1-\delta)}{\alpha\beta(1-\alpha)(1-\beta)}} \left\{ 1 + \sqrt[4]{\frac{\gamma\delta}{\alpha\beta}} + \sqrt[4]{\frac{(1-\gamma)(1-\delta)}{(1-\alpha)(1-\beta)}} \right\} = \frac{1+(\frac{1}{2})^2\alpha+\beta c}{1+(\frac{1}{2})^2\beta+\alpha c} \cdot \frac{1+(\frac{1}{2})^2\delta+\gamma c}{1+(\frac{1}{2})^2\gamma+\delta c}$$

$$\text{iii. } \sqrt[4]{\frac{\alpha\beta}{\gamma\delta}} + \sqrt[3]{\frac{(1-\alpha)(1-\beta)}{(1-\gamma)(1-\delta)}} + \sqrt[5]{\frac{\alpha\beta(1-\delta)(1-\alpha)}{\gamma\delta(1-\gamma)(1-\delta)}} - 2\sqrt[8]{\frac{\alpha\beta(1-\alpha)(1-\beta)}{\gamma\delta(1-\gamma)(1-\delta)}} \times$$

$$\left\{ 1 + \sqrt[4]{\frac{\alpha\beta}{\gamma\delta}} + \sqrt[3]{\frac{(1-\alpha)(1-\beta)}{(1-\gamma)(1-\delta)}} \right\} = 25 \cdot \frac{1 + (\alpha)^2\gamma + \alpha\epsilon}{1 + (\alpha)^2\alpha + 2\alpha\epsilon} \cdot \frac{1 + (\beta)^2\delta + 2\alpha\epsilon}{1 + (\beta)^2\beta + 2\alpha\epsilon}$$

$$\text{iv. } \sqrt{\alpha\beta\gamma\delta} + \sqrt{(1-\alpha)(1-\beta)(1-\gamma)(1-\delta)} + \sqrt{2}\sqrt{\alpha\beta\gamma\delta(1-\alpha)(1-\gamma)(1-\delta)} = 1.$$

$$\text{xv. } P/P = \sqrt[8]{256\alpha\beta\gamma\delta(1-\alpha)(1-\beta)(1-\gamma)(1-\delta)} \text{ and}$$

$$Q = \sqrt[16]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}}, \text{ then } Q + \frac{1}{Q} = \sqrt{2} (P + \frac{1}{P})$$

$$\text{12. i. } \frac{f(-x^{17})}{x^{17}f(x^{17})} = \frac{f(-x^6, -x^{11})}{x^{17}f(-x^3, -x^{14})} - \frac{1}{x^{17}} \frac{f(-x^4, -x^{13})}{f(-x^7, -x^{10})}$$

$$- \frac{1}{x^{17}} \frac{f(-x^9, -x^8)}{f(-x^5, -x^{12})} + \frac{1}{x^{17}} \frac{f(-x^2, -x^{15})}{f(-x, -x^{16})} + \frac{1}{x^{17}} \frac{f(-x^3, -x^{14})}{f(-x^7, -x^{10})}$$

$$- 1 - x^{17} \frac{f(-x^5, -x^{12})}{f(-x^6, -x^{11})} + x^{17} \frac{f(-x^3, -x^{14})}{f(-x^7, -x^{10})} - x^{17} \frac{f(-x^2, -x^{15})}{f(-x^9, -x^8)}$$

$$= u_1 - u_2 - u_3 + u_4 + u_5 - 1 - u_6 + u_7 - u_8 \text{ where}$$

$$u_1 u_5 u_6 u_7 = u_2 u_8 u_3 u_4 = 1, \text{ and}$$

$$u_3 u_6 + u_2 u_8 - u_3 u_4 - u_6 u_7 = -1.$$

$$\text{ii. } f(x, -x^{16}) f(x^2, -x^{11}) f(x^3, -x^{14}) f(x^4, -x^{13}) f(x^5, -x^{12}) \times$$

$$f(x^6, -x^{11}) f(x^7, -x^{10}) f(x^8, -x^9) = f(x) f(x^{17})$$

iii If  $\beta$  be of the 17th degree,

$$m = \sqrt[4]{\frac{\alpha}{\beta}} + \sqrt[4]{\frac{1-\alpha}{1-\beta}} + \sqrt[5]{\frac{\alpha(1-\beta)}{\beta(1-\alpha)}} - 2\sqrt[8]{\frac{\alpha(1-\beta)}{\beta(1-\alpha)}} \left\{ 1 + \sqrt[4]{\frac{\alpha}{\beta}} + \sqrt[4]{\frac{1-\beta}{1-\alpha}} \right\}.$$

$$\text{iv. } \frac{17}{m} = \sqrt[4]{\frac{\alpha}{\beta}} + \sqrt[4]{\frac{1-\alpha}{1-\beta}} + \sqrt[5]{\frac{\alpha(1-\beta)}{\beta(1-\alpha)}} - 2\sqrt[8]{\frac{\alpha(1-\beta)}{\beta(1-\alpha)}} \left\{ 1 + \sqrt[4]{\frac{\alpha}{\beta}} + \sqrt[4]{\frac{1-\beta}{1-\alpha}} \right\}$$

N.B. Thus we see that  $\phi(x^{1/n}), \psi(x^{1/n})$  or  $f(x^{1/n})$  n being any prime number can be expressed as the sum of  $\frac{n-1}{2}$ , nth roots of several functions and  $\phi(x^{-1}), \psi(x^{-1})$  and  $f(x^{-1})$ . In finding the values of these functions, quadratics only appear in case of the 5th, 17th, 257th &c degrees and cubics in case of the 7th, 13th, 19th, 37th, 73rd, 97th, 109th, 163rd, 193rd &c degrees not as cube roots but as  $\sin(\frac{1}{3} \sin^{-1} \theta)$  and quintics in case of the 11th, 16th, 101st &c degrees.  $f(x^{1/n})$  can also be similarly expressed.

13. If  $\alpha, \beta, \gamma$  &  $\delta$  be of the 1st, 3rd, 7th and 21<sup>st</sup> degree,

i.  $\sqrt[4]{\frac{\alpha\gamma}{\beta\delta}} + \sqrt[4]{\frac{(1-\beta)(1-\gamma)}{(1-\alpha)(1-\delta)}} - \sqrt[4]{\frac{\beta\gamma(1-\alpha)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} + \sqrt[4]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}}$   
 $= \frac{1 + (\zeta^2)^4 \alpha + \beta\epsilon}{1 + (\zeta^2)^4 \beta + \alpha\epsilon} \cdot \frac{1 + (\zeta^2)^4 \delta + \gamma\epsilon}{1 + (\zeta^2)^4 \gamma + \delta\epsilon}$

ii.  $\sqrt[4]{\frac{\alpha\delta}{\beta\gamma}} + \sqrt[4]{\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}} - \sqrt[4]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} + \sqrt[4]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}}$   
 $= \frac{1 + (\zeta^2)^4 \beta + \alpha\epsilon}{1 + (\zeta^2)^4 \alpha + \beta\epsilon} \cdot \frac{1 + (\zeta^2)^4 \gamma + \delta\epsilon}{1 + (\zeta^2)^4 \delta + \gamma\epsilon}$

iii.  $\sqrt[8]{\frac{\alpha\delta}{\beta\gamma}} + \sqrt[8]{\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}} - \sqrt[8]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} - 2\sqrt[8]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}}$   
 $= \sqrt{\frac{1 + (\zeta^2)^4 \alpha + \beta\epsilon}{1 + (\zeta^2)^4 \gamma + \delta\epsilon}} \sqrt{\frac{1 + (\zeta^2)^4 \beta + \alpha\epsilon}{1 + (\zeta^2)^4 \delta + \gamma\epsilon}}$

iv.  $\sqrt[8]{\frac{\alpha\beta}{\gamma\delta}} + \sqrt[8]{\frac{(1-\alpha)(1-\beta)}{(1-\gamma)(1-\delta)}} - \sqrt[8]{\frac{\alpha\beta(1-\alpha)(1-\beta)}{\gamma\delta(1-\gamma)(1-\delta)}} - 2\sqrt[8]{\frac{\alpha\beta(1-\alpha)(1-\beta)}{\gamma\delta(1-\gamma)(1-\delta)}}$   
 $= 7\sqrt{\frac{1 + (\zeta^2)^4 \gamma + \delta\epsilon}{1 + (\zeta^2)^4 \alpha + \beta\epsilon}} \sqrt{\frac{1 + (\zeta^2)^4 \delta + \gamma\epsilon}{1 + (\zeta^2)^4 \beta + \alpha\epsilon}}$

v.  $\sqrt[4]{\frac{\beta\delta}{\alpha\gamma}} + \sqrt[4]{\frac{(1-\beta)(1-\delta)}{(1-\alpha)(1-\gamma)}} + \sqrt[4]{\frac{\beta\delta(1-\beta)(1-\delta)}{\alpha\gamma(1-\alpha)(1-\gamma)}} - 2\sqrt[4]{\frac{\beta\delta(1-\beta)(1-\delta)}{\alpha\gamma(1-\alpha)(1-\gamma)}}$   
 $\left. \sqrt[8]{\frac{\alpha\delta}{\beta\gamma}} + \sqrt[8]{\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}} \right\} = \frac{1 + (\zeta^2)^4 \alpha + \beta\epsilon}{1 + (\zeta^2)^4 \beta + \alpha\epsilon} \cdot \frac{1 + (\zeta^2)^4 \gamma + \delta\epsilon}{1 + (\zeta^2)^4 \delta + \gamma\epsilon}$

vi.  $\sqrt[4]{\frac{\alpha\gamma}{\beta\delta}} + \sqrt[4]{\frac{(1-\alpha)(1-\gamma)}{(1-\beta)(1-\delta)}} + \sqrt[4]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta\delta(1-\beta)(1-\delta)}} - 2\sqrt[4]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta\delta(1-\beta)(1-\delta)}}$   
 $\left. \sqrt[8]{\frac{\alpha\delta}{\beta\gamma}} + \sqrt[8]{\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}} \right\} = 9 \cdot \frac{1 + (\zeta^2)^4 \beta + \alpha\epsilon}{1 + (\zeta^2)^4 \alpha + \beta\epsilon} \cdot \frac{1 + (\zeta^2)^4 \delta + \gamma\epsilon}{1 + (\zeta^2)^4 \gamma + \delta\epsilon}$

14. If  $\alpha, \beta, \gamma, \delta$  be of the 1<sup>st</sup>, 3<sup>rd</sup>, 11<sup>th</sup> and 33<sup>rd</sup> degree.

i.  $\sqrt[8]{\frac{\beta\delta}{\alpha\gamma}} + \sqrt[8]{\frac{(1-\beta)(1-\delta)}{(1-\alpha)(1-\gamma)}} - \sqrt[8]{\frac{\beta\delta(1-\beta)(1-\delta)}{\alpha\gamma(1-\alpha)(1-\gamma)}} - 2\sqrt[8]{\frac{\beta\delta(1-\beta)(1-\delta)}{\alpha\gamma(1-\alpha)(1-\gamma)}}$   
 $= \sqrt{\frac{1 + (\zeta^2)^4 \alpha + \beta\epsilon}{1 + (\zeta^2)^4 \gamma + \delta\epsilon}} \sqrt{\frac{1 + (\zeta^2)^4 \beta + \alpha\epsilon}{1 + (\zeta^2)^4 \delta + \gamma\epsilon}}$

ii.  $\sqrt[8]{\frac{\alpha\gamma}{\beta\delta}} + \sqrt[8]{\frac{(1-\alpha)(1-\gamma)}{(1-\beta)(1-\delta)}} - \sqrt[8]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta\delta(1-\beta)(1-\delta)}} - 2\sqrt[8]{\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta\delta(1-\beta)(1-\delta)}}$   
 $= 3\sqrt{\frac{1 + (\zeta^2)^4 \beta + \alpha\epsilon}{1 + (\zeta^2)^4 \alpha + \beta\epsilon}} \sqrt{\frac{1 + (\zeta^2)^4 \delta + \gamma\epsilon}{1 + (\zeta^2)^4 \gamma + \delta\epsilon}}$

15. If  $\beta$  be of the 23<sup>rd</sup> degree,

i.  $\sqrt[23]{\frac{\alpha\gamma}{\beta}} + \sqrt[23]{(1-\alpha)(1-\beta)} + \sqrt[23]{\frac{\alpha\beta(1-\alpha)(1-\beta)}{\gamma}} = 1$

ii.  $1 + \sqrt[23]{\alpha\beta} + \sqrt[23]{(1-\alpha)(1-\beta)} + 2\sqrt[23]{\frac{\alpha\beta(1-\alpha)(1-\beta)}{\gamma}} =$

$$\sqrt{1 + \sqrt{\alpha\beta} + \sqrt{1 - \alpha(1-\beta)}}$$

iii.  $\frac{1}{m} = 2 \left( \sqrt{\alpha\beta} - \sqrt{1 - \alpha(1-\beta)} \right) \left\{ 1 - 2 \sqrt[4]{\alpha\beta(1-\alpha)(1-\beta)} + 7 \sqrt[4]{\alpha\beta(1-\alpha)(1-\beta)} \right\}$

16. If  $\beta$  be of the 19th degree,

i.  $\sqrt{\frac{1-\beta}{1-\alpha}} - \sqrt{\frac{\beta}{\alpha}} + \sqrt{\frac{\beta^2(1-\beta)^2}{\alpha(1-\alpha)}} - 2 \sqrt{\frac{\beta^3(1-\beta)^3}{\alpha(1-\alpha)}} \times$

$$\sqrt{\frac{1-\beta}{1-\alpha}} - 1 - \sqrt{\frac{\beta}{\alpha}} = m \sqrt{\frac{1 + \sqrt{\alpha\beta} + \sqrt{1 - \alpha(1-\beta)}}{2}}$$

ii.  $\sqrt{\frac{\alpha^2}{\beta}} - \sqrt{\frac{1-\alpha}{1-\beta}} + \sqrt{\frac{\alpha^2(1-\alpha)^2}{\beta(1-\beta)}} - 2 \sqrt{\frac{16\alpha^3(1-\alpha)^3}{7\beta(1-\beta)}} \times$

$$\sqrt{\frac{\alpha^2}{\beta}} - 1 - \sqrt{\frac{1-\alpha}{1-\beta}} = \frac{12}{m} \sqrt{\frac{1 + \sqrt{\alpha\beta} + \sqrt{1 - \alpha(1-\beta)}}{2}}$$

17. i.  $\phi(\alpha) \phi(\alpha^{35}) = \phi(\alpha) \phi(\alpha^{35}) + 4x f(\alpha^{10}) f(\alpha^{14}) + 4x^9 \psi(\alpha^7) \psi(\alpha^{70})$

ii.  $\phi(\alpha^7) \phi(\alpha^7) = \phi(\alpha^7) \phi(\alpha^7) + 4x^2 \psi(\alpha^{10}) \psi(\alpha^{14}) - 4x^3 f(\alpha^5) f(\alpha^{70})$

iii.  $\phi(\alpha^{10}) \phi(\alpha^{54}) + 2x f(\alpha^7) f(\alpha^{15}) + 2x^4 \psi(\alpha^5) \psi(\alpha^{27}) = \phi(\alpha) \phi(\alpha^{135})$

iv.  $\phi(\alpha^2) \phi(\alpha^{270}) + 2x^{17} \psi(\alpha) \psi(\alpha^{135}) + 2x^2 f(\alpha^3) f(\alpha^{45}) = \phi(\alpha^5) \phi(\alpha^{27})$

18. If  $\alpha, \beta, \gamma$  &  $\delta$  be of the 1st, 5th, 7th & 35th degree,

i.  $\frac{\sqrt{\alpha\delta} + \sqrt{1-\alpha(1-\delta)} + 2 \sqrt[4]{\alpha\delta(1-\alpha)(1-\delta)}}{\sqrt{\beta\gamma} + \sqrt{1-\beta(1-\gamma)} + 2 \sqrt[4]{\beta\gamma(1-\beta)(1-\gamma)}} = \frac{1 + \left\{ 1 + 2 \sqrt[4]{\alpha\beta\gamma\delta(1-\alpha)(1-\beta)(1-\gamma)(1-\delta)} \right\}}{2}$

ii.  $\left\{ \sqrt{\alpha\delta} + \sqrt{1-\alpha(1-\delta)} + 2 \sqrt[4]{\alpha\delta(1-\alpha)(1-\delta)} \right\} \times \left\{ \sqrt{\beta\gamma} + \sqrt{1-\beta(1-\gamma)} + 2 \sqrt[4]{\beta\gamma(1-\beta)(1-\gamma)} \right\} = 1 - 4 \sqrt[4]{\alpha\beta\gamma\delta(1-\alpha)(1-\beta)(1-\gamma)(1-\delta)} \left\{ \sqrt{\alpha\beta\gamma\delta} + \sqrt{1-\alpha(1-\beta)(1-\gamma)(1-\delta)} \right\}$

$$\text{iii. } \frac{\sqrt{\alpha\delta} + \sqrt{(1-\alpha)(1-\delta)}}{\sqrt{\beta\gamma}} + 2\sqrt[3]{2} \sqrt[4]{\alpha\delta(1-\beta)(1-\gamma)} \cdot \sqrt{\frac{1+(\frac{1}{2})^2\beta+2c}{1+(\frac{1}{2})^2\alpha+2c} \cdot \frac{1+(\frac{1}{2})^2\gamma+2c}{1+(\frac{1}{2})^2\delta+2c}} = 1$$

$$\text{iv. } \frac{\sqrt{\beta\gamma} + \sqrt{(1-\beta)(1-\gamma)}}{\sqrt{\alpha\delta}} - 2\sqrt[3]{2} \sqrt[4]{\alpha\delta(1-\alpha)(1-\delta)} \sqrt{\frac{1+(\frac{1}{2})^2\alpha+2c}{1+(\frac{1}{2})^2\beta+2c} \cdot \frac{1+(\frac{1}{2})^2\delta+2c}{1+(\frac{1}{2})^2\gamma+2c}} = 1$$

$$\text{v. } \sqrt{\frac{1+(\frac{1}{2})^2\beta+2c}{1+(\frac{1}{2})^2\alpha+2c} \cdot \frac{1+(\frac{1}{2})^2\gamma+2c}{1+(\frac{1}{2})^2\delta+2c}} = \frac{\sqrt[4]{\beta\gamma(1-\beta)(1-\gamma)} - \sqrt[4]{\alpha\delta(1-\alpha)(1-\delta)}}{\sqrt[4]{\beta\alpha\gamma(1-\beta)(1-\gamma)} + \sqrt[4]{\alpha\delta(1-\alpha)(1-\delta)}} \\ = \frac{\sqrt[4]{\beta\delta(1-\alpha)(1-\delta)} + \sqrt[4]{\beta\alpha\delta(1-\alpha)(1-\delta)}}{\sqrt[4]{\beta\alpha\gamma(1-\beta)(1-\gamma)} - \sqrt[4]{\beta\alpha\delta(1-\alpha)(1-\delta)}}$$

$$\text{vi. } \frac{\sqrt[3]{\alpha\delta}}{\sqrt[3]{\beta\gamma}} + \frac{\sqrt[3]{(1-\alpha)(1-\delta)}}{\sqrt[3]{(1-\beta)(1-\gamma)}} - \sqrt[3]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} + 2\sqrt[4]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} \\ = \sqrt{\frac{1+(\frac{1}{2})^2\beta+2c}{1+(\frac{1}{2})^2\alpha+2c} \cdot \frac{1+(\frac{1}{2})^2\gamma+2c}{1+(\frac{1}{2})^2\delta+2c}}$$

$$\text{vii. } \frac{\sqrt[3]{\beta\gamma}}{\sqrt[3]{\alpha\delta}} + \frac{\sqrt[3]{(1-\beta)(1-\gamma)}}{\sqrt[3]{(1-\alpha)(1-\delta)}} - \sqrt[3]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} + 2\sqrt[4]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} \\ = \sqrt{\frac{1+(\frac{1}{2})^2\alpha+2c}{1+(\frac{1}{2})^2\beta+2c} \cdot \frac{1+(\frac{1}{2})^2\delta+2c}{1+(\frac{1}{2})^2\gamma+2c}}$$

$$\text{19. i. } \phi(\alpha)\phi(\alpha^2) - \phi(\alpha^7)\phi(\alpha^7) = z^x f(\alpha^3)f(\alpha^2)$$

$$\text{ii. } \psi(\alpha^7)\psi(\alpha^7) - x^6\psi(\alpha)\psi(\alpha^2) = f(-x^6)f(x^2)$$

iii. If  $\alpha, \beta, \gamma, \delta$  be of the 1st, 3rd, 13th and 39th degree or 1st, 5th, 11th and 55th degree or 1st, 7th, 9th and 63rd degree

$$\frac{1 + \sqrt{(1-\alpha)(1-\delta)} + \sqrt{\alpha\delta}}{1 + \sqrt{(1-\beta)(1-\gamma)} + \sqrt{\beta\gamma}} = \frac{\sqrt[3]{(1-\alpha)(1-\delta)} - \sqrt[3]{\alpha\delta}}{\sqrt[3]{(1-\beta)(1-\gamma)} - \sqrt[3]{\beta\gamma}} =$$

$$\sqrt{\frac{1+(\frac{1}{2})^2\beta+2c}{1+(\frac{1}{2})^2\alpha+2c} \cdot \frac{1+(\frac{1}{2})^2\gamma+2c}{1+(\frac{1}{2})^2\delta+2c}} = \frac{\sqrt[3]{\alpha\delta} \pm \sqrt[3]{\alpha\delta(1-\alpha)(1-\delta)}}{\sqrt[3]{\beta\gamma} - \sqrt[3]{\beta\gamma(1-\beta)(1-\gamma)}}$$

(+ in the first two cases and - in the last case.)

iv. If  $\alpha, \beta, \gamma, \delta$  be of the 1st, 3rd, 13th and 39th degree or 1st, 5th, 7th and 55th degree

$$\frac{\sqrt[3]{\alpha\delta}}{\sqrt[3]{\beta\gamma}} + \frac{\sqrt[3]{(1-\alpha)(1-\delta)}}{\sqrt[3]{(1-\beta)(1-\gamma)}} - \sqrt[3]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} + 2\sqrt[4]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} \\ = \sqrt{\frac{1+(\frac{1}{2})^2\beta+2c}{1+(\frac{1}{2})^2\alpha+2c}} \sqrt{\frac{1+(\frac{1}{2})^2\gamma+2c}{1+(\frac{1}{2})^2\delta+2c}} \text{ and}$$

$$\sqrt{\frac{\alpha\gamma}{\alpha\delta}} + \sqrt{\frac{(1-\alpha)(1-\gamma)}{(1-\alpha)(1-\delta)}} - \sqrt{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} + 2\sqrt{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} \quad (24)$$

$$= \pm \sqrt{\frac{1+(6)^2\alpha+8\epsilon}{1+(6)^2\beta+8\epsilon}} \cdot \sqrt{\frac{1+(6)^2\delta+8\epsilon}{1+(6)^2\gamma+8\epsilon}} \quad (+ \text{ in the I case } \& - \text{ in the II})$$

20. If  $\alpha, \beta, \gamma, \delta$  be of the 1st-3rd 21st and 63rd degree or 1st-5th 19th 95th or 1st-11th 15th 143rd or 1st-7th 17th 119th or 1st-9th 15th 135th degree then

$$\sqrt{\frac{1+\sqrt{\alpha\delta} + \sqrt{(1-\alpha)(1-\delta)}}{2}} = \sqrt[3]{\alpha\delta} + \sqrt[3]{(1-\alpha)(1-\delta)} \pm \sqrt[3]{\alpha\delta(1-\alpha)(1-\delta)} +$$

$$2\sqrt[3]{\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}} \sqrt{\frac{1+(6)^2\alpha+8\epsilon}{1+(6)^2\beta+8\epsilon}} \cdot \sqrt{\frac{1+(6)^2\delta+8\epsilon}{1+(6)^2\gamma+8\epsilon}}$$

(+ in the first 3 cases and - in the last 2 cases)

ii. If  $\alpha, \beta, \gamma, \delta$  be of the 1st-5th 19th 95th degree or 1st-7th 17th 119th or 1st-11th 13th 143rd degree, then

$$\sqrt{\frac{1+\sqrt{\beta\gamma} + \sqrt{(1-\beta)(1-\gamma)}}{2}} = \sqrt[3]{\beta\gamma} + \sqrt[3]{(1-\beta)(1-\gamma)} - \sqrt[3]{\beta\gamma(1-\beta)(1-\gamma)}$$

$$\pm 2\sqrt[3]{\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}} \cdot \sqrt{\frac{1+(6)^2\alpha+8\epsilon}{1+(6)^2\beta+8\epsilon}} \cdot \sqrt{\frac{1+(6)^2\delta+8\epsilon}{1+(6)^2\gamma+8\epsilon}}$$

(- in the first 2 cases and + in the last one)

21. i. If  $\alpha, \beta$  be of the 1st and 7th or 3rd and 5th or 1st-3-5-7th

$$\sqrt{\frac{1+\sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}} = \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} \pm \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)}$$

(- in the 1st 2 cases and + in the last)

ii. If  $\alpha, \beta, \gamma, \delta$  be of the 1st-3rd 13th 39th or 1st-5th 11th 55th or 1st-7th 9th 63rd degree, then

$$\sqrt{\frac{1+\sqrt{\alpha\delta} + \sqrt{(1-\alpha)(1-\delta)}}{2}} = \sqrt[3]{(1-\alpha)(1-\delta)} +$$

$$\left(\sqrt[3]{\beta\gamma} + \sqrt[3]{\beta\gamma(1-\beta)(1-\gamma)}\right) \sqrt{\frac{1+(6)^2\alpha+8\epsilon}{1+(6)^2\beta+8\epsilon}} \cdot \sqrt{\frac{1+(6)^2\delta+8\epsilon}{1+(6)^2\gamma+8\epsilon}}$$

and also  $\sqrt{\frac{1+\sqrt{\beta\gamma} + \sqrt{(1-\beta)(1-\gamma)}}{2}} = \sqrt[3]{(1-\beta)(1-\gamma)} +$

$$\left(\sqrt[3]{\alpha\delta} \pm \sqrt[3]{\alpha\delta(1-\alpha)(1-\delta)}\right) \sqrt{\frac{1+(6)^2\alpha+8\epsilon}{1+(6)^2\beta+8\epsilon}} \cdot \sqrt{\frac{1+(6)^2\delta+8\epsilon}{1+(6)^2\gamma+8\epsilon}}$$

(- in the 1st 2 cases and + in the last)

22. If  $\beta$  be of the 31st degree

$$i. \sqrt[3]{\alpha\beta} \left\{ \sqrt[3]{(1+\sqrt{\alpha}\alpha)(1+\sqrt{\beta})} \sqrt[3]{1+\sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\sqrt{\alpha}\alpha)(1-\sqrt{\beta})}} + \sqrt[3]{(1-\sqrt{\alpha}\alpha)(1-\sqrt{\beta})} \sqrt[3]{1+\sqrt[3]{\alpha\beta} + \sqrt[3]{(1+\sqrt{\alpha}\alpha)(1+\sqrt{\beta})}} \right\} + \sqrt[3]{(1-\alpha)(1-\beta)} \left\{ \dots \dots \dots \right\} = \sqrt[3]{8}$$

$$ii. 1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} - 2 \left( \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} + \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)} \right) = 2 \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)} \sqrt[3]{1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)}} \quad \checkmark$$

$$iii. 1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} - \sqrt[3]{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}} = \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} + \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)}$$

13. i. If  $\beta$  be of the 47th degree,

$$2 \sqrt[3]{\frac{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}} = \left( 1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} \right) + \frac{3}{4} \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)} \left\{ 1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} \right\}$$

ii. If  $\beta$  be of the 71st degree

$$1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} - \sqrt[3]{\frac{1 + \sqrt{\alpha\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}} = \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} + \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)}$$

$$+ \frac{3}{4} \sqrt[3]{\alpha\beta(1-\alpha)(1-\beta)} \left\{ 1 + \sqrt[3]{\alpha\beta} + \sqrt[3]{(1-\alpha)(1-\beta)} \right\}$$

24. If  $\alpha, \beta, \gamma, \delta$  be of the 1st 3rd 29th 87th or 1st 5th 27th 135th or 1st 11th 21st 231st or 1st 13th 19th 247th or 1st 7th 35th 175th or 1st 9th 27th 207th or 1st 15th 17th 255th degree, then

$$i. \sqrt[3]{\frac{1 + \sqrt{\alpha\gamma} + \sqrt{(1-\alpha)(1-\gamma)}}{2}} + \sqrt[3]{\beta\gamma} + \sqrt[3]{(1-\alpha)(1-\gamma)} + \sqrt[3]{\alpha\beta\gamma(1-\alpha)(1-\gamma)} = \left( 1 + \sqrt[3]{\alpha\delta} + \sqrt[3]{(1-\alpha)(1-\delta)} \right) \sqrt{\frac{1 + (\beta^2)\delta + \beta\epsilon}{1 + (\beta^2)\gamma + \beta\epsilon} \cdot \frac{1 + (\beta^2)\delta + \beta\epsilon}{1 + (\beta^2)\gamma + \beta\epsilon}}$$

$$ii. \sqrt[3]{\frac{1 + \sqrt{\alpha\delta} + \sqrt{(1-\alpha)(1-\delta)}}{2}} + \sqrt[3]{\alpha\delta} + \sqrt[3]{(1-\alpha)(1-\delta)} \pm \sqrt[3]{\alpha\delta(1-\alpha)(1-\delta)} = \left( 1 + \sqrt[3]{\beta\gamma} + \sqrt[3]{(1-\alpha)(1-\gamma)} \right) \sqrt{\frac{1 + (\beta^2)\delta + \beta\epsilon}{1 + (\beta^2)\alpha + \beta\epsilon} \cdot \frac{1 + (\beta^2)\gamma + \beta\epsilon}{1 + (\beta^2)\delta + \beta\epsilon}}$$

(- in the 1st 4 cases and + in the last 3).

i.  $1 - \frac{3}{4} = 24 \left( \frac{1}{e^{12}-1} + \frac{2}{e^{24}-1} + \frac{3}{e^{36}-1} + \frac{4}{e^{48}-1} + \dots \right)$  is a complete series which when divided by  $2^2$  can be expressed as radicals precisely in the same manner as the series  $1 + 240 \left( \frac{1^3}{e^{20}-1} + \frac{2^3}{e^{40}-1} + \frac{3^3}{e^{60}-1} + \frac{4^3}{e^{80}-1} + \dots \right)$  and the series  $1 - 504 \left( \frac{1^5}{e^{15}-1} + \frac{2^5}{e^{30}-1} + \frac{3^5}{e^{45}-1} + \frac{4^5}{e^{60}-1} + \dots \right)$  when divided by  $2^4$  and  $2^6$  respectively.

$$\text{ii. } 1 - 24 \left( \frac{1}{e^{12}+1} + \frac{2}{e^{24}+1} + \frac{5}{e^{36}+1} + \dots \right) = 2^2 (1 - 2x)$$

$$\text{iii. } 1 - 240 \left( \frac{1^3}{e^{20}+1} - \frac{2^3}{e^{40}-1} + \frac{3^3}{e^{60}+1} - \dots \right) = 2^4 (1 - 16x \cdot 1 - x)$$

$$\text{iv. } 1 + 504 \left( \frac{1^5}{e^{15}+1} - \frac{2^5}{e^{30}-1} + \frac{3^5}{e^{45}+1} - \dots \right) = 2^6 (1 - 3x)(1 + 32x \cdot 1 - x)$$

$$\text{2.112 } x \frac{d\phi(x)}{dx} / \phi(x) = \left\{ 1 - 24 \left( \frac{x^2}{1-x^4} + \frac{2x^4}{1-x^8} + \dots \right) \right\} \\ - \left\{ 1 - 24 \left( \frac{x}{1+x} + \frac{3x^2}{1+x^2} + \dots \right) \right\}$$

$$\text{ii. } 24x \frac{d x^{1/2} \psi(x)}{dx} / x^{1/2} \psi(x) = \left\{ 1 - 24 \left( \frac{x^6}{1-x^8} + \frac{2x^8}{1-x^{16}} + \dots \right) \right\} \\ - \left\{ 1 - 24 \left( \frac{x}{1+x} + \frac{3x^2}{1+x^2} + \dots \right) \right\}$$

$$\text{iii. } 24x \frac{d x^{1/4} f(x)}{dx} / x^{1/4} f(x) = 1 - 24 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^4} + \frac{3x^3}{1-x^2} + \dots \right)$$

$$\text{iv. } 24x \frac{d x^{1/2} / \chi(x)}{dx} / x^{1/2} \chi(x) = 1 - 24 \left( \frac{x}{1+x} + \frac{2x^2}{1+x^2} + \frac{5x^3}{1+x^4} + \dots \right)$$

v. By differentiating the equation for  $\alpha$  once or the equation for  $\beta$  twice we can calculate the value of the first series.

$$\text{2.1 } 1 + 12 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^4} + \frac{3x^3}{1-x^2} + \dots \right) - 12 \left( \frac{2x^4}{1-x^8} + \frac{6x^6}{1-x^{16}} + \dots \right)$$

$$= \left\{ 1 + 6 \left( \frac{x}{1-x} - \frac{x^4}{1-x^8} + \frac{x^3}{1-x^2} - \dots \right) \right\}^2$$

$$= \left\{ 1 + 24 \left( \frac{1^3}{1-x} + \frac{2^3 x^2}{1-x^4} + \frac{3^3 x^3}{1-x^2} + \dots \right) + 8 \left( \frac{2^3 x^4}{1-x^8} + \frac{6^3 x^6}{1-x^{16}} + \dots \right) \right\}^2$$

$$= \left\{ \frac{\psi^4(x) + 3x\psi^4(x^2)}{\psi(x)\psi(x^2)} \right\}^2 = \left\{ \frac{f^{1/2}(x) + 27f^{1/2}(x^2)}{f(x)f(x^2)} \right\}^2$$

$$\text{ii. } 1 + 12 \left( \frac{x^6}{1-x^8} + \frac{2x^8}{1-x^{16}} + \dots \right) - 12 \left( \frac{2x^4}{1-x^8} + \frac{6x^6}{1-x^{16}} + \dots \right)$$

$$= \left\{ \frac{\phi^4(x) + 3\phi^4(x^2)}{\phi(x)\phi(x^2)} \right\}^2 = \phi^4(x)\phi^4(x^2) - 4x\psi^4(x)\psi^4(x^2)$$



$$150$$

$$\text{iii. } 1 + 12 \left( \frac{1}{e^{2y}} + \frac{2}{e^{4y}} + 2c \right) - 12 \left( \frac{3}{e^{6y}} + \frac{6}{e^{8y}} + 2c \right)$$

$$= \phi^2(e^{-y}) \phi^2(e^{-2y}) \cdot \frac{1 + \sqrt{3\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}$$

$$4. \text{i. } 1 + 6 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^2} + \frac{3x^3}{1-x^3} + 2c \right) - 6 \left( \frac{5x^4}{1-x^4} + \frac{10x^{10}}{1-x^{10}} + 2c \right)$$

$$= \sqrt{f^{10}(x) + 22x f^6(x) f^4(x^2) + 125x^4 f^8(x^2)} / f(x) f(x^2)$$

$$= \frac{\psi^4(x) + 2x \psi^4(x) \psi^4(x^2) + 5x^4 \psi^4(x^2)}{\psi(x) \psi(x^2)} \sqrt{\psi^4(x) - 2x \psi^4(x) \psi^4(x^2) + 5x^4 \psi^4(x^2)}$$

$$\text{ii. } 1 + 6 \left( \frac{x^2}{1-x^2} + \frac{2x^4}{1-x^4} + \frac{3x^6}{1-x^6} + 2c \right) - 6 \left( \frac{5x^{10}}{1-x^{10}} + \frac{10x^{20}}{1-x^{20}} + 2c \right)$$

$$= \left\{ \phi^2(x) \phi^2(x^2) - 2x f^4(x^2) f^4(x^{10}) \right\} \sqrt{1 - 2x / \chi^4(x) \chi^4(x^2)}$$

$$\text{iii. } 1 + 6 \left( \frac{1}{e^{2y}} + \frac{2}{e^{4y}} + 2c \right) - 6 \left( \frac{6}{e^{10y}} + \frac{10}{e^{20y}} + 2c \right)$$

$$= \phi^2(e^{-y}) \phi^2(e^{-5y}) \cdot \frac{3 + \sqrt{3\beta} + \sqrt{(1-\alpha)(1-\beta)}}{4} \sqrt{\frac{1 + \sqrt{3\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}}$$

$$= \phi^2(e^{-y}) \phi^2(e^{-5y}) \sqrt{\frac{1 + \sqrt{3\beta} + (1-\alpha)(1-\beta)}{2}} - \frac{3}{4} \sqrt{10\beta(1-\alpha)(1-\beta)}$$

$$5. \text{i. } 1 + 4 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^2} + \frac{3x^3}{1-x^3} + 2c \right) - 4 \left( \frac{7x^7}{1-x^7} + \frac{14x^{14}}{1-x^{14}} + 2c \right)$$

$$= \left\{ 1 + 2 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^2} - \frac{x^2}{1-x^2} + \frac{x^4}{1-x^4} - \frac{x^5}{1-x^5} - \frac{x^6}{1-x^6} + \frac{x^8}{1-x^8} + 2c \right) \right\}^2$$

$$= \left\{ \frac{f^8(x^7) + 13x f^6(x) f^4(x^7) + 49x^4 f^8(x^7)}{f(x) f(x^7)} \right\}^2$$

$$\text{ii. } 1 + 4 \left( \frac{x^2}{1-x^2} + \frac{2x^4}{1-x^4} + \frac{3x^6}{1-x^6} + 2c \right) - 4 \left( \frac{7x^{14}}{1-x^{14}} + \frac{14x^{28}}{1-x^{28}} + 2c \right)$$

$$= \left\{ \phi(x) \phi(x^7) - 2x \psi(x^7) \psi(x^{14}) \right\}^2$$

$$\text{iii. } 1 + 4 \left( \frac{1}{e^{2y}} + \frac{2}{e^{4y}} + 2c \right) - 4 \left( \frac{7}{e^{14y}} + \frac{14}{e^{28y}} + 2c \right)$$

$$= \phi^2(e^{-y}) \phi^2(e^{-7y}) \cdot \frac{1 + \sqrt{3\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2}$$

$$6. \text{i. } 1 - 12 \left( \frac{1}{e^{2y}} + \frac{2}{e^{4y}} + \frac{3}{e^{6y}} - 2c \right) + 12 \left( \frac{3}{e^{10y}} + \frac{6}{e^{20y}} + 2c \right)$$

$$= \phi^2(e^{-y}) \phi^2(e^{-2y}) \left\{ \sqrt{3\beta} - \sqrt{(1-\alpha)(1-\beta)} \right\}^2$$

$$\text{ii. } 1 - 6 \left( \frac{1}{e^{4y}} + \frac{2}{e^{8y}} + \frac{3}{e^{12y}} - 2c \right) + 6 \left( \frac{6}{e^{20y}} + \frac{10}{e^{40y}} + 2c \right)$$

- =  $\phi^4(e^{-7y}) \phi^4(e^{-11y}) (\sqrt{ab} + \sqrt{a-2)(b-2}) \sqrt{1 + \sqrt{ab} + \sqrt{a-2)(b-2)}$
- iii.  $1 - 4 \left( \frac{1}{e^{4y} - 1} - \frac{2}{e^{12y} - 1} + \frac{1}{e^{20y} - 1} - 2c \right) + 4 \left( \frac{7}{e^{28y} - 1} - \frac{14}{e^{40y} - 1} + 2c \right)$   
 =  $\phi^2(e^{-7y}) \phi^2(e^{-11y}) \left\{ \sqrt{ab} + \sqrt{a-2)(b-2) \right\}^2$
- 7. i.  $1 + 3 \left( \frac{x}{1-x} + \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} + 2c \right) - 3 \left( \frac{9x^2}{1-x^6} + \frac{19x^{14}}{1-x^{14}} + 2c \right)$   
 =  $\frac{f^6(x^3)}{f^2(x)f^2(x^9)} \left\{ f^6(x) + 9x f^3(x) f^3(x^9) + 27x^2 f^6(x^9) \right\}^{\frac{1}{3}}$   
 =  $\left\{ \frac{\psi^6(\alpha^3) + 3 \psi^3(\alpha) \psi^3(\alpha^9)}{\psi(\alpha^3) \psi(\alpha^9)} \right\}^{\frac{1}{3}} \cdot \frac{\psi^2(\alpha^3)}{\psi(\alpha) \psi(\alpha^9)}$
- ii.  $1 + 3 \left( \frac{x^2}{1-x^2} + \frac{x^4}{1-x^4} + 2c \right) - 3 \left( \frac{9x^{18}}{1-x^{18}} + \frac{19x^{26}}{1-x^{26}} + 2c \right)$   
 =  $\left\{ \frac{\phi^6(\alpha^3) + 3 \phi^3(\alpha) \phi^3(\alpha^9)}{4} \right\}^{\frac{1}{3}} \cdot \frac{\phi^2(\alpha^3)}{f^2(\alpha) \phi^2(\alpha^9)}$
- iii.  $1 + \left( \frac{x}{1-x} + \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} + 2c \right) - \left( \frac{25x^{14}}{1-x^{14}} + \frac{59x^{50}}{1-x^{50}} + 2c \right)$   
 =  $\frac{f^5(x^5)}{f(x)f(x^5)} \sqrt{f^5(x) + 2x f(x) f(x^5) + 5x^2 f^2(x^5)}$
- 8. i.  $5 + 12 \left( \frac{x^2}{1-x^2} + \frac{x^4}{1-x^4} + \frac{x^6}{1-x^6} + 2c \right) - 12 \left( \frac{11x^{11}}{1-x^{11}} + \frac{22x^{55}}{1-x^{55}} + 2c \right)$   
 =  $5 \psi^2(\alpha) \psi^2(\alpha^{11}) - 20x f^2(\alpha) f^2(\alpha^{11}) + 32x^2 f^2(x) f^2(x^{11}) - 20x^2 \psi^2(x) \psi^2(x^{11})$
- ii.  $5 + 12 \left( \frac{1}{e^{2y} - 1} + \frac{2}{e^{4y} - 1} + \frac{3}{e^{6y} - 1} + 2c \right) - 12 \left( \frac{11}{e^{11y} - 1} + \frac{22}{e^{55y} - 1} + 2c \right)$   
 =  $\phi^2(e^{-y}) \phi^2(e^{-11y}) \left\{ 2(1 + \sqrt{ab} + \sqrt{a-2)(b-2}) + \sqrt{ab} + \sqrt{a-2)(b-2) - \sqrt{ab(a-2)(b-2)} \right\}$
- iii.  $3 + 4 \left( \frac{1}{e^{12y} - 1} + \frac{2}{e^{24y} - 1} + \frac{3}{e^{36y} - 1} + 2c \right) - 4 \left( \frac{19}{e^{39y} - 1} + \frac{38}{e^{78y} - 1} + 2c \right)$   
 =  $\phi^2(e^{-y}) \phi^2(e^{-19y}) \left\{ 1 + \sqrt{ab} + \sqrt{a-2)(b-2) + \sqrt{ab} + \sqrt{a-2)(b-2) - \sqrt{ab(a-2)(b-2)} \right\}$
- 9. i.  $11 + 12 \left( \frac{1}{e^{6y} - 1} + \frac{2}{e^{12y} - 1} + 2c \right) - 12 \left( \frac{13}{e^{13y} - 1} + \frac{46}{e^{26y} - 1} + 2c \right)$   
 =  $\phi^2(e^{-y}) \phi^2(e^{-23y}) \left\{ \frac{11}{2} (1 + \sqrt{ab} + \sqrt{a-2)(b-2}) - 8 \sqrt{16ab(a-2)(b-2)} (1 + \sqrt{ab} + \sqrt{a-2)(b-2}) - 10 \sqrt{16ab(a-2)(b-2)} \right\}$
- ii.  $7 + 12 \left( \frac{1}{e^{6y} - 1} + \frac{2}{e^{12y} - 1} + 2c \right) - 12 \left( \frac{15}{e^{15y} - 1} + \frac{30}{e^{30y} - 1} + 2c \right)$

$$= \phi^2(e^{-2}) \phi^2(e^{-16}) \left\{ \frac{1}{2} (1 + \sqrt{4\beta} + \sqrt{(1-\alpha)(1-\beta)})^2 - \frac{1 + \sqrt{4\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2} \right\}$$

iii.  $5 + 4 \left( \frac{1}{e^{4\gamma}}, + \frac{2}{e^{8\gamma}}, + 2\epsilon \right) - 4 \left( \frac{31}{e^{64\gamma}}, + \frac{62}{e^{128\gamma}}, + 2\epsilon \right)$

$$= \phi^4(e^{-2}) \phi^4(e^{-32}) \left\{ \frac{1 + \sqrt{4\beta} + \sqrt{(1-\alpha)(1-\beta)}}{2} + (1 + \sqrt{4\beta} + \sqrt{(1-\alpha)(1-\beta)})^2 - 2 \sqrt{4\beta(1-\alpha)(1-\beta)} (1 + \sqrt{4\beta} + \sqrt{(1-\alpha)(1-\beta)}) \right\}$$

10. i.  $1 + 6 \left( \frac{1}{e^{4\gamma}}, + \frac{2}{e^{8\gamma}}, + 2\epsilon \right) - 6 \left( \frac{5}{e^{16\gamma}}, + \frac{10}{e^{32\gamma}}, + 2\epsilon \right)$

$$= \phi^4(e^{-2}) \phi^4(e^{-8}) \sqrt{\left\{ \frac{1 + \alpha\beta + (1-\alpha)(1-\beta)}{2} - \frac{3}{16} (1 - \sqrt{4\beta} - \sqrt{(1-\alpha)(1-\beta)})^2 \right\}}$$

ii.  $1 + 3 \left( \frac{1}{e^{4\gamma}}, + \frac{2}{e^{8\gamma}}, + 2\epsilon \right) - 3 \left( \frac{9}{e^{16\gamma}}, + \frac{18}{e^{32\gamma}}, + 2\epsilon \right)$

$$= \phi^4(e^{-2}) \phi^4(e^{-6}) \sqrt{\left\{ \frac{1 + \alpha\beta + (1-\alpha)(1-\beta)}{2} - \frac{9}{32} (1 - \sqrt{4\beta} - \sqrt{(1-\alpha)(1-\beta)})^2 + \frac{3}{8} \frac{\sqrt{4\beta(1-\alpha)(1-\beta)}}{1 - \sqrt{4\beta} - \sqrt{(1-\alpha)(1-\beta)}} \right\}}$$

iii.  $2 + 3 \left( \frac{1}{e^{4\gamma}}, + \frac{2}{e^{8\gamma}}, + 2\epsilon \right) - 3 \left( \frac{17}{e^{16\gamma}}, + \frac{34}{e^{32\gamma}}, + 2\epsilon \right)$

$$= \phi^4(e^{-2}) \phi^4(e^{-12}) \sqrt{\left\{ 2(1 + \alpha\beta + (1-\alpha)(1-\beta)) - \frac{21}{16} (1 - \sqrt{4\beta} - \sqrt{(1-\alpha)(1-\beta)})^2 - \frac{21}{32} (1 - \sqrt{4\beta} - \sqrt{(1-\alpha)(1-\beta)}) \sqrt{16\alpha\beta(1-\alpha)(1-\beta)} - 3 \sqrt{16\alpha\beta(1-\alpha)(1-\beta)} \right\}}$$

11. i.  $17 + 12 \left( \frac{1}{e^{4\gamma}}, + \frac{2}{e^{8\gamma}}, + 2\epsilon \right) - 12 \left( \frac{35}{e^{16\gamma}}, + \frac{70}{e^{32\gamma}}, + 2\epsilon \right)$

$$= \phi^4(e^{-2}) \phi^4(e^{-25}) \left\{ \frac{(1 - \sqrt{4\beta} - \sqrt{(1-\alpha)(1-\beta)})^3}{2 \sqrt{16\alpha\beta(1-\alpha)(1-\beta)}} + \frac{\sqrt{4\beta} + \sqrt{(1-\alpha)(1-\beta)}}{\sqrt{16\alpha\beta(1-\alpha)(1-\beta)}} \right\}$$

$$\frac{\phi(x) - \phi(-x)}{\phi(x) + \phi(-x)} = \sqrt{\frac{\phi^2(x) - \phi^2(-x)}{\phi^2(x) + \phi^2(-x)}} = \sqrt{\frac{\phi^2(x) - \phi^2(-x)}{\phi^2(x)}}$$

$$\sqrt{\phi(x) + \phi(-x)} = \sqrt{\frac{\phi(x) + \phi(-x)\sqrt{3}}{2}} + \sqrt{\frac{\phi(x) - \phi(-x)\sqrt{3}}{2}}$$

$$y = e^{-x} \left( \frac{1}{3} + \frac{1}{3}x + \frac{1}{6}x^2 + \dots \right)$$

$$1 + 240 \left( \frac{1}{1-y} + \frac{2y}{1-y^2} + \frac{3y^2}{1-y^3} + \dots \right) = 2^2(1+2x)$$

where  $x = 1 + \frac{1}{3}x + \frac{1}{6}x^2 + \dots$

$$1 - 504 \left( \frac{1}{1-y} + \frac{2y}{1-y^2} + \frac{3y^2}{1-y^3} + \dots \right)$$

$$= 2^6(1 - 20x - 9x^2)$$

$$8\sqrt{1-y}(1-y^2)(1-y^4)2x = \frac{2\sqrt{x} \cdot 2\sqrt{1-x} \cdot \sqrt{2}}{\sqrt{3}}$$

$$8\sqrt{1-y}(1-y^2)(1-y^4)(1-y^8)2x = \sqrt{\frac{x}{3}} \cdot \sqrt{1-x} \cdot \sqrt{2}$$

$$1 + 240 \left( \frac{1}{1-y} + \frac{2y}{1-y^2} + \frac{3y^2}{1-y^3} + \dots \right) = 2^4(1 - \frac{2}{3}x)$$

$$1 - 504 \left( \frac{1}{1-y} + \frac{2y}{1-y^2} + \dots \right) = 2^6(1 - \frac{1}{3}x - \frac{2}{3}x^2)$$

$$1 + 6 \left( \frac{1}{e^x + e^{-x}} + \frac{e^x + e^{-x}}{e^{2x} + e^{-2x}} + \dots \right) = 2$$

$$\frac{1}{e^x + e^{-x}} + \frac{2x}{e^{2x} + e^{-2x}} + \frac{3x^2}{e^{3x} + e^{-3x}} + \dots = \frac{2}{27} 2^3$$

$$\frac{1}{e^x + e^{-x}} + \frac{2x}{e^{2x} + e^{-2x}} + \dots = \frac{2}{27} 2^5$$

$$\frac{1}{e^x + e^{-x}} + \frac{2x}{e^{2x} + e^{-2x}} + \dots = \frac{2}{27} (1 + \frac{2}{3}) 2^7$$

$$\frac{1}{e^x + e^{-x}} + \frac{2x}{e^{2x} + e^{-2x}} + \dots = \frac{2}{27} (1 + 2x) 2^9$$

$$\int f(x) = \int_0^{\phi} \left( 1 + \frac{1}{3}x + \frac{1}{6}x^2 + \dots \right) dx = \dots$$

$$\text{then } \phi = 0 + 3 \left\{ \frac{\sin^{-1} \theta}{1 + 2 \cosh y} + \frac{5 - \theta}{2(1 + 2 \cosh y)} + \dots \right\}$$

$$\text{If } d = \frac{p^3(2+p)}{1+2p} \text{ and } B = \frac{27}{4} \cdot \frac{(p+1)^2}{(1+p+p^2)^2}$$

$$\text{then } 1 + \frac{1 \cdot 2}{3^2} d + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} d^2 + 2c$$

$$= \left\{ 1 + \left(\frac{2}{3}\right) d + \left(\frac{1 \cdot 2}{2 \cdot 6}\right)^2 d^2 + 2c \right\} \cdot \frac{1+p+p^2}{\sqrt{1+2p}}$$

$$1 + 6 \left( \frac{2}{1-p} - \frac{2^2}{1-p^2} + \frac{4^2}{1-p^4} - \frac{4^2}{1-p^4} + 2c \right) = 2$$

$$1 + 12 \left( \frac{2}{1-p} + \frac{2 \cdot 2^2}{1-p^2} + \frac{4 \cdot 4^2}{1-p^4} + \frac{4^2}{1-p^4} + 2c \right) = 2^2$$

$$2 = \frac{\psi^3(\psi^2)}{\psi(\psi)} (1+4p+p^2) = 4 \frac{\psi^3(\psi)}{\psi(\psi)} - 3 \frac{\psi^3(\psi)}{\psi(\psi)}$$

$$1 + 4 \left( \frac{x}{1-x} + \frac{2x^2}{1-x^2} + \frac{x^2}{1-x^2} \right)$$

$$\text{If } d = \frac{p \cdot (2+p)^2}{2(1+p)^2} \text{ and } B = \frac{p^2(2+p)}{4}$$

$$\text{or } 1-d = \frac{(1-p)^2(2+p)}{2(1+p)^2} \text{ and } 4-B = \frac{(1-p)(2+p)^2}{2}$$

$$\text{then } 1 + \frac{1 \cdot 2}{3^2} d + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} d^2 + 2c$$

$$= (1+p) \left\{ 1 + \frac{1 \cdot 2}{3^2} d + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} d^2 + 2c \right\}$$

$$1 + \frac{1 \cdot 2}{3^2} \left\{ 1 - \left(\frac{1-p}{1+2c}\right)^2 \right\} + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} \left\{ 1 - \left(\frac{1-p}{1+2c}\right)^2 \right\}^2 + 2c$$

$$= (1+2c) \left\{ 1 + \frac{1 \cdot 2}{3^2} c^2 + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} c^2 + 2c \right\}$$

$$\text{If } d = \frac{27p(1+p)^2}{2(1+4p+p^2)^2} \text{ and } B = \frac{27p^2(1+p)}{2(2+2p-p^2)^2}$$

$$\text{then } \left(1+p - \frac{p^2}{2}\right) \left\{ 1 + \frac{1 \cdot 2}{3^2} d + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} d^2 + 2c \right\}$$

$$= (1+4p+p^2) \left\{ 1 + \frac{1 \cdot 2}{3^2} p + \frac{1 \cdot 2 \cdot 4 \cdot 1^2}{3^2 \cdot 6^2} p^2 + 2c \right\}$$

I degree  $\sqrt[3]{d\beta} + \sqrt[3]{(1-d)(1-\beta)} = 1$   
 $\sqrt[3]{\frac{d^2}{\beta}} + \sqrt[3]{\frac{(1-d)^2}{1-\beta}} = \frac{2}{1+\beta} = \frac{2}{m}$

$\sqrt[3]{\frac{(1-d)^2}{1-d}} - \sqrt[3]{\frac{d^2}{d}} = m$

$\sqrt[3]{\frac{d^2}{\beta}} + \sqrt[3]{\frac{(1-d)^2}{1-\beta}} = \frac{4}{m}$

$\sqrt[3]{\frac{(1-d)^2}{1-d}} + \sqrt[3]{\frac{d^2}{d}} = m^2$

II degree  $\sqrt[3]{d\beta} + \sqrt[3]{(1-d)(1-\beta)} + \sqrt[3]{d\beta(1-d)(1-\beta)} = 1$

III degree  $\sqrt[3]{d\beta} + \sqrt[3]{(1-d)(1-\beta)} + \sqrt[3]{d\beta(1-d)(1-\beta)} + 3\sqrt[3]{\beta^2 d\beta(1-d)(1-\beta)} \left\{ \sqrt[3]{\frac{d}{\beta}} + \sqrt[3]{\frac{(1-d)(1-\beta)}{1-d}} \right\} = 1$

~~IV degree  $\sqrt[4]{(1-d)^3} + \sqrt[4]{\frac{d^3}{d}}$~~

V degree  $m = 3 \cdot \frac{1 + 2\sqrt[3]{d\beta}}{1 + 2\sqrt[3]{1-d}}$

VI degree  $m = \sqrt[3]{\frac{d}{\beta}} + \sqrt[3]{\frac{1-d}{1-\beta}} - \frac{4}{m} \sqrt[3]{\frac{d(1-d)}{\beta(1-\beta)}}$

VII degree  $m = \sqrt[3]{\frac{d}{\beta}} + \sqrt[3]{\frac{1-d}{1-\beta}} - \frac{7}{m} \sqrt[3]{\frac{\beta(1-d)}{\alpha(1-\beta)}} - 3\sqrt[3]{\frac{d(1-d)}{\alpha(1-d)}}$

I, II, IV and VIII

$\frac{1 - \sqrt[3]{d\beta} - \sqrt[3]{(1-d)(1-\beta)}}{2\sqrt[3]{d\beta(1-d)(1-\beta)}} = \frac{1 + \frac{1-d}{3}\beta + 2\alpha + \frac{1+\beta}{3}\sqrt{1-d}}{1 + \frac{1-d}{3}d + 2\alpha + 1 + \frac{1+\beta}{3}d + \dots}$

II, III, VII, VIII or I, IV, V, VI

$\frac{1 + \alpha(\sqrt[3]{d\beta} + \sqrt[3]{(1-d)(1-\beta)})}{1 + 3(\sqrt[3]{d\beta} + \sqrt[3]{(1-d)(1-\beta)})} = \frac{1 + \frac{1-d}{3}\beta + 2\alpha + \frac{1+\beta}{3}\sqrt{1-d}}{1 + \frac{1-d}{3}d + 2\alpha + 1 + \frac{1+\beta}{3}d + \dots}$

$$256$$

$$y = e^{-\frac{27x}{4}} \cdot \frac{1 + \frac{1.3}{4}(1-x) + \frac{1.2.5.7}{4^2.8^2}(1-x)^2 + \dots}{1 + \frac{1.3}{4}x + \frac{1.2.5.7}{4^2.8^2}x^2 + \dots} = F\left(\frac{27x}{4}\right)$$

$$1 + 240 \left( \frac{1^3 y^3}{1-y} + \frac{2^3 y^6}{1-y^2} + \frac{3^3 y^9}{1-y^3} + \dots \right) = 2^4 (1+3x)$$

$$1 - 504 \left( \frac{1^5 y^5}{1-y} + \frac{1^5 y^5}{1-y^2} + \frac{2^5 y^5}{1-y^3} + \dots \right) = 2^6 (1-9x)$$

$$\frac{24 y^3}{\sqrt{y}} (1-y)(1-y^2)(1-y^3) \dots = \frac{24 \sqrt{x} \sqrt{1-x} \sqrt{2}}{\sqrt{2}}$$

$$\frac{192 y^5}{\sqrt{y}} (1-y^2)(1-y^4)(1-y^6) \dots = \frac{192 \sqrt{x} \sqrt{1-x} \sqrt{2}}{\sqrt{2}}$$

$$1 + 240 \left( \frac{1^3 y^3}{1-y^2} + \frac{2^3 y^6}{1-y^4} + \frac{3^3 y^9}{1-y^6} + \dots \right) = 2^4 \left( 1 - \frac{3}{2}x \right)$$

$$1 - 504 \left( \frac{1^5 y^5}{1-y^2} + \frac{1^5 y^5}{1-y^4} + \frac{2^5 y^5}{1-y^6} + \dots \right) = 2^6 \left( 1 - \frac{9}{8}x \right)$$

$$1 + \frac{1.3}{4} \left\{ 1 - \left( \frac{1-t}{1+3t} \right)^2 \right\} + \frac{1.2.5.7}{4^2.8^2} \left\{ 1 - \left( \frac{1-t}{1+3t} \right)^4 \right\} + \dots$$

$$= \sqrt{1+3t} \left\{ 1 + \frac{1.3}{4} t + \frac{1.2.5.7}{4^2.8^2} t^2 + \dots \right\}$$

$$\text{If } \alpha = \frac{64t}{(3+6t-t^2)^2} \text{ and } \beta = \frac{64t^3}{(27+18t-t^2)^2}$$

$$\text{then } \sqrt{1+3t} = \frac{t}{3} \left( 1 + \frac{1.3}{4} \beta + \dots \right)$$

$$= \sqrt{1 - \frac{2}{3}t - \frac{t^2}{27}} \left( 1 + \frac{1.3}{4} \alpha + \dots \right)$$

$$1 + \left( \frac{1}{2} \right)^2 \frac{2x}{1+x} + \left( \frac{1.3}{2.6} \right)^2 \cdot \left( \frac{2x}{1+x} \right)^2 + \dots$$

$$= \sqrt{1+x} \left\{ 1 + \frac{1.3}{4} x + \frac{1.2.5.7}{4^2.8^2} x^2 + \dots \right\}$$

$$\left( x = 1 + \frac{1.3}{4} x + \frac{1.2.5.7}{4^2.8^2} x^2 + \dots \right)$$

III degree  $\sqrt{a\beta} + \sqrt{(1-a)(1-\beta)} + 4\sqrt[4]{a\beta(1-a)(1-\beta)} = 1$

VII degree  $\sqrt{a\beta} + \sqrt{(1-a)(1-\beta)} + 20\sqrt[5]{a\beta(1-a)(1-\beta)} + 8\sqrt{2}\sqrt[8]{a\beta(1-a)(1-\beta)} (\sqrt[4]{a\beta} + \sqrt[4]{(1-a)(1-\beta)}) = 1$

V degree  $\sqrt{a\beta} + \sqrt{(1-a)(1-\beta)} + 8\sqrt[6]{a\beta(1-a)(1-\beta)} (\sqrt[3]{a\beta} + \sqrt[3]{(1-a)(1-\beta)}) = 1$

XI degree  $\sqrt{a\beta} + \sqrt{(1-a)(1-\beta)} + 68\sqrt[7]{a\beta(1-a)(1-\beta)} + 16\sqrt[14]{a\beta(1-a)(1-\beta)} (\sqrt[7]{a\beta} + \sqrt[7]{(1-a)(1-\beta)}) + 48\sqrt[6]{a\beta(1-a)(1-\beta)} (\sqrt[3]{a\beta} + \sqrt[3]{(1-a)(1-\beta)}) = 1$

III degree  $m^2 = \sqrt{\frac{a}{2}} + \sqrt{\frac{1-a}{1-a}} - \frac{9}{m} \sqrt{\frac{a(1-a)}{2(1-a)}}$

V degree  $m = \sqrt[4]{\frac{a}{2}} + \sqrt[4]{\frac{1-a}{1-a}} - \frac{5}{m} \sqrt[5]{\frac{a(1-a)}{2(1-a)}}$

IX degree  $\sqrt[9]{m} = \sqrt[9]{\frac{a}{2}} + \sqrt[9]{\frac{1-a}{1-a}} - \frac{3}{\sqrt[9]{m}} \sqrt[9]{\frac{a(1-a)}{2(1-a)}}$

VII degree  $m^2 = \sqrt{\frac{a}{2}} + \sqrt{\frac{1-a}{1-a}} - \frac{49}{m} \sqrt{\frac{a(1-a)}{2(1-a)}} - 8\sqrt[6]{\frac{a(1-a)}{2(1-a)}} (\sqrt[3]{\frac{a}{2}} + \sqrt[3]{\frac{1-a}{1-a}})$

XIII degree  $m = \sqrt[4]{\frac{a}{2}} + \sqrt[4]{\frac{1-a}{1-a}} - \frac{13}{m} \sqrt[5]{\frac{a(1-a)}{2(1-a)}} - 4\sqrt[13]{\frac{a(1-a)}{2(1-a)}} (\sqrt[5]{\frac{a}{2}} + \sqrt[5]{\frac{1-a}{1-a}})$

XXV degree  $\sqrt[25]{m} = \sqrt[25]{\frac{a}{2}} + \sqrt[25]{\frac{1-a}{1-a}} - \frac{5}{\sqrt[25]{m}} \sqrt[25]{\frac{a(1-a)}{2(1-a)}} - 2\sqrt[25]{\frac{a(1-a)}{2(1-a)}} (\sqrt[5]{\frac{a}{2}} + \sqrt[5]{\frac{1-a}{1-a}})$



$$\text{ff } y = e^{-2\pi} \frac{1 + \frac{1.5}{6}(1-x) + \frac{1.5 \cdot 7 \cdot 11}{6^2 \cdot 12^2}(1-x)^2 + \dots}{1 + \frac{1.5}{6}x + \frac{1.5 \cdot 7 \cdot 11}{6^2 \cdot 12^2}x^2 + \dots}$$

and  $z = 1 + \frac{1.5}{6}x + \frac{1.5 \cdot 7 \cdot 11}{6^2 \cdot 12^2}x^2 + \dots$ , then

$$1 + 240 \left( \frac{1^2 y}{1-y} + \frac{2^2 y^2}{1-y^2} + \frac{3^2 y^3}{1-y^3} + \dots \right) = z^4.$$

$$1 - 504 \left( \frac{1^5 y}{1-y} + \frac{2^5 y^2}{1-y^2} + \frac{3^5 y^3}{1-y^3} + \dots \right) = z^6(1-2x)$$

$$\frac{24/y}{(1-y)(1-y^2)(1-y^3)(1-y^4)} z^4 = \sqrt{\frac{24x(1-x)}{432}} \sqrt{2}.$$

ff  $u = x(1-x)$  and  $v = y(1-y)$

and  $u = \frac{27v^2}{16(1-v)^3}$ , then

$$1 + \frac{1.5}{6}x + \frac{1.5 \cdot 7 \cdot 11}{6^2 \cdot 12^2}x^2 + \dots = \left\{ 1 + \left(\frac{1.5}{6}\right)^2 y + \left(\frac{1.5 \cdot 7 \cdot 11}{6^2 \cdot 12^2}\right)^2 y^2 + \dots \right\}^4 \sqrt{1-y+y^2}.$$


ff  $y = \frac{p(2+p)}{1+2p}$ , then  $x = \frac{27}{4} \cdot \frac{(p+p^2)^2}{(1+p+p^2)^3}$

$$\sqrt{21} \cdot \frac{1}{2} \cdot \left(\frac{2-\sqrt{7}}{\sqrt{2}}\right)^2 \left(\frac{\sqrt{5+\sqrt{7}}}{4} - \frac{\sqrt{1+\sqrt{7}}}{4}\right)^4 \left(\frac{\sqrt{2+\sqrt{7}}}{4} - \frac{\sqrt{\sqrt{7}}}{4}\right)^4 \left(\frac{\sqrt{7}-\sqrt{3}}{2}\right)^2$$

$$\sqrt{33} \cdot \frac{1}{2} (2-\sqrt{3})^3 \left(\frac{\sqrt{7+3\sqrt{3}}}{4} - \frac{\sqrt{3+3\sqrt{3}}}{4}\right)^4 \left(\frac{\sqrt{5+\sqrt{3}}}{4} - \frac{\sqrt{1+\sqrt{3}}}{4}\right)^4 \left(\frac{2\sqrt{3}-1}{\sqrt{2}}\right)^2$$

$$\sqrt{45} \cdot \frac{1}{2} (\sqrt{5}-2)^3 \left(\frac{\sqrt{7+3\sqrt{5}}}{4} - \frac{\sqrt{3+3\sqrt{5}}}{4}\right)^4 \left(\frac{\sqrt{2+\sqrt{5}}}{2} - \frac{\sqrt{1+\sqrt{5}}}{2}\right)^4 \left(\frac{\sqrt{5}-\sqrt{3}}{\sqrt{2}}\right)^2$$

$$\sqrt{15} \cdot 2 \cdot \frac{1}{16} \cdot \left(\frac{\sqrt{5}-1}{2}\right)^6 \cdot (2-\sqrt{3})^2 (4-\sqrt{15})^2.$$



$$PQ^2 = 2 BP \cdot QC.$$

$$(\alpha + b - \sqrt{\alpha^2 + b^2})^2 = 2(\sqrt{\alpha^2 + b^2} - \alpha)(\sqrt{\alpha^2 + b^2} - b)$$

$$\left\{ \sqrt[3]{(\alpha + b)^3} - \sqrt[3]{\alpha^3 - \alpha b + b^3} \right\}^3 = 3(\sqrt[3]{\alpha^2 + b^2} - \alpha)(\sqrt[3]{\alpha^2 + b^2} - b)$$

$$\sqrt{A + B\sqrt[3]{p}} = \sqrt{\frac{B}{p + k^3}} \left( \frac{k^2}{2} + k\sqrt[3]{p} - \sqrt[3]{k^2} \right)$$

where  $Bk^4 - 4Ak^3 - 8Bkp - 4Ap = 0$ .

F.  $\frac{1 - \sqrt{1 - t^{24}}}{2} = e^{-\pi\sqrt{29}}$ . Then

$$t^{24} + 9t^{20} + 5t^{16} - 2t^{12} - 5t^8 + 9t^4 - 1 = 0$$

$$\frac{t^6 + t^2}{1 - t^4} = \sqrt{\frac{\sqrt{29} - 5}{2}}$$

$$\frac{t^3 + t\sqrt{\sqrt{29} - 2}}{1 + t^2\sqrt{\sqrt{29} + 2}} = \sqrt[4]{\frac{\sqrt{29} - 5}{2}}$$

If  $\sqrt[4]{1 - t^8} = t(1 + u^2)$ , then  $u^3 + u = \sqrt{2}$ .

F.  $\frac{1 - \sqrt{1 - \frac{1}{64}t^{24}}}{2} = e^{-\pi\sqrt{79}}$ . Then

$$t^5 - t^4 + t^3 - 2t^2 + 3t - 1 = 0.$$

F.  $\frac{1 - \sqrt{1 - \frac{1}{64}t^{24}}}{2} = e^{-\pi\sqrt{47}}$ , then

$$t^5 + 2t^4 + 2t^3 + t^2 - 1 = 0$$

$$(1) \phi^2(x) = 1 - \frac{4x}{1+x} + \frac{4x^2}{1+x^2} - \frac{4x^4}{1+x^4} + \frac{4x^{10}}{1+x^4} - \dots$$

$$(2) \psi(x)\phi(x) = \frac{1+x}{1-x} - x \cdot \frac{1+x^3}{1-x^3} + x^3 \cdot \frac{1+x^5}{1-x^5} - x^6 \cdot \frac{1+x^7}{1-x^7} + \dots$$

$$(3) \psi^2(x) = \frac{1+x}{1-x} - x^2 \cdot \frac{1+x^3}{1-x^3} + x^6 \cdot \frac{1+x^5}{1-x^5} - x^{12} \cdot \frac{1+x^7}{1-x^7} + \dots$$

$$(4) \frac{x}{1-x} + \frac{2x^2}{1-x^2} + \frac{3x^3}{1-x^3} + \frac{4x^6}{1-x^4} + \dots$$
  
$$= x \cdot \frac{1+x}{(1-x)^2} - x^3 \cdot \frac{1+x^2}{(1-x^2)^2} + x^6 \cdot \frac{1+x^3}{(1-x^3)^2} - x^{10} \cdot \frac{1+x^4}{(1-x^4)^2} + \dots$$

$$(5) x\psi(x)\psi(x) = \frac{x}{1-x} - \frac{x^3}{1-x^3} + \frac{x^6}{1-x^5} - \frac{x^{10}}{1-x^7} + \dots$$

$$(6) \frac{x}{1+x} - \frac{3^2 x^3}{1+x^2} + \frac{3^4 x^6}{1+x^3} - \frac{4^2 x^{10}}{1+x^4} + \dots$$
  
$$= \phi^2(x) \left\{ x \cdot \frac{1+x}{(1-x)^2} + x^6 \cdot \frac{1+x^3}{(1-x^3)^2} + x^{10} \cdot \frac{1+x^5}{(1-x^5)^2} + \dots \right\}$$

$$(7) \frac{1+x}{1-x} - 3^2 x^2 \cdot \frac{1+x^3}{1-x^3} + 5^2 x^6 \cdot \frac{1+x^5}{1-x^5} - \dots$$
  
$$= \psi^2(x) \left\{ 1 - \frac{8x^2}{(1+x)^2} + \frac{8x^6}{(1+x^3)^2} - \frac{8x^{12}}{(1+x^5)^2} + \dots \right\}$$

$$(8) \frac{x^2}{(1-x)^2} - \frac{3x^6}{(1-x^3)^2} + \frac{5x^{12}}{(1-x^5)^2} - \frac{7x^{20}}{(1-x^7)^2} + \dots$$
  
$$= x\psi^2(x) \left\{ x \cdot \frac{1+x^2}{1-x^2} - 3x^4 \cdot \frac{1+x^6}{1-x^6} + 3x^9 \cdot \frac{1+x^8}{1-x^8} + \dots \right\}$$

$$(9) x \cdot \frac{1-x}{(1+x)^2} - 2x^3 \cdot \frac{1-x^2}{(1+x^3)^2} + 3x^6 \cdot \frac{1-x^3}{(1+x^5)^2} - \dots$$
  
$$= \phi^2(x) \left( \frac{x}{1-x} + \frac{2x^3}{1-x^2} + \frac{3x^6}{1-x^3} + \frac{4x^{10}}{1-x^4} + \dots \right)$$

$$\frac{1}{2 \log 2} + \frac{1}{3 \log 3} + \dots + \frac{1}{x \log x}$$

$$= .1015314 + \log \log(x^{-1} + x + \theta)$$

$$x = \infty \quad \theta = \frac{1}{3}$$

$$x = 1 \quad \theta = .46811$$

$$\int_0^{\infty} e^{-nx} \sin nx \cot x dx$$

$$= \frac{\pi^2}{2} \left\{ \frac{1}{1^2 + \frac{n^2}{2}} + \frac{2}{2^2 + \frac{n^2}{2}} + \frac{3}{3^2 + \frac{n^2}{2}} \dots \right\}$$

$$\int_0^{\infty} e^{-x} \frac{\sin^2 x}{x} \left\{ A_0 - \frac{2\beta_2 A_2 x^2}{L^2} - \frac{2^2 \beta_4 A_4 x^4}{L^4} \dots \right\}$$

130489 quadrats  
 - lion wants to  
 get the value  
 of 5

$$= (A_2 - A_6 + A_{10} - \dots)$$

$$+ (2A_2 - 5A_6 + 2A_{10} - \dots)$$

$$+ (A_2 - 7A_6 + 3^2 A_{10} - \dots)$$

$$\int_0^{\infty} e^{-ax} e^{-bx} \cot x dx$$

$$= \frac{1}{2a} + 2a \left\{ \frac{1}{a^2 + (b+1)^2} + \frac{1}{a^2 + (b+3)^2} \dots \right\}$$

$$\int_0^{\infty} e^{-ax} \sin ax (\cot x + \coth x) dx$$

$$= \frac{\pi}{2} \frac{\sinh \pi a}{\cosh \pi a - \cos \pi}$$

*nbv*  
If  $x + na^2 = y + nab = z + nb^2 = (a+b)^2$

then  $x^2 + (n-2)xy + y^2 = ny^2$

---

If  $p, q, r$  are quantities so taken that

$$p + 3a^2 = q + 3ab = r + 3b^2 = (a+b)^2$$

and  $m$  &  $n$  are any two quantities, then

$$\begin{aligned} n(mp + nq)^3 + m(mq + nr)^3 \\ = m(np + mq)^3 + n(nq + mr)^3 \end{aligned}$$

---

A particular case of the above theorem is

$$\begin{aligned} (3a^2 + 5ab - 5b^2)^3 + (4a^2 - 4ab + 6b^2)^3 + (3a^2 + ab - 7b^2)^3 \\ = 6a^2 + 11ab + \dots \end{aligned}$$

$$\begin{aligned} (3a^2 + 5ab - 5b^2)^3 + (4a^2 - 4ab + 6b^2)^3 + (5a^2 - 5ab - 3b^2)^3 \\ = (6a^2 + 4ab + 6b^2)^3 \end{aligned}$$

$$\begin{aligned} (2x^2 + 3xy + 5y^2)(2p^2 + 3pq + 5q^2) \\ = 2u^2 + 3uv + 5v^2 \quad \text{where} \end{aligned}$$

$$u = \frac{5}{2}(x+y)(p+q) - 2xp \quad \& \quad v = 2qy - \frac{(x+y)^2}{2} + (p+q)^2$$

203 284

Let  $I(P)$  be the integer equal to or just less than  $P$  and  $G(P)$ , equal to or just greater than  $P$ , and let  $N(P)$  be the nearest integer to  $P$ . Then,

(1)  $N(P) = I(P + \frac{1}{2})$ .

(2)  $I(\frac{P}{q})$  is the coeff<sup>t</sup> of  $x^n$  in  $\frac{x^P}{(1-x)(1-x^q)}$

(3)  $I \phi(n)$  is the coeff<sup>t</sup> of  $x^n$  in  $\sum_{m=1}^{n=\infty} \frac{x^{G_m} \phi^{-1}(G_m)}{1-x}$

(4) The sum of the nos<sup>s</sup> of divisors of  $P$  natural nos.

$$= I(\frac{P}{1}) + I(\frac{P}{2}) + I(\frac{P}{3}) + I(\frac{P}{4}) + \dots + I(\frac{P}{P})$$

$$= 2 \{ I(\frac{P}{1}) + I(\frac{P}{2}) + \dots \text{ to } I(\sqrt{P}) \text{ terms} \} - (I \sqrt{P})^2$$

(5) The above sum is odd or even according as  $I(\sqrt{P})$  is odd or even and is approximately equal to  $P(2c-1) \log P + \frac{1}{2}$  the no of factors of  $P + \frac{1}{2}$ .

(6) If  $n > \sqrt{P}$  and  $m = I(\frac{P}{n})$ , then

$$I(\frac{P}{1}) + I(\frac{P}{2}) + I(\frac{P}{3}) + \dots + I(\frac{P}{n})$$

$$= n I(\frac{P}{n}) + I(\frac{P}{1+m}) + I(\frac{P}{2+m}) + I(\frac{P}{3+m}) + \dots + I(\frac{P}{P})$$

(7) If  $P$  be the  $n$ th Prime no. then  $\frac{dP}{dn} = \log P$  nearly and hence  $n = \frac{P}{\log P - 1}$  nearly.

(8)  $\phi(2) + \phi(3) + \phi(5) + \phi(7) + \phi(11) + \dots$  and  $\frac{\phi(2)}{\log 2} + \frac{\phi(3)}{\log 3} + \frac{\phi(4)}{\log 4} + \dots$  are both convergent or both divergent.

$$(1) \text{ If } \alpha\beta = \pi^2, \text{ then } \frac{1}{\sqrt{\alpha}} \left\{ 1 + 4\alpha \int_0^{\infty} \frac{x e^{-\alpha x^2}}{e^{2\pi x} - 1} dx \right\}$$

$$= \frac{1}{\sqrt{\beta}} \left\{ 1 + 4\beta \int_0^{\infty} \frac{x e^{-\beta x^2}}{e^{2\pi x} - 1} dx \right\} = \sqrt{\frac{1}{\alpha} + \frac{1}{\beta} + \frac{2}{3}} \text{ nearly}$$

$$(2) \text{ If } \alpha\beta = \pi^2, \text{ then}$$

$$\frac{1}{\sqrt{\alpha}} \left\{ \phi(0) + \frac{\alpha}{1} \phi(2) B_2 - \frac{\alpha^2}{2} \phi(4) B_4 + \frac{\alpha^3}{3} \phi(6) B_6 - \dots \right\}$$

$$= \frac{1}{\sqrt{\beta}} \left\{ \phi(0) + \frac{\beta}{1} \phi(-1) B_2 - \frac{\beta^2}{2} \phi(-3) B_4 + \frac{\beta^3}{3} \phi(-5) B_6 - \dots \right\}$$

$$(3) \text{ If } \alpha\beta = 4\pi^2, \text{ then } 2\alpha^{\frac{m+1}{2}} \int_0^{\infty} \frac{x^m}{e^{2\pi x} - 1} \frac{dx}{e^{\alpha x}} =$$

$$\alpha^{\frac{m-1}{2}} \left\{ \frac{B_m}{m} - \frac{\alpha}{2} \frac{B_{m+1}}{m+1} + \alpha^2 \frac{B_2}{2} \frac{B_{m+2}}{m+2} - \alpha^3 \frac{B_4}{4} \frac{B_{m+4}}{m+4} + \dots \right\}$$

$$= \beta^{\frac{m-1}{2}} \left\{ \frac{B_m}{m} - \frac{\beta}{2} \frac{B_{m+1}}{m+1} + \beta^2 \frac{B_2}{2} \frac{B_{m+2}}{m+2} - \beta^3 \frac{B_4}{4} \frac{B_{m+4}}{m+4} + \dots \right\}$$

$$(4) \text{ If } \alpha\beta = \pi^2, \text{ then } -\frac{\pi}{2} \frac{\alpha^{\frac{x}{2}}}{\sin \frac{\pi x}{2}} \frac{B_x}{\Gamma \frac{x}{2}} \phi(x) =$$

$$\frac{\phi(0)}{x} - \frac{\alpha}{4} \frac{\phi(2)}{2-x} B_2 + \frac{\alpha^2}{12} \frac{\phi(4)}{4-x} B_4 - \frac{\alpha^3}{12} \frac{\phi(6)}{6-x} B_6 + \dots$$

$$+ \sqrt{\frac{\alpha}{\pi}} \left\{ \frac{\phi(1)}{1-x} - \frac{\beta}{4} \frac{\phi(-1)}{1+x} B_2 + \frac{\beta^2}{12} \frac{\phi(-3)}{3+x} B_4 - \dots \right\}$$

$$(5) \frac{\pi}{2} \frac{\alpha^x B_x \phi(x)}{\sin \frac{\pi x}{2}} + \frac{\phi(0)}{x} + \frac{\alpha \phi(1)}{2(1-x)} =$$

$$\frac{\alpha^2 \phi(2) B_2}{2-x} - \frac{\alpha^4 \phi(4) B_4}{4-x} + \frac{\alpha^6 \phi(6) B_6}{6-x} - \dots$$

$$+ \frac{3_2 \phi(-1)}{\alpha(1+x)} - \frac{2_3 \phi(-2)}{\alpha^2(2+x)} + \frac{3_4 \phi(-3)}{\alpha^3(3+x)} - \dots$$

(1) If  $\beta = 4\pi^2$ , then

$$\sqrt{\beta^n} \left\{ \frac{B_n}{2^n} + \cos \frac{\pi n}{2} \left( \frac{1^{n-1}}{e^{\beta} - 1} + \frac{2^{n-1}}{e^{2\beta} - 1} + \frac{3^{n-1}}{e^{3\beta} - 1} + \dots \right) \right\}$$

$$= \sqrt{\beta^n} \left\{ \frac{B_n}{2^n} \cos \frac{\pi n}{2} - \sin \frac{\pi n}{2} \int_0^{\infty} \frac{x^{n-1} \cot \frac{\beta x}{2}}{e^{2\pi x} - 1} dx + \right.$$

$$\left. \frac{1^{n-1}}{e^{\beta} - 1} + \frac{2^{n-1}}{e^{2\beta} - 1} + \frac{3^{n-1}}{e^{3\beta} - 1} + \dots \right\}$$

2).  $\frac{1^{n+1}}{1^4 + 4x^4} + \frac{2^{n+1}}{2^4 + 4x^4} + \frac{3^{n+1}}{3^4 + 4x^4} + \frac{4^{n+1}}{4^4 + 4x^4} + \dots$

$$= \frac{\pi}{4} (x\sqrt{2})^{n-2} \sec \frac{\pi n}{4} - 2 \cos \frac{\pi n}{2} \int_0^{\infty} \frac{x^{n+1}}{e^{2\pi x} - 1} \cdot \frac{dx}{2^4 + 4x^4}$$

$$+ \frac{\pi}{2} (x\sqrt{2})^{n-2} \frac{\cos(\frac{\pi n}{4} + 2\pi x) - e^{-2\pi x} \cos \frac{\pi n}{4}}{\cosh 2\pi x - \cos 2\pi x}$$

(3).  $\int_0^{\infty} \frac{x \sin 2\pi x}{e^{x^2} - 1} dx = \frac{n\sqrt{\pi}}{2} \left( \frac{e^{-n^2}}{1\sqrt{1}} + \frac{e^{-\frac{n^2}{2}}}{2\sqrt{2}} + \dots \right)$

$$= \frac{\pi}{2} \left( 1 + 2e^{-2n^2\sqrt{\pi}} \cos 2n\sqrt{\pi} + 2e^{-2n^2\sqrt{2\pi}} \cos 2n\sqrt{2\pi} + \dots \right)$$

4).  $\int_0^{\infty} \frac{x \sin 2\pi x}{e^{x^2} + e^{-x^2}} dx = \frac{n\sqrt{\pi}}{2} \left( e^{-n^2} \frac{e^{-\frac{n^2}{3}}}{3\sqrt{3}} + \frac{e^{-\frac{n^2}{5}}}{5\sqrt{5}} + \dots \right)$

$$= \frac{\pi}{2} \left( e^{-n^2\sqrt{\pi}} \sin n\sqrt{\pi} - e^{-n^2\sqrt{3\pi}} \sin n\sqrt{3\pi} + \dots \right)$$

(5) If  $n$  is a positive integer, then

$$\frac{1^{4n}}{(e^{\pi} - e^{-\pi})^2} + \frac{2^{4n}}{(e^{2\pi} - e^{-2\pi})^2} + \frac{3^{4n}}{(e^{3\pi} - e^{-3\pi})^2} + \dots =$$

$$\frac{\pi}{11} \left( \frac{B_{4n}}{8n} + \frac{1^{4n-1}}{e^{2\pi}} + \frac{2^{4n-1}}{e^{4\pi}} + \frac{3^{4n-1}}{e^{6\pi}} + \dots \right)$$



$$(1) \frac{1}{p+1} + \frac{1}{(p+2)^2} + \frac{2}{(p+3)^3} + \frac{4^2}{(p+4)^4} + \frac{5^3}{(p+5)^5} + \dots$$

$$= \frac{1-e^{-p}}{p} + e^{-(p+1)} \left\{ \frac{1}{p+1} \right\} + \frac{e^{-(p+2)}}{2} \left\{ \frac{1}{p+2} + \frac{1}{(p+2)^2} \right\}$$

$$+ \frac{3e^{-(p+3)}}{6} \left\{ \frac{1}{p+3} + \frac{2}{(p+3)^2} + \frac{2}{(p+3)^3} \right\} + \dots$$

the  $n^{\text{th}}$  term within the brackets being  $\frac{1}{p+n} + \frac{n-1}{(p+n)^2}$

$$+ \frac{(n-1)(n-2)}{(p+n)^3} + \frac{(n-1)(n-2)(n-3)}{(p+n)^4} + \frac{(n-1)(n-2)(n-3)(n-4)}{(p+n)^5} + \dots$$

$$= \frac{1}{p} - \frac{1}{p^2} + \frac{2}{p^3} - \frac{6}{p^4} + \frac{24}{p^5} - \frac{120}{p^6} + \dots$$

$$- n \left( \frac{1}{p^3} - \frac{5}{p^4} + \frac{36}{p^5} - \frac{154}{p^6} + \dots \right)$$

$$+ n^2 \left( \frac{3}{p^5} - \frac{35}{p^6} + \frac{340}{p^7} - \frac{3304}{p^8} + \dots \right)$$

$$- n^3 \left( \frac{15}{p^7} - \frac{315}{p^8} + \dots \right) + \dots$$

$$154 = 4 \cdot 6 + 5 \cdot 26; \quad 340 = 5 \cdot 26 + 6 \cdot 35; \quad 3304 = 6 \cdot 154 + 7 \cdot 340$$

&c      &c      &c

$$1, \quad 1, \quad 1, \quad 1, \quad 1, \quad \dots$$

$$\frac{1}{2}, \quad \frac{1}{2} + \frac{1}{3}, \quad \frac{1}{2} + \frac{1}{3} + \frac{1}{4}, \quad \dots$$

$$\frac{1}{2} \cdot \frac{1}{3}, \quad \frac{1}{2} \cdot \frac{1}{3} + \left( \frac{1}{2} + \frac{1}{3} \right) \frac{1}{5}, \quad \dots$$

$$\frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4}, \quad \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4} + \left\{ \frac{1}{2} \cdot \frac{1}{3} + \left( \frac{1}{2} + \frac{1}{3} \right) \frac{1}{5} \right\} \frac{1}{7}, \quad \dots$$

&c      &c      &c      &c      &c

$$\begin{aligned}
 (1) \quad & \frac{1}{p+1} + \frac{1}{(p+2)^2} + \frac{1}{(p+3)^3} + \frac{1}{(p+4)^4} + \frac{1}{(p+5)^5} + \dots \\
 & = \frac{1-e^{-p}}{p} + e^{-p} \left\{ \frac{1}{p+1} - \frac{1}{(p+1)(p+2)} + \frac{1}{2(p+1)(p+2)(p+4)} \right. \\
 & \quad \left. - \frac{1}{(p+1)(p+2)(p+4)(3p+23+\theta)} \right\}
 \end{aligned}$$

where  $\theta_{-1} = -2.5856$ ;  $\theta_0 = .0069$ ;  $\theta_1 = .4137$   
 and  $\theta_{\infty} = \frac{3}{5}$ .

$$\begin{aligned}
 (2) \quad & \frac{1}{a(p+a)} + \frac{1}{(p+a+1)^2} + \frac{a+2}{(p+a+2)^3} + \frac{(a+3)^4}{(p+a+3)^4} + \dots \\
 & = u_0(a) - \frac{p}{a} u_1(a) + \frac{p^2}{a^2} u_2(a) - \frac{p^3}{a^3} u_3(a) + \dots
 \end{aligned}$$

where  $u_n(a) = \frac{1}{a^{n+1}} + \frac{1}{(a+1)^{n+2}} + \frac{1}{(a+2)^{n+3}} + \dots$

and  $\frac{u_{n-1}(a) - u_n(a+1)}{u_n(a) - u_{n+1}(a+1)} = \frac{a}{n}$ .

$$\begin{aligned}
 (4) \quad u_n(a) &= \frac{1}{a(n+1)} + \frac{1}{2a^2} + \frac{1}{a^3} \left\{ \frac{1}{6} + \frac{n}{4} \right\} + \frac{1}{a^4} \left\{ \frac{n(n-1)}{24} + \frac{n(n-1)(n-2)}{8} \right\} \\
 &+ \frac{1}{a^5} \left\{ -\frac{1}{30} + \frac{n}{12} + \frac{n(n-1)}{4} + \frac{n(n-1)(n-2)}{16} \right\} \\
 &+ \frac{1}{a^6} \left\{ -\frac{n}{12} + \frac{5n(n-1)}{24} + \frac{5n(n-1)(n-2)}{24} + \frac{n(n-1)(n-2)(n-3)}{32} \right\} \\
 &+ \frac{1}{a^7} \left\{ \frac{1}{42} - \frac{n}{12} - \frac{n(n-1)}{18} + \frac{5n(n-1)(n-2)}{16} + \right. \\
 &\quad \left. \frac{5n(n-1)(n-2)(n-3)}{32} + \frac{n(n-1)(n-2)(n-3)(n-4)}{64} \right\} \\
 &+ \frac{1}{a^8} \left\{ \frac{n}{12} - \frac{7n(n-1)}{24} + \frac{7n(n-1)(n-2)}{72} + \frac{35n(n-1)(n-2)(n-3)}{96} \right. \\
 &\quad \left. + \frac{7n(n-1)(n-2)(n-3)(n-4)}{64} + \frac{n(n-1)(n-2)(n-3)(n-4)(n-5)}{128} \right\} \\
 &+ \dots \quad \quad \quad \dots \quad \quad \quad \dots
 \end{aligned}$$

$$(5) \frac{1}{a(2p+a)} + \frac{1}{(2p+a)^2} + \frac{a+2}{(2p+a+2)^3} + \frac{(a+3)^4}{(2p+a+3)^4} + \dots$$

$$= \frac{1}{2ap} - e^{-2p} \left\{ \frac{1}{2p(a+p)} - \frac{P_2}{(a+p)^3} + \frac{P_4}{(a+p)^5} - \dots \right\}$$

$$P_2 = \frac{1}{6}$$

$$P_4 = \frac{1}{30} + \frac{p}{6}$$

$$P_6 = \frac{1}{42} + \frac{p}{6} + \frac{5p^2}{18}$$

$$P_8 = \frac{1}{30} + \frac{3p}{10} + \frac{7p^2}{9} + \frac{35p^3}{54}$$

$$P_{10} = \frac{5}{66} + \frac{5p}{6} + \frac{17p^2}{6} + \frac{35p^3}{9} + \frac{35p^4}{18}$$

$$P_{12} = \frac{691}{2730} + \frac{691p}{210} + \frac{616p^2}{45} + \frac{451p^3}{18} + \frac{385p^4}{18} + \frac{385p^5}{54}$$

$$P_{14} = \frac{7}{6} + \frac{35p}{2} + \frac{7709p^2}{90} + \frac{26026p^3}{185} + \frac{2002p^4}{9} + \frac{7007p^5}{54}$$

$$\dots \dots \dots + \frac{5005p^6}{162}$$

$$P_{2n} = \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{2 \cdot 3^n} \left\{ p^{n+1} + \frac{n(n-1)}{10} p^{n-2} + \right.$$

$$\frac{n(n-1)(n-2)}{200} \left[ (n-3) + \frac{20}{7} \right] p^{n-3} +$$

$$\frac{n(n-1)(n-2)(n-3)}{6000} \left[ (n-4)(n-5) + \frac{60}{7}(n-4) + \frac{90}{7} \right] p^{n-4} +$$

$$\frac{n(n-1)(n-2)(n-3)(n-4)}{240000} \left[ \frac{(n-5)(n-6)(n-7) + \frac{130}{7}(n-5)(n-6)}{+ \frac{3730}{49}(n-5) + \frac{6000}{77}} \right] p^{n-5}$$

$$+ \dots \dots \dots \left. \right\}$$

$$\begin{aligned}
 P_n = & B_n + (n+1) B_{n-1} p + \left\{ \frac{(n+1)(n+2)}{3} B_{n-2} - \frac{n(n-1)}{6} B_{n-2} \right\} p^2 \\
 & + \left\{ \frac{(n+1)(n+2)(n+3)}{18} B_{n-3} - \frac{n^2(n-1)}{9} B_{n-2} \right\} p^3 + \\
 & \left\{ \frac{(n+1)(n+2)(n+3)(n+4)}{180} B_{n-4} - \frac{n^2(n-1)}{36} B_{n-2} + \frac{n(n-1)(n-2)(n-3)}{120} B_{n-4} \right\} p^4 \\
 & + \left\{ \frac{(n+1)(n+2)(n+3)(n+4)(n+5)}{2700} B_{n-5} - \frac{n^2(n-1)(n+2)}{270} B_{n-2} \right. \\
 & \left. + \frac{n(n-1)(n-2)(n-3)(23n-25)}{5400} B_{n-4} \right\} p^5 + \&c
 \end{aligned}$$

which is got from  $\frac{(a-p+n)^{n-1}}{(a+p+n)^{n+1}} = \frac{1}{(a+n)^{-1}} \exp\left\{\frac{2ap}{a+n}\right\}$

$$\begin{aligned}
 & + \left\{ \frac{p^2}{(a+n)^{-1}} - \frac{2np^3}{3(a+n)^3} + \frac{p^4}{2(a+n)^4} - \frac{2np^5}{5(a+n)^5} + \frac{p^6}{3(a+n)^6} - \&c \right\} \\
 = & 1 + 2p \cdot \frac{a}{a+n} + 2p^2 \cdot \frac{a^2 + \frac{1}{2}}{(a+n)^{-1}} + \frac{4p^3}{3} \left\{ \frac{a^3 + 2a}{(a+n)^2} - \frac{1}{2(a+n)} \right\} \\
 & + \frac{2p^4}{3} \left\{ \frac{a^4 + 5a^2 + \frac{7}{2}}{(a+n)^4} - \frac{2a}{(a+n)^3} \right\} + \\
 & \frac{4p^5}{15} \left\{ \frac{a^5 + 10a^3 + \frac{23}{2}}{(a+n)^5} - \frac{5a^2 + 4}{(a+n)^4} \right\} + \&c.
 \end{aligned}$$

$$\begin{aligned}
 6). & \frac{\text{Cott } \pi}{p^n} + \frac{\text{Cott } 2\pi}{2^n} + \frac{\text{Cott } 3\pi}{3^n} + \frac{\text{Cott } 4\pi}{4^n} + \&c \\
 = & \frac{1}{2} \left( \frac{3}{\pi} S_{n+1} + \frac{\pi}{8} S_{n-1} \right) + \frac{2^{n-3} \cdot \pi \cdot n \cdot v_{n+1}}{n-3} \text{ where}
 \end{aligned}$$

$$v_4 = -\frac{3}{2}, v_8 = 0, v_{12} = \frac{1}{2730}, v_{16} = \frac{1}{340}$$

$$v_{20} = \frac{191}{2310}, v_{24} = \frac{907}{294}, \&c \ \&c.$$

270

$$(1) \frac{\theta}{\sqrt{\pi}} B_2 - \frac{\theta^3}{\sqrt{3}} B_6 + \frac{\theta^5}{\sqrt{5}} B_{10} - \dots$$

$$= \sqrt{\frac{\theta}{2\pi}} \left\{ 1 + \frac{\pi^4}{\theta^2 \sqrt{2}} B_4 - \frac{\pi^8}{\theta^4 \sqrt{4}} B_8 + \frac{\pi^{12}}{\theta^6 \sqrt{6}} B_{12} - \dots \right\}$$

$$- \sqrt{\frac{\theta}{2\pi}} \left\{ \frac{\pi^2}{\theta \sqrt{2}} B_2 - \frac{\pi^6}{\theta^3 \sqrt{3}} B_6 + \frac{\pi^{10}}{\theta^5 \sqrt{5}} B_{10} - \dots \right\}$$

(2) If  $\int_0^\infty \frac{\cos nx}{e^{2\pi\sqrt{x}} - 1} dx = \phi(n)$ , then

$$\int_0^\infty \frac{\sin nx}{e^{2\pi\sqrt{x}} - 1} dx = \phi(n) - \frac{1}{2n} + \phi\left(\frac{\pi^2}{n}\right) \sqrt{\frac{2\pi^3}{n^3}}$$

(3)  $\frac{1}{4\pi} + \frac{2 \cos \pi}{e^{2\pi} - 1} + \frac{4 \cos 4\pi}{e^{4\pi} - 1} + \frac{6 \cos 9\pi}{e^{6\pi} - 1} + \dots$

$= \phi(n) + \psi(n)$ , where,

$$\int_0^\infty e^{-2a^2\pi n} \psi(n) dn = \frac{\pi}{e^{4\pi a} - 2e^{2\pi a} \cos 2\pi a + 1}$$

(4) The part without the transcendental part of  $\phi(\pi n)$  can be found from the series

$$\frac{1}{n\sqrt{4\pi}} \left\{ \sin\left(\frac{\pi}{4} + \frac{\pi}{n}\right) + 2 \sin\left(\frac{\pi}{2} + \frac{2\pi}{n}\right) + 3 \sin\left(\frac{3\pi}{4} + \frac{3\pi}{n}\right) + \dots \right\}$$

$$- (\cos \pi n + 2 \cos 4\pi n + 3 \cos 9\pi n + \dots)$$

$$\phi(0) = \frac{1}{2}; \quad \phi\left(\frac{\pi}{2}\right) = \frac{1}{4\pi}; \quad \phi(\pi) = \frac{2-\sqrt{2}}{8}; \quad \phi(2\pi) = \frac{1}{16}$$

$$\phi\left(\frac{3\pi}{2}\right) = \frac{8-3\sqrt{5}}{16}; \quad \phi\left(\frac{\pi}{3}\right) = \frac{6+\sqrt{5}}{4} - \frac{5\sqrt{10}}{8}; \quad \phi(\infty) = 0$$

$$\phi\left(\frac{2\pi}{3}\right) = \frac{1}{3} - \sqrt{3} \left( \frac{2}{16} - \frac{1}{8\pi} \right)$$

$$5) \int_0^{\infty} e^{-2a^2 x} f(x) dx = \pi e^{-4ap} \quad 271$$

$$\text{then } f(x) = \frac{p\sqrt{2\pi}}{x\sqrt{x}} e^{-\frac{2p^2}{x}} \quad n\sqrt{\frac{x}{2}} \Psi(n\pi) =$$

$$6) \frac{p\sqrt{2x}}{\pi\sqrt{x}} \phi(n\pi) = \left\{ e^{-\frac{2\pi}{2x}} + e^{-\frac{4\pi}{x}} \left( 3 \cos \frac{3\pi}{x} + \sin \frac{3\pi}{x} \right) \right. \\ \left. + e^{-\frac{6\pi}{x}} \left( 4 \cos \frac{8\pi}{x} + 2 \sin \frac{8\pi}{x} \right) + e^{-\frac{8\pi}{x}} \left( 5 \cos \frac{15\pi}{x} + 3 \sin \frac{15\pi}{x} \right) \right. \\ \left. + e^{-\frac{10\pi}{x}} \left( 6 \cos \frac{24\pi}{x} + 4 \sin \frac{24\pi}{x} \right) + \&c \right\} + \left\{ 2e^{-\frac{8\pi}{x}} + \right. \\ \left. e^{-\frac{12\pi}{x}} \left( 5 \cos \frac{5\pi}{x} + \sin \frac{5\pi}{x} \right) + e^{-\frac{14\pi}{x}} \left( 6 \cos \frac{14\pi}{x} + 2 \sin \frac{14\pi}{x} \right) \right. \\ \left. + e^{-\frac{20\pi}{x}} \left( 7 \cos \frac{21\pi}{x} + 3 \sin \frac{21\pi}{x} \right) + \&c \right\} + \left\{ 3e^{-\frac{18\pi}{x}} \right. \\ \left. + e^{-\frac{24\pi}{x}} \left( 7 \cos \frac{7\pi}{x} + \sin \frac{7\pi}{x} \right) + \&c \right\} + \left\{ 4e^{-\frac{32\pi}{x}} + \&c \right\} + \&c$$

The pth term being,

$$pe^{-\frac{2\pi p^2}{x}} + e^{-\frac{2\pi p(p+1)}{x}} \left\{ (2p+1) \cos \frac{\pi(2p+1)}{x} + \sin \frac{\pi(2p+1)}{x} \right\} \\ + e^{-\frac{2\pi p(p+2)}{x}} \left\{ (2p+2) \cos \frac{2\pi(2p+2)}{x} + 2 \sin \frac{2\pi(2p+2)}{x} \right\} \\ + e^{-\frac{2\pi p(p+3)}{x}} \left\{ (2p+3) \cos \frac{3\pi(2p+3)}{x} + 3 \sin \frac{3\pi(2p+3)}{x} \right\} \\ + e^{-\frac{2\pi p(p+4)}{x}} \left\{ (2p+4) \cos \frac{4\pi(2p+4)}{x} + 4 \sin \frac{4\pi(2p+4)}{x} \right\} \\ + \&c \quad \&c \quad \&c \quad \&c \quad \&c$$

$$(1) \frac{\pi}{2} \frac{a^x \sin x}{\sin \pi x} + \frac{1}{2x} + \frac{\pi a}{2(1-x)} = \frac{a^x \sin x}{2-x} - \frac{a^x \sin x}{4-x} + \dots$$

$$+ \frac{e^{-2\pi a}}{2} \phi(2\pi a) + \frac{e^{-4\pi a}}{4} \phi(4\pi a) + \frac{e^{-6\pi a}}{6} \phi(6\pi a) + \dots$$

where  $\phi(x) = 1 - \frac{x}{2} + \frac{x(x+1)}{2^2} - \frac{x(x+1)(x+2)}{2^3} + \dots$

$$(2) x \left\{ \frac{1}{2} + e^{-ax-6x^2} + e^{-2ax-46x^2} + e^{-3ax-96x^2} + \dots \right\}$$

$$= \frac{1}{a} + \frac{26}{a} + \frac{46}{a} + \frac{66}{a} + \dots + \frac{B_2}{2} x^2 A_1 - \frac{B_4}{4} x^4 A_3 + \dots$$

where  $A_n = a^n - \frac{n(n-1)}{2} a^{n-2} + \frac{n(n-1)(n-2)}{6} a^{n-4} - \dots$

$$(3) \text{ when } x \text{ is small, } \frac{1}{1+x} = \frac{1}{1} - \frac{x}{1} + \frac{x^2}{1} - \frac{x^3}{1} + \frac{x^4}{1} + \dots =$$

$$x e^{\frac{1}{2}} \left\{ e^{-\frac{(1+x)^2}{2}} + e^{-\frac{(1+3x)^2}{2}} + e^{-\frac{(1+5x)^2}{2}} + \dots \right\} +$$

$$\frac{x}{2} - \frac{x^2}{12} - \frac{x^4}{360} - \frac{x^6}{5040} - \frac{x^8}{60480} - \frac{x^{10}}{1710720} \dots \text{ nearly}$$

$$(4) 2(2^2-1) \frac{B_2}{2} - 2(2^4-1) \frac{B_4}{3 \cdot 2^2} + 2(2^{10}-1) \frac{B_{10}}{5 \cdot 2^5} - \dots$$

$$= \frac{1}{2} + \frac{1}{2} + \frac{30}{2} + \frac{150}{2} + \frac{493}{2} + \dots$$

$$(5) \text{ If } m = \frac{n(n+1)}{2}, \text{ then } 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} =$$

$$c + \frac{1}{2} \log 2m + \frac{1}{12m} - \frac{1}{120m^2} + \frac{1}{630m^3} - \frac{1}{1680m^4}$$

$$+ \frac{1}{2310m^5} - \frac{171}{360360m^6} + \frac{29}{30030m^7} - \frac{2839}{1166880m^8}$$

$$+ \frac{140051}{17459472m^9} - \dots$$

$$6) x \coth x = 1 + \frac{x^2}{3} - \frac{x^4}{9} + \frac{x^6}{5} - \frac{4.5 x^8}{7} + \frac{2.3 x^{10}}{9} + \dots$$

$$\frac{6.7 x^{12}}{11} - \frac{4.5 x^{14}}{13} + \dots$$

$$7) \frac{x}{n} - \frac{x}{n+1} + \frac{x}{n+2} - \frac{x}{n+3} + \frac{x}{n+4} - \dots$$

$$= \frac{x}{n} - \frac{x}{n+1} + \frac{x}{n+2} - \frac{2(n+1)x}{1(n^2)} + \frac{1 \cdot n}{2(n+1)} x - \frac{3(n+1)x}{2(n+1)} + \frac{2(n+1)x}{3(n+3)} - \dots$$

$$8) \frac{1}{a(p+a)} + \frac{n}{(p+a+1)^2} + \frac{n^2(a+2)}{(p+a+2)^3} + \dots$$

$$\frac{n^3(a+3)^2}{(p+a+3)^4} + \frac{n^4(a+4)^3}{(p+a+4)^5} + \dots$$

$$= \int_0^1 \frac{x^{a-1} (1 - x^{\frac{p}{1-nx}})}{p} dx$$

$$9) \frac{1^{n-1}}{e^{2\pi i}} + \frac{2^{n-1}}{e^{4\pi i}} + \frac{3^{n-1}}{e^{6\pi i}} + \dots$$

$$= \frac{B_n}{2^n} + \frac{B_n}{n} \cos \frac{\pi n}{4} \left\{ \frac{1}{2^{n/2}} + \frac{2 \cos(n \tan^{-1} \frac{1}{2})}{5^{n/2}} + \dots \right\}$$

$$\frac{2 \cos(n \tan^{-1} \frac{1}{2})}{10^{n/2}} + \frac{2 \cos(n \tan^{-1} \frac{1}{5})}{13^{n/2}} + \dots \text{ where } \dots$$

2, 5, 10, 13 &c are sum of sqs. of numbers that are prime to each other



$$(10). \frac{1^{n-1}}{\cosh \frac{\pi}{2}} - \frac{3^{n-1}}{\cosh \frac{3\pi}{2}} + \frac{5^{n-1}}{\cosh \frac{5\pi}{2}} - \&c$$

$$= (2^n - 1) \frac{B_n}{n} \sin \frac{\pi n}{4} \left\{ \frac{1}{2^{\frac{n}{2}}} - \frac{2 \cos(n \tan^{-1} \frac{1}{2})}{10^{\frac{n}{2}}} + \frac{2 \cos(n \tan^{-1} \frac{2}{3})}{26^{\frac{n}{2}}} - \&c \right\}$$

$$(11) \frac{1^{n-1}}{e^{\pi} e^{-\pi}} - \frac{2^{n-1}}{e^{2\pi} e^{-2\pi}} + \frac{3^{n-1}}{e^{3\pi} e^{-3\pi}} - \&c$$

$$= -(2^n - 1) \frac{B_n}{n} \cos \frac{\pi n}{4} \left\{ \frac{1}{2^{\frac{n}{2}}} + \frac{2 \cos(n \tan^{-1} \frac{1}{2})}{10^{\frac{n}{2}}} + \frac{2 \cos(n \tan^{-1} \frac{2}{3})}{26^{\frac{n}{2}}} + \&c \right\}$$

$$(12) \frac{1^{n-1}}{\cosh \frac{\pi\sqrt{3}}{2}} - \frac{3^{n-1}}{\cosh \frac{3\pi\sqrt{3}}{2}} + \frac{5^{n-1}}{\cosh \frac{5\pi\sqrt{3}}{2}} - \&c$$

$$= (2^n - 1) \frac{B_n}{n} \sin \frac{\pi n}{6} \left\{ 1 - \frac{2 \cos \frac{\pi n}{6}}{3^{\frac{n}{2}}} + \frac{2 \cos(n \tan^{-1} \frac{\sqrt{3}}{2})}{7^{\frac{n}{2}}} - \&c \right\}$$

$$(14) \frac{1^{n-1}}{e^{\pi\sqrt{3}} + 1} - \frac{2^{n-1}}{e^{2\pi\sqrt{3}} - 1} + \frac{3^{n-1}}{e^{3\pi\sqrt{3}} + 1} - \frac{4^{n-1}}{e^{4\pi\sqrt{3}} - 1} + \&c$$

$$= - \frac{B_n}{n} \cos \frac{\pi n}{6} - \frac{B_n}{n} \left( \frac{1}{2} + \cos \frac{\pi n}{3} \right) \left\{ \frac{1}{3^{\frac{n}{2}}} + \frac{2 \cos(n \tan^{-1} \frac{1}{\sqrt{3}})}{7^{\frac{n}{2}}} + \&c \right\}$$

$$(15) \frac{16^n}{\cosh \pi\sqrt{3} + 1} - \frac{2^{6n}}{\cosh 2\pi\sqrt{3} - 1} + \frac{3^{6n}}{\cosh 3\pi\sqrt{3} - 1} - \&c$$

$$+ \frac{2\pi\sqrt{3}}{\pi} \left\{ \frac{B_{6n}}{12n} \cos 3\pi n - \left( \frac{16^{6n-1}}{e^{6\pi\sqrt{3}} + 1} - \frac{2^{6n-1}}{e^{2\pi\sqrt{3}} - 1} + \&c \right) \right\}$$

$n$  being a positive integer.

$$11) \int_0^{\infty} \frac{e^{-x} \cos x}{\sin x} x^{m-1} dx = \frac{(m-1)}{2^{m/2}} \cdot \frac{\cos \frac{\pi m}{4}}{\sin \frac{\pi m}{4}}$$

$$12) \int_0^{\infty} \frac{\sinh ax}{\sinh \pi x} \cdot \frac{dx}{x^2+x^2} = \int_0^1 \frac{x^n}{n} \cdot \frac{\sin a}{1+2x \cos a+x^2} dx$$

$$13) \frac{1}{2} \log \left[ 2\pi(n^2+x^2)^{\frac{1}{2}} \left\{ 1 + \left(\frac{x}{n+1}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+2}\right)^2 \right\} \left\{ 1 + \left(\frac{x}{n+3}\right)^2 \right\} \&c \right]$$

$$= \log \Gamma(n) + n + \tan^{-1} \frac{x}{n} - \frac{\pi}{2} \log(n^2+x^2) - \int_0^{\infty} \frac{\tan^{-1} \frac{2m x}{n^2+x^2}}{e^{2\pi x} - 1} dx$$

$$14) \text{ If } \alpha\beta = \pi, \text{ then } d \left\{ e^{-n} + e^{-ne^{\alpha}} + e^{-ne^{9\alpha}} + \&c \right\}$$

$$= d \left\{ \frac{1}{2} + \frac{n}{\Gamma} \cdot \frac{1}{e^{\alpha}} - \frac{n^2}{\Gamma^2} \cdot \frac{1}{e^{2\alpha}} + \frac{n^3}{\Gamma^3} \cdot \frac{1}{e^{3\alpha}} - \&c \right\}$$

$$+ C - \log n + 2\phi(\beta) + 2\phi(3\alpha) + 2\phi(3\beta) + \&c$$

where  $\phi(\alpha) = \sqrt{\frac{\pi}{\beta \sinh \pi \beta}} \cos \left( \beta \log \frac{\beta}{n} - \beta - \frac{\pi}{4} - \frac{\beta^2}{12\beta} - \&c \right)$

$$15) \text{ If } \alpha\beta = \frac{\pi}{2}, \text{ then } d \left\{ e^{-n e^{\alpha}} - e^{-n e^{3\alpha}} + e^{-n e^{5\alpha}} - \&c \right\}$$

$$= d \left\{ \frac{1}{2} - \frac{n}{\Gamma} \cdot \frac{1}{e^{\alpha} + e^{\alpha}} + \frac{n^2}{\Gamma^2} \cdot \frac{1}{e^{3\alpha} + e^{3\alpha}} - \&c \right\}$$

$$+ \phi(\beta) - \phi(3\alpha) + \phi(5\alpha) - \phi(7\alpha) + \&c, \text{ where}$$

$$\phi(\alpha) = \sqrt{\frac{\pi}{\beta \sinh \pi \beta}} \sin \left( \beta \log \frac{\beta}{n} - \beta - \frac{\pi}{4} - \frac{\beta^2}{12\beta} - \frac{\beta^4}{360\beta^3} - \&c \right)$$

$$16) \frac{(m)^2}{(n-1+x)(n-x)} = \left\{ 1 + \frac{x^2}{(n+1)^2} \right\} \left\{ 1 + \frac{x^2}{(n+2)^2} \right\} \left\{ 1 + \frac{x^2}{(n+3)^2} \right\} \&c$$

$$\begin{aligned}
 (1) \quad & \frac{1}{2x^2} + \frac{1}{(x+1)^2} + \frac{1}{(x+2)^2} + \frac{1}{(x+3)^2} + \dots \\
 &= \frac{1}{2\pi x^3} + \frac{\pi}{3x} - \frac{\pi^2}{\sin 2\pi x (e^{2\pi x} - 1)} + \\
 & 4x \left\{ \frac{1}{e^{4\pi} - 1} \cdot \frac{1}{(1^2 - x^2)^2} + \frac{2}{e^{8\pi} - 1} \cdot \frac{1}{(2^2 - x^2)^2} + \dots \right\} \\
 & + 8\pi x^3 \left\{ \frac{1}{(e^{2\pi} - e^{-2\pi})^2} \cdot \frac{1}{1^2 - x^4} + \frac{1}{(e^{4\pi} - e^{-4\pi})^2} \cdot \frac{1}{2^2 - x^4} + \dots \right\}
 \end{aligned}$$

$$\begin{aligned}
 (2) \quad & 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{x} = C + \frac{\pi}{3} \log x + \frac{1}{2x} - \frac{1}{4\pi x^2} \\
 & + \frac{\pi \cot \pi x}{e^{2\pi x} - 1} + \frac{2\pi \log(2 \sin \pi x)}{(e^{\pi x} - e^{-\pi x})^2} + \\
 & 2 \left( \frac{1}{e^{2\pi} - 1} \cdot \frac{1}{1^2 - x^2} + \frac{2}{e^{4\pi} - 1} \cdot \frac{1}{2^2 - x^2} + \frac{3}{e^{6\pi} - 1} \cdot \frac{1}{3^2 - x^2} + \dots \right) \\
 & - 2\pi \left\{ \frac{\log(1^2 - x^4)}{(e^{\pi} - e^{-\pi})^2} + \frac{\log(2^2 - x^4)}{(e^{2\pi} - e^{-2\pi})^2} + \frac{\log(3^2 - x^4)}{(e^{3\pi} - e^{-3\pi})^2} + \dots \right\} \\
 & - 2\pi \sum_{n=1}^{\infty} e^{-2\pi n x} \left\{ n^2 \left( \frac{\sin 2\pi x}{1^2 + n^2} + \frac{\sin 4\pi x}{2^2 + n^2} + \frac{\sin 6\pi x}{3^2 + n^2} + \dots \right) \right. \\
 & \left. - n^3 \left( \frac{\cos 2\pi x}{1^2 + n^2} + \frac{1}{2} \cdot \frac{\cos 4\pi x}{2^2 + n^2} + \frac{1}{3} \cdot \frac{\cos 6\pi x}{3^2 + n^2} + \dots \right) \right\}
 \end{aligned}$$

$$\begin{aligned}
 (3) \quad & \frac{\pi}{x^2 \sqrt{3}} \frac{\sinh \pi x \sqrt{3} \sinh \pi x + \sin \pi x \sqrt{3} \sin \pi x}{(\cosh \pi x \sqrt{3} - \cos \pi x)(\cosh \pi x - \cos \pi x \sqrt{3})} = \\
 & \frac{1}{2\pi x^4} + \coth \pi \left( \frac{1}{1+x^2+x^4} + \frac{1}{1-x^2+x^4} \right) + 2 \coth 2\pi \\
 & \times \left( \frac{1}{2^4+x^2x^2+x^4} + \frac{1}{2^4-x^2x^2+x^4} \right) + 3 \coth 3\pi \left( \frac{1}{3^4+3^2x^2+x^4} + \frac{1}{3^4-3^2x^2+x^4} \right) \\
 & + \dots
 \end{aligned}$$

$$(4) \text{ If } S_n = \frac{B_m}{2^n} + \frac{1^{n-1}}{e^{2\pi i}} + \frac{2^{n-1}}{e^{4\pi i}} + \frac{3^{n-1}}{e^{6\pi i}} + \dots$$

then if  $n-2$  be a multiple of 4,

$$\frac{(n+3)(n-4)}{24} S_{n+2} = \frac{n(n-1)(n-2)(n-3)}{12} S_4 S_{n-2} +$$

$$\frac{n(n-1)(n-2)(n-3)(n-4)(n-5)(n-6)(n-7)}{16} S_8 S_{n-6} + \dots$$

$$(5) \sqrt{1} + \sqrt{2} + \sqrt{3} + \sqrt{4} + \dots + \sqrt{x} = C + \frac{2}{3} x\sqrt{x} + \frac{1}{2} \sqrt{x} + \frac{1}{6} \left\{ \frac{1}{(\sqrt{x} + \sqrt{x+1})^3} + \frac{1}{(\sqrt{x+1} + \sqrt{x+2})^3} + \dots \right\}$$

$$(6) 1\sqrt{1} + 2\sqrt{2} + 3\sqrt{3} + \dots + x\sqrt{x} = C + \frac{2}{5} x^2\sqrt{x} + \frac{x}{2}\sqrt{x} + \frac{1}{8}\sqrt{x} + \frac{1}{40} \left\{ \frac{1}{(\sqrt{x} + \sqrt{x+1})^5} + \frac{1}{(\sqrt{x+1} + \sqrt{x+2})^5} + \dots \right\}$$

$$(7) (1^2\sqrt{1} + 2^2\sqrt{2} + 3^2\sqrt{3} + \dots + x^2\sqrt{x}) + \frac{1}{16} (\sqrt{1} + \sqrt{2} + \dots + \sqrt{x}) = C + \frac{2}{7} x^2\sqrt{x} + \frac{x^2}{2}\sqrt{x} + \frac{x}{4}\sqrt{x} + \frac{1}{32}\sqrt{x} + \frac{1}{224} \left\{ \frac{1}{(\sqrt{x} + \sqrt{x+1})^7} + \frac{1}{(\sqrt{x+1} + \sqrt{x+2})^7} + \dots \right\}$$

$$(8) \sum \frac{1}{x} - \sum \frac{1}{x/3} + \frac{1}{x} - \log 3 =$$

$$\frac{2}{3} \cdot \frac{1}{x^2} + \frac{2^2-2}{6} + \frac{4^2-4}{3x^2} + \frac{5^2-5}{6} + \frac{7^2-7}{5x^2} + \dots$$

$$(9) \int_0^{\infty} \cos mx \log(1+x^2) dx = \dots$$

278

$$(1). \frac{x^n}{\Gamma} \left\{ 1 + \frac{x^2}{\Gamma} \cdot \frac{1}{(1+n)} + \frac{x^4}{\Gamma} \cdot \frac{1}{(1+n)(1+n)} + \dots \right\}$$

$$- \frac{x^{-n}}{\Gamma} \left\{ 1 + \frac{x^2}{\Gamma} \cdot \frac{1}{(1-n)} + \frac{x^4}{\Gamma} \cdot \frac{1}{(1-n)(1-n)} + \dots \right\}$$

$$= - \frac{e^{-2x}}{\sqrt{\pi x}} \sin \pi n \left\{ 1 + \frac{n^2 - \frac{1}{2}}{4x} + \frac{(n^2 - \frac{1}{2})(n^2 - \frac{3}{2})}{6,8 x^2} + \dots \right\}$$

$$= - \frac{\sin \pi n}{\Gamma} \int_0^{\infty} x^{n-1} e^{-x(2+\frac{1}{2})} dx.$$

$$(2). \int_0^{\infty} x^{2n} e^{-x^2} \frac{a^x}{x^2} dx = \frac{\sqrt{\pi}}{2} e^{-2a} a^n \left\{ 1 + \frac{n(n+1)}{4a} + \frac{(n-1)n(n+1)(n+2)}{6,8 a^2} + \frac{(n-2)(n-1)n(n+1)(n+2)(n+3)}{6,8 \cdot 12 a^3} + \dots \right\}$$

N.B. Integrate partially and add.

$$(3) \log \left( 1 + \frac{x^2}{\Gamma} \right) - 3 \log \left( 1 + \frac{x^2}{3\Gamma} \right) + 5 \log \left( 1 + \frac{x^2}{5\Gamma} \right) - \dots$$

$$+ 2x \tan^{-1} e^{-\frac{\pi x}{2}} =$$

$$\frac{4}{\pi} \left( \frac{1 - e^{-\frac{\pi x}{2}}}{1^2} - \frac{1 - e^{-\frac{3\pi x}{2}}}{3^2} + \frac{1 - e^{-\frac{5\pi x}{2}}}{5^2} - \dots \right)$$

$$(4). \log \left\{ 1 + \left( \frac{2}{\pi} \log \sqrt{2+\sqrt{3}} \right)^2 \right\} - 3 \log \left\{ 1 + \left( \frac{2}{3\pi} \log \sqrt{2+\sqrt{3}} \right)^2 \right\} +$$

$$\log \left\{ 1 + \left( \frac{2}{5\pi} \log \sqrt{2+\sqrt{3}} \right)^2 \right\} - \dots = \frac{4}{3\pi} \left( \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots \right)$$

$$5) \frac{1^m}{1-x^2} + \frac{2^m}{2-x^2} + \frac{3^m}{3-x^2} + \dots$$

$$= \frac{\pi}{2} x^{n+1} (\tan \frac{\pi n}{2} - \cot \pi x) + 2 \sin \frac{\pi n}{2} \int_0^\infty \frac{z^n}{e^{2\pi z} - 1} \cdot \frac{dz}{2+x^2}$$

$$6) \left( \frac{1^m}{1+x^2} + \frac{2^m}{2+x^2} + \frac{3^m}{3+x^2} + \dots \right) - \frac{\pi}{2} x^{n+1} \sec \frac{\pi n}{2}$$

$$= \frac{\pi x^{n+1} \cos \frac{\pi n}{2}}{e^{2\pi x} - 1} + 2 \sin \frac{\pi n}{2} \int_0^\infty \frac{z^n}{e^{2\pi z} - 1} \cdot \frac{dz}{2-x^2}$$

(7) If  $\int_0^\infty e^{-p^2 x} \phi(x) dx = \frac{e^{-2ap}}{p^{n+1}}$ , then

$$\phi(x) = \frac{x^n}{\sqrt{\pi x}} e^{-\frac{a^2}{x}} \int_0^\infty e^{-az - \frac{xz^2}{4}} \frac{z^{n+1}}{z^2} dz$$

$$= \frac{x^n}{a^n \sqrt{\pi x}} e^{-\frac{a^2}{x}} \left\{ 1 - \frac{n(n+1)}{4a^2} x + \frac{n(n+1)(n+3)(n+2)}{4 \cdot 8 \cdot a^4} x^2 - \frac{n(n+1)(n+2)(n+3)(n+4)(n+5)}{4 \cdot 8 \cdot 12 \cdot a^6} x^3 + \dots \right\}$$

(8) If  $\frac{d}{d\theta} \theta u = u + \frac{1}{2} \frac{u^7}{7} + \frac{1 \cdot 3}{2 \cdot 4} \frac{u^{13}}{13} + \dots$

then  $\frac{1}{9} \cdot \frac{u^2}{u^2} = \frac{1}{3} \sin \theta - \frac{2}{\pi \sqrt{3}} + 8 \left( \frac{\cos 2\theta}{e^{\pi \sqrt{3}} + 1} - \frac{2 \cos 4\theta}{e^{2\pi \sqrt{3}} - 1} + \frac{3 \cos 6\theta}{e^{3\pi \sqrt{3}} + 1} - \dots \right)$  where  $u = \frac{\sqrt{\pi}}{[-\frac{1}{3}]^{-\frac{1}{3}}}$

(9)  $\frac{\beta_1}{2} - \frac{\beta_2}{8} + \dots, \frac{\beta_2}{2} \cos + \frac{\beta_4}{4} \cos \dots$

(10)  $\int_0^x \left( \frac{x}{x} \right)^x dx = \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots$

(11) The difference between  $\frac{\sqrt{\beta-m}}{\alpha+\beta-m}$  and

$$\frac{\sqrt{\beta}}{\alpha+\beta} + \frac{\alpha}{\Gamma} \cdot m \cdot \frac{\sqrt{\beta+m}}{\alpha+\beta+m+1} + \frac{\alpha(\alpha+1) \cdot m(m+2\alpha+1)}{\Gamma^2}$$

$$\times \frac{\sqrt{\beta+2m}}{\alpha+\beta+2m+2} + \frac{\alpha(\alpha+1)(\alpha+2) m(m+3\alpha+1)(m+2\alpha+2)}{\Gamma^3}$$

$$\times \frac{\sqrt{\beta+3m}}{\alpha+\beta+3m+3} + \dots$$

(12)  $e^{-\frac{x}{2\alpha}} (1-e^{-a})^{\frac{1}{2}} (1-e^{-a-x}) (1-e^{-a-2x}) (1-e^{-a-3x}) \dots$

$$= \frac{\left(\frac{a}{x}\right)^{\frac{a}{x}} \sqrt{\frac{2\pi a}{x}}}{e^{\frac{a}{x}} \Gamma^{\frac{a}{x}}} e^{-\frac{1}{x} \left( \frac{e^{-a}}{\Gamma} + \frac{e^{-2a}}{\Gamma^2} + \dots \right) - \theta}$$

where  $\theta = \sum_{n=1}^{\infty} \frac{B_{2n}}{2n} \cdot \frac{B_{2n}}{2n} \cdot \frac{a}{\Gamma} \cdot \frac{x^{2n-1}}{\Gamma^{2n-1}}$

$$\frac{B_{2m}}{\Gamma^{2m}} x^{2m-1} \left\{ \frac{B_{2m}}{2m} \cdot \frac{a}{\Gamma} - \frac{B_{2m+2}}{2m+2} \cdot \frac{a^2}{\Gamma^2} + \dots \right\}$$

(13) The property of the function

$$\frac{\log 1}{1^2+x^2} + \frac{\log 2}{2^2+x^2} + \frac{\log 3}{3^2+x^2} + \dots \text{ and}$$

the integral  $\int_0^{\infty} \frac{x}{e^{2\pi x} - 1} \cdot \frac{dx}{x+x}$

(1) If  $\frac{\theta u}{\sqrt{v}} = v + \frac{1}{2} \cdot \frac{v^5}{5} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{v^9}{9} + \dots$  where  $\theta$  is the constant obtained by putting  $v=1$  and  $\theta = \frac{\pi}{2}$ , then

$$(1) \frac{u^2}{2v^2} = \frac{1}{\sin 2\theta} - \frac{1}{\pi} - 8 \left( \frac{\cos 2\theta}{e^{4\pi}} + \frac{2\cos 4\theta}{e^{8\pi}} + \dots \right)$$

$$(2) \frac{u}{\sqrt{v}} \left( \frac{1}{v} - \frac{1}{2} \cdot \frac{v^3}{3} - \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{v^7}{7} - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \cdot \frac{v^{11}}{11} - \dots \right) \\ = \cot \theta + \frac{\theta}{\pi} + 4 \left( \frac{\sin 2\theta}{e^{2\pi}} + \frac{\sin 4\theta}{e^{4\pi}} + \frac{\sin 6\theta}{e^{6\pi}} + \dots \right)$$

$$(3) \log \frac{v\sqrt{v}}{u} + \frac{1}{3} \cdot \frac{v^4}{4} + \frac{1 \cdot 5}{3 \cdot 7} \cdot \frac{v^8}{8} + \frac{1 \cdot 5 \cdot 9}{3 \cdot 7 \cdot 11} \cdot \frac{v^{12}}{12} + \dots \\ = \log \sin \theta + \frac{\theta^2}{2\pi} - 2 \left\{ \frac{\cos 2\theta}{1(e^{2\pi}-1)} + \frac{\cos 6\theta}{2(e^{4\pi}-1)} + \dots \right\}$$

$$(4) \frac{1}{2} \tan^{-1} v^2 = \frac{\sin \theta}{\cosh \frac{\pi}{2}} + \frac{\sin 3\theta}{3 \cosh \frac{3\pi}{2}} + \frac{\sin 5\theta}{5 \cosh \frac{5\pi}{2}} + \dots$$

$$(5) \frac{1}{2} \cos^{-1} v^2 = \frac{\cos \theta}{\cosh \frac{\pi}{2}} - \frac{\cos 3\theta}{3 \cosh \frac{3\pi}{2}} + \frac{\cos 5\theta}{5 \cosh \frac{5\pi}{2}} - \dots$$

$$(6) \frac{\sqrt{v}}{1 \cdot u} \left\{ \frac{v^3}{3} + \frac{2}{3} \cdot \frac{v^7}{7} + \frac{2 \cdot 4}{3 \cdot 5} \cdot \frac{v^{11}}{11} + \dots \right\} \\ = \frac{\pi \theta}{8} - \frac{\sin \theta}{2 \cosh \frac{\pi}{2}} + \frac{\sin 3\theta}{3 \cosh \frac{3\pi}{2}} - \frac{\sin 5\theta}{5 \cosh \frac{5\pi}{2}} + \dots$$

If  $\frac{\theta u}{v} = v - \frac{1}{2} \cdot \frac{v^5}{5} + \frac{1 \cdot 3}{2 \cdot 4} \cdot \frac{v^9}{9} - \dots$ , then



$$2.82$$

$$(7) 2 \tan^{-1} v = \theta + \frac{\sin 2\theta}{\cosh \pi} + \frac{\sin 4\theta}{2 \cosh 2\pi} + \dots$$

$$(8) \frac{\pi}{8} - \frac{1}{2} \tan^{-1} v = \frac{\cos \theta}{\cosh \frac{\pi}{2}} - \frac{\cos 3\theta}{3 \cosh \frac{3\pi}{2}} + \dots$$

$$(9) \frac{1}{2} \log \frac{1+v}{1-v} = \log \tan \left( \frac{\pi}{4} + \frac{\theta}{2} \right) + \frac{1}{4} \left\{ \frac{\sin \theta}{e^{\pi} - 1} - \frac{\sin 3\theta}{3(e^{3\pi} - 1)} + \dots \right\}$$

$$(10) \log \left( 1 - \frac{x^2}{1^2} \right) - 3 \log \left( 1 - \frac{x^2}{3^2} \right) + 5 \log \left( 1 - \frac{x^2}{5^2} \right) - \dots$$

$$= \frac{4}{\pi} \left\{ \frac{1 - \cos \frac{\pi x}{2}}{1^2} - \frac{1 - \cos \frac{3\pi x}{2}}{3^2} + \dots \right\} +$$

$$x \log \tan \frac{\pi - \pi x}{4}$$

$$= \frac{4}{\pi} \left\{ \frac{1 - \tan \left( \frac{\pi - \pi x}{4} \right)}{1^2} - \frac{1 - \tan \left( \frac{3(\pi - \pi x)}{4} \right)}{3^2} + \dots \right\}$$

$$+ \log \tan \frac{\pi - \pi x}{4}$$

$$(11) \text{ If } \frac{\pi \alpha}{2} = \log \tan \left( \frac{\pi}{4} + \frac{\pi \beta}{4} \right), \text{ then}$$

$$\log \left( 1 + \frac{\alpha^2}{1^2} \right) - 3 \log \left( 1 + \frac{\alpha^2}{3^2} \right) + 5 \log \left( 1 + \frac{\alpha^2}{5^2} \right) - \dots$$

$$= \frac{\pi \alpha \beta}{2} + \log \left( 1 - \frac{\alpha^2}{1^2} \right) - 3 \log \left( 1 - \frac{\alpha^2}{3^2} \right) + 5 \log \left( 1 - \frac{\alpha^2}{5^2} \right) - \dots$$

— &c

283

$$(1) \text{ If } \phi(m, n) = \left\{ 1 + \left( \frac{m+n}{1+m} \right)^3 \right\} \left\{ 1 + \left( \frac{m+n}{2+m} \right)^3 \right\} \&c$$

then  $\phi(m, n) \phi(n, m) =$

$$\frac{(m)^3 (n)^3}{|2m+n| |2n+m|} \cdot \frac{\cosh \pi(m+n)\sqrt{3} - \cos \pi(m-n)}{2\pi^2 (m^2 + mn + n^2)}$$

$$(2) \left\{ 1 + \left( \frac{n}{1} \right)^3 \right\} \left\{ 1 + \left( \frac{n}{2} \right)^3 \right\} \left\{ 1 + \left( \frac{n}{3} \right)^3 \right\} \&c$$

$$\times \left\{ 1 + 3 \cdot \left( \frac{n}{n+2} \right)^4 \right\} \left\{ 1 + 3 \left( \frac{n}{n+4} \right)^4 \right\} \left\{ 1 + 3 \left( \frac{n}{n+6} \right)^4 \right\} \&c$$

$$= \frac{\left| \frac{n}{2} - 1 \right|}{\left| \frac{n}{2} \right|} \cdot \frac{\cosh \pi n \sqrt{3} - \cos \pi n}{2^{n+2} \pi n \sqrt{\pi}}$$

$$(3) \frac{3}{2} \log 2 \pi n + \log \left( 1 + \frac{n^2}{1} \right) \left( 1 + \frac{n^2}{2^2} \right) \left( 1 + \frac{n^2}{3^2} \right) \&c$$

$$- \log \left( e^{\pi n \sqrt{3}} + e^{-\pi n \sqrt{3}} - 2 \cos \pi n \right)$$

$$= - \frac{\pi n}{\sqrt{3}} + \frac{B_2}{2} \cdot \frac{1}{n^2} - \frac{B_4}{10} \cdot \frac{1}{3n^4} + \frac{B_6}{16} \cdot \frac{1}{5n^6} \&c$$

$$(4) \frac{B_2}{1 \cdot 2 \cdot 2n} + \frac{B_4}{3! \cdot 4 \cdot 2^2 n^3} - \frac{B_6}{5! \cdot 6 \cdot 2^3 n^5} - \frac{B_8}{7! \cdot 8 \cdot 2^4 n^7} + \&c$$

$$= \log \frac{e^{\pi n \sqrt{3}}}{\pi^n \sqrt{2\pi n}} + \frac{n}{2} \left( \frac{\pi}{2} - \log 2 \right) - \frac{1}{2} \log 2$$

$$- \frac{1}{2} \log \left\{ 1 + \left( \frac{n}{n+1} \right)^2 \right\} \left\{ 1 + \left( \frac{n}{n+2} \right)^2 \right\} \left\{ 1 + \left( \frac{n}{n+3} \right)^2 \right\} \&c$$

(1) The difference between the two series ( $d\beta = \frac{\pi^2}{8}$ )

$$\alpha^2 \left\{ \frac{\operatorname{sech} \frac{\pi}{2}}{\cosh \alpha + \cos \alpha} - \frac{3^3 \operatorname{sech} \frac{3\pi}{2}}{\cosh 3\alpha + \cos 3\alpha} + \dots \right\} \text{ and}$$

$$\beta^2 \left\{ \frac{\operatorname{sech} \frac{\pi}{2}}{\cosh \beta + \cos \beta} - \frac{3^3 \operatorname{sech} \frac{3\pi}{2}}{\cosh 3\beta + \cos 3\beta} \right\} \text{ is } 0?$$

$$(2) \int_0^{\infty} \frac{\sin 2\pi x}{x(\cosh \pi x + \cos \pi x)} dx = \frac{\pi}{4} - 2 \left( \frac{e^{-\pi} \cos \pi}{\cosh \frac{\pi}{2}} \right. \\ \left. - \frac{e^{-3\pi} \cos 3\pi}{3 \cosh \frac{3\pi}{2}} + \frac{e^{-5\pi} \cos 5\pi}{5 \cosh \frac{5\pi}{2}} - \dots \right)$$

$$(3) \text{ If } d\beta = \frac{\pi^2}{4}, \text{ then, } \frac{1}{\cosh d + \cos d} + \\ - \frac{1}{3(\cosh 3d + \cos 3d)} + \frac{1}{5(\cosh 5d + \cos 5d)} - \dots \\ = \frac{\pi}{8} - \frac{2 \cos \beta \cosh \beta}{\cosh \frac{\pi}{2} (\cosh 2\beta + \cos 2\beta)} + \\ \frac{2 \cos 3\beta \cosh 3\beta}{3 \cosh \frac{3\pi}{2} (\cosh 6\beta + \cos 6\beta)} + \dots$$

$$(4) \text{ If } d\beta = \frac{\pi^2}{2}, \text{ then } \frac{\pi}{8} - \frac{\pi^3}{32d^2} + \\ \frac{\cos d}{\cosh d - \cos d} - \frac{\cos 3d}{3(\cosh 3d - \cos 3d)} + \dots = \\ \frac{\sin \beta \sinh \beta}{\cosh 2\beta + \cos 2\beta} \cdot \frac{\cot \frac{\pi}{2}}{1} + \frac{\sin 2\beta \sinh 2\beta}{\cosh 4\beta + \cos 4\beta} \cdot \frac{\cot \frac{\pi}{2}}{2}$$

(1) The difference between the series

$$\frac{\theta}{8\pi} + \frac{\sin \theta}{1(e^{2\pi}-1)} + \frac{\sin 4\theta}{2(e^{4\pi}-1)} + \frac{\sin 9\theta}{3(e^{6\pi}-1)} + \dots$$

and  $\frac{1}{4} \left\{ \frac{B_2}{14} \theta - \frac{B_6}{315} \theta^3 + \frac{B_{10}}{515} \theta^5 - \dots \right\}$

(2)  $\frac{\pi}{2n} \cdot \frac{\sec \frac{\pi m}{2n}}{e^{\frac{\pi m}{2n}} - 1} + \frac{1}{m+n} - \frac{1}{m+3n} + \frac{1}{m+5n} - \dots$

$$= \frac{1}{2} + \frac{\operatorname{sech} \frac{\pi m}{2n}}{1+(2n)^2} + \frac{\operatorname{sech} 2\pi m}{1+(4n)^2} + \dots$$

$$- 2m \left\{ \frac{1}{m^2-n^2} \cdot \frac{1}{e^{\frac{\pi m}{2n}} - 1} - \frac{1}{m^2-(3n)^2} \cdot \frac{1}{e^{\frac{3\pi m}{2n}} - 1} + \dots \right\}$$

(3) If  $\phi = \frac{x}{1+x} + \frac{x^5}{1+x} + \frac{x^{10}}{1+x} + \frac{x^{15}}{1+x} + \dots$  and

$$f = \frac{\sqrt{x}}{1+x} + \frac{x}{1+x} + \frac{x^2}{1+x} + \frac{x^3}{1+x} + \dots, \text{ then}$$

$$f^5 = \phi \cdot \frac{1-2\phi+4\phi^2-3\phi^3+\phi^4}{1+3\phi+4\phi^2+2\phi^3+\phi^4}$$

(4)  $1 - \frac{ax}{1+x} + \frac{a^2}{1+x} - \frac{a^3x}{1+x} + \frac{a^4}{1+x} - \frac{a^5x}{1+x} + \dots$  } *Convergent only.*

$$= \frac{a}{x} + \frac{a^4}{x} + \frac{a^8}{x} + \frac{a^{12}}{x} + \dots \text{ nearly.}$$

(5)  $\frac{\pi}{2} \int_0^{\infty} \frac{dx}{e^{x^n} + e^{-x^n}} =$

$$\frac{\pi}{2} \left[ \frac{1}{n} - 1 \cos \frac{\pi}{2n} \right] \int_0^{\infty} \frac{x^{n-2}}{e^{x^n} + e^{-x^n}} dx$$

286

$$(1) \frac{x}{4n+2} + \frac{x^2}{4n+6} + \frac{x^2}{4n+10} + \dots$$

$$+ \frac{2n}{x} + \frac{n-1}{1} - \frac{n+1}{x} + \frac{n-2}{k} - \frac{n+2}{x} + \dots$$

$= 1$  nearly.

$$(2) 1 - \frac{ax}{1+a} + \frac{a^2x}{1+a^2} - \frac{a^4x}{1+a^4} + \frac{a^6x}{1+a^6} - \frac{a^8x}{1+a^8} + \dots$$

$$\frac{a^9x}{1+a^6} - \dots = \frac{1}{x} + \frac{a}{x} + \frac{a^3}{x} + \frac{a^5}{x} + \dots \text{ nearly.}$$

$$(3) \frac{1-a^2}{1-a} \cdot \frac{1-a^4}{1-a^4} \cdot \frac{1-a^8}{1-a^8} \cdot \frac{1-a^{16}}{1-a^{16}} \dots$$

$$= \frac{1}{1-a} \cdot \frac{a}{1+a} - \frac{a^3}{1+a^2} - \frac{a^5}{1+a^3} - \frac{a^7}{1+a^4} - \dots$$

$$(4) \frac{1-a^3}{1-a} \cdot \frac{1-a^7}{1-a^5} \cdot \frac{1-a^{11}}{1-a^9} + \dots =$$

$$\frac{1}{1-a} \cdot \frac{a}{1+a^2} - \frac{a^3}{1+a^4} - \frac{a^5}{1+a^6} - \dots$$

$$(5) \frac{1+a^2}{1+a} \cdot \frac{1+a^4}{1+a^3} \cdot \frac{1+a^6}{1+a^5} \dots =$$

$$\frac{1}{1+a} \cdot \frac{a}{1+a} \cdot \frac{a^2+a}{1+a} \cdot \frac{a^3}{1+a} \cdot \frac{a^4+a^2}{1+a} \cdot \frac{a^5}{1+a} \dots$$

$$(6) \frac{(1-a)(1-a^3)(1-a^5)(1-a^7) \dots}{(1-a^2)(1-a^4)(1-a^6)(1-a^8) \dots} =$$

$$\frac{1}{1+a} \cdot \frac{a+a^2}{1+a} \cdot \frac{a^3}{1+a} \cdot \frac{a^3+a^6}{1+a} \cdot \frac{a^8}{1+a} \dots$$

$$(1) \text{ If } \phi(\alpha, \beta) = \frac{\pi}{e^{4\pi\alpha} - 2e^{2\pi\alpha} \cos 2\pi\beta + 1} + \dots \quad 287$$

$$\alpha \left\{ \frac{1}{2(\alpha^2 + \beta^2)} + \frac{1}{\alpha^2 + (1+\beta)^2} + \frac{1}{\alpha^2 + (2+\beta)^2} + \dots \right\}$$

$$- 4\alpha\beta \left\{ \frac{1}{e^{4\pi} - 1} \cdot \frac{1}{\alpha^2 + (1+\beta)^2} - \frac{1}{\alpha^2 + (1-\beta)^2} + \right.$$

$$\left. \frac{2}{e^{8\pi} - 1} \cdot \frac{1}{\alpha^2 + (2+\beta)^2} - \frac{1}{\alpha^2 + (2-\beta)^2} + \dots \right\}$$

$$\text{then } \phi(\alpha, \beta) + \phi(\beta, \alpha) = \frac{\pi}{2} + \frac{\alpha\beta}{\pi(\alpha^2 + \beta^2)^2} +$$

$$\frac{\pi}{2} \cdot \frac{\cosh 2\pi(\alpha - \beta) - \cos 2\pi(\alpha - \beta)}{(\cosh 2\pi\alpha - \cos 2\pi\beta)(\cosh 2\pi\beta - \cos 2\pi\alpha)}$$

$$(2) \text{ If } \phi(\alpha, \beta) = \frac{\pi/2}{e^{2\pi\alpha} + 2e^{\pi\alpha} \cos \pi\beta + 1} +$$

$$\alpha \left\{ \frac{1}{\alpha^2 + (1+\beta)^2} + \frac{1}{\alpha^2 + (3+\beta)^2} + \frac{1}{\alpha^2 + (5+\beta)^2} + \dots \right\}$$

$$+ 4\alpha\beta \left\{ \frac{1}{e^{\pi} + 1} \cdot \frac{1}{\alpha^2 + (1+\beta)^2} - \frac{1}{\alpha^2 + (1-\beta)^2} + \dots \right\}$$

$$\text{then } \phi(\alpha, \beta) + \phi(\beta, \alpha) = \frac{\pi}{4} +$$

$$\frac{\pi}{4} \cdot \frac{\cosh \pi(\alpha - \beta) - \cos \pi(\alpha - \beta)}{(\cosh \pi\alpha + \cos \pi\beta)(\cosh \pi\beta + \cos \pi\alpha)}$$

286  
 $\int \frac{y}{x} = \frac{\sqrt{1+x^2}-1}{x}$  and  $m = \frac{n}{\sqrt{1+x^2}}$ , then

$$(1) \frac{x}{1+n} + \frac{(x)^2}{3+n} + \frac{(3x)^2}{5+n} + \frac{(3x)^2}{7+n} + \dots$$

$$= 2 \left( \frac{y}{m+1} - \frac{y^3}{m+3} + \frac{y^5}{m+5} - \dots \right)$$

$$(2) \frac{x}{2+n} + \frac{1.2x^2}{4+n} + \frac{2.3x^2}{6+n} + \frac{3.4x^2}{8+n} + \dots$$

$$= y - m(y + \frac{1}{y}) \left( \frac{y^2}{m+2} - \frac{y^4}{m+4} + \frac{y^6}{m+6} - \dots \right)$$

$$(3) \frac{1}{n} + \frac{1.p}{n} + \frac{2(p+1)}{n} + \frac{3(p+2)}{n} + \frac{4(p+3)}{n} + \dots$$

$$= 2^p \left\{ \frac{1}{n+p} - \frac{p}{4} \cdot \frac{1}{n+p+2} + \frac{p(p+1)}{16} \cdot \frac{1}{n+p+4} - \dots \right\}$$

$$(4) \frac{x}{p+n} + \frac{1.p x^2}{p+2+n} + \frac{2(p+1)x^2}{p+4+n} + \frac{3(p+2)x^2}{p+6+n} + \dots$$

$$= \left(1 + \frac{1}{2x}\right)^{\frac{p-1}{2}} (2y)^p \left\{ \frac{1}{m+p} - \frac{p}{4} \cdot \frac{y^2}{m+p+2} + \frac{p(p+1)}{16} \cdot \frac{y^4}{m+p+4} - \dots \right\}$$

$$(5) \frac{1}{2x^2} + \frac{1}{(x+1)^2} + \frac{1}{(x+2)^2} + \frac{1}{(x+3)^2} + \dots$$

$$= \frac{1}{2} + \frac{1}{2x^2} \cdot \frac{1}{3x} + \frac{3}{5x} + \frac{18}{7x} + \frac{60}{9x} + \dots$$

$$3 = 2^2(2^2-1)/4; \quad 18 = 3^2(3^2-1)/4; \quad 60 = 4^2(4^2-1)/4 \dots$$

$$(6). \frac{1}{2x^2} + \frac{1}{(x+1)^2} + \frac{1}{(x+2)^2} + \frac{1}{(x+3)^2} + \dots$$

$$= \frac{1}{2x^2} + \frac{1}{4x^2} + \frac{1}{x} + \frac{1}{3x} + \frac{2}{x} + \frac{6}{5x} + \frac{9}{x} + \frac{18}{7x} + \dots$$

$$(7) 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} = \frac{1}{2m} - \frac{1}{2\pi n^2} +$$

$$\frac{\pi \cot \pi n}{e^{2\pi n} - 1} + \frac{n^2}{1(1^2+n^2)} + \frac{n^2}{2(2^2+n^2)} + \frac{n^2}{3(3^2+n^2)} + \dots$$

$$+ \frac{4n^2}{1^4-n^4} \cdot \frac{1}{e^{2\pi} - 1} + \frac{8n^2}{2^4-n^4} \cdot \frac{1}{e^{4\pi} - 1} + \frac{12n^2}{3^4-n^4} \cdot \frac{1}{e^{6\pi} - 1} + \dots$$

$$(8) 1 + \frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{n-1} = \frac{\pi}{2} \cdot \frac{\tan \frac{\pi n}{2}}{e^{\pi n} + 1}$$

$$+ \frac{n^2}{1(1^2+n^2)} + \frac{n^2}{3(3^2+n^2)} + \frac{n^2}{5(5^2+n^2)} + \dots$$

$$- \left( \frac{4n^2}{1^4-n^4} \cdot \frac{1}{e^{\pi} + 1} + \frac{12n^2}{3^4-n^4} \cdot \frac{1}{e^{3\pi} + 1} + \dots \right)$$

$$(9) \frac{1}{n+1} - \frac{1}{n+3} + \frac{1}{n+5} - \frac{1}{n+7} + \dots$$

$$= \frac{1}{2m} - \frac{\pi}{2} \cdot \frac{\sec \frac{\pi n}{2}}{e^{\pi n} - 1} +$$

$$2m \left\{ \frac{1}{1^2-n^2} \cdot \frac{1}{e^{\pi} - 1} - \frac{1}{3^2-n^2} \cdot \frac{1}{e^{3\pi} - 1} + \dots \right\}$$

$$+ 2m \left\{ \frac{1}{2^2+n^2} \cdot \frac{1}{e^{\pi} + e^{\pi}} + \frac{1}{4^2+n^2} \cdot \frac{1}{e^{2\pi} + e^{2\pi}} + \dots \right\}$$



$$(1+e^{-\pi n})(1+e^{-3\pi n})(1+e^{-5\pi n}) \&c$$

$$= \frac{4/2}{24 \sqrt{G_n} e^{\pi n}}$$

$$(1-e^{-\pi n})(1-e^{-3\pi n})(1-e^{-5\pi n}) \&c$$

$$= \frac{4/2}{24 \sqrt{g_n} e^{\pi n}} \quad \text{then}$$

$$g_n G_n = 64 g_{2n} \quad \text{and} \quad h_1 = 4 \sqrt[3]{\frac{G}{g}} + \sqrt[3]{\frac{g}{G}}$$

$$\sqrt{1}. \quad G = 1.$$

$$\sqrt{3}. \quad G = \frac{1}{4}$$

$$\sqrt{5}. \quad G = (\sqrt{5}-2)^2$$

$$\sqrt{7}. \quad G = \frac{1}{64}$$

$$\sqrt{9}. \quad G = (2-\sqrt{3})^4$$

$$\sqrt{11}. \quad G^3 - G^2 + G = \frac{1}{2}$$

$$\sqrt{13}. \quad G = \left(\frac{\sqrt{13}-3}{2}\right)^6$$

$$\sqrt{15}. \quad G = \frac{1}{64} \left(\frac{\sqrt{5+1}}{2}\right)^8$$

$$\sqrt{17}. \quad G = \left(\frac{5+\sqrt{17}}{8} - \sqrt{\frac{12-3}{9}}\right)^{24}$$

$$\sqrt{19}. \quad G^3 + G^2 = \frac{1}{2}$$

$$\sqrt{21}. \quad G = (2-3\sqrt{3})^2 \left(\frac{5 \pm \sqrt{41}}{2}\right)^3$$

$$\sqrt{23}. \quad G^3 + G^2 = 1$$

$$\sqrt{25}. \quad G = (\sqrt{5}-2)^8$$

$$\sqrt{27}. \quad G = \frac{1}{4} (\sqrt[3]{2}-1)^8$$

$$\text{or } \{G^3 + G^2 \sqrt{3} = \frac{1}{2}\}$$

$$\sqrt{31} \quad \{ G_1^3 + G_2 = 1 \}$$

$$\sqrt{33} \quad G_2 = (2 - \sqrt{3})^6 (10 \pm 3\sqrt{11})^2$$

$$\sqrt{37} \quad G_1 = (\sqrt{37} - 6)^6$$

$$\sqrt{39} \quad G_1 = \frac{1}{64} \left( \frac{\sqrt{13} - 3}{2} \right)^4 \left( \sqrt{\frac{5 + \sqrt{13}}{8}} \pm \sqrt{\frac{\sqrt{13} - 3}{8}} \right)^{24}$$

$$\sqrt{43} \quad \{ G_1^3 + G_2 = \frac{1}{2} \}$$

$$\sqrt{45} \quad G_1 = (\sqrt{5} - 2)^6 (4 \pm \sqrt{15})^4$$

$$\sqrt{49} \quad G_1 = \left( \frac{\sqrt{4 + \sqrt{7}} - \sqrt{7}}{2} \right)^{24}$$

$$\sqrt{55} \quad G_1 = \frac{1}{64} (\sqrt{5} - 2)^4 \left( \sqrt{\frac{7 + \sqrt{5}}{8}} \pm \sqrt{\frac{\sqrt{5} - 1}{8}} \right)^{24}$$

$$\sqrt{57} \quad G_1 = \left( \frac{3\sqrt{19} - 13}{\sqrt{2}} \right)^4 (2 \pm \sqrt{3})^6$$

$$\sqrt{63} \quad G_1 = \frac{1}{64} \left( \frac{5 - \sqrt{21}}{2} \right)^4 \left( \sqrt{\frac{5 + \sqrt{21}}{8}} - \sqrt{\frac{\sqrt{21} - 3}{8}} \right)^{24}$$

$$\sqrt{65} \quad G_1 = \left( \frac{\sqrt{13} \pm 3}{2} \right)^6 (\sqrt{5} \pm 2)^2 \left( \sqrt{\frac{9 + \sqrt{65}}{8}} - \sqrt{\frac{1 + \sqrt{65}}{8}} \right)^{12}$$

$$\sqrt{67} \quad \{ G_1^3 + G_2 + G_3 = \frac{1}{2} \}$$

$$\sqrt{69} \quad G_1 = \left( \frac{5 \pm \sqrt{23}}{\sqrt{2}} \right)^2 \left( \frac{3\sqrt{3} \pm \sqrt{23}}{2} \right)^3 \left( \sqrt{\frac{6 + 3\sqrt{3}}{4}} - \sqrt{\frac{3 + \sqrt{3}}{4}} \right)^{12}$$

$$\sqrt{73} \quad G_1 = \left( \sqrt{\frac{9+\sqrt{73}}{8}} - \sqrt{\frac{1+\sqrt{73}}{8}} \right)^{24}$$

$$\sqrt{77} \quad G_1 = (8 \pm 3\sqrt{7})^3 \left( \frac{11 \pm \sqrt{7}}{2} \right)^3 \left( \sqrt{\frac{6+\sqrt{11}}{4}} - \sqrt{\frac{3+\sqrt{11}}{4}} \right)^{12}$$

$$\sqrt{81} \quad G_1 = \left( \frac{\sqrt[3]{2(\sqrt{3}-1)} - 1}{\sqrt[3]{2(\sqrt{3}+1)} + 1} \right)^8$$

$$\sqrt{85} \quad G_1 = (\sqrt{5} \pm 2)^8 \left( \frac{\sqrt{85}-9}{2} \right)^6$$

$$\sqrt{93} \quad G_1 = \left( \frac{39-7\sqrt{31}}{\sqrt{2}} \right)^4 \left( \frac{\sqrt{31} \pm 3\sqrt{3}}{2} \right)^6$$

$$\sqrt{97} \quad G_1 = \left( \sqrt{\frac{13+\sqrt{97}}{8}} - \sqrt{\frac{5+\sqrt{97}}{8}} \right)^{24}$$

$$\sqrt{105} \quad \left( \frac{5-\sqrt{21}}{2} \right)^6 (2 \pm \sqrt{3})^6 (\sqrt{5} \pm 2)^4 (6 \pm \sqrt{35})^2$$

$$\sqrt{165} \quad (4-\sqrt{15})^6 (3\sqrt{5} \pm 2\sqrt{11})^4 \left( \frac{\sqrt{15} \pm \sqrt{11}}{2} \right)^6 (\sqrt{5} \pm 2)^4$$

$$\sqrt{273} \quad \left( \frac{15\sqrt{7}-11\sqrt{13}}{\sqrt{2}} \right)^4 \left( \frac{13 \pm 3}{2} \right)^{12} \left( \frac{\sqrt{7} \pm \sqrt{3}}{2} \right)^{12} (2 \pm \sqrt{3})^6$$

$$\sqrt{301} \quad (8 \pm 3\sqrt{7})^3 \left( \frac{23\sqrt{43} \pm 57\sqrt{7}}{2} \right)^3 \times \left( \sqrt{\frac{46+7\sqrt{43}}{4}} - \sqrt{\frac{42+7\sqrt{43}}{4}} \right)^{12}$$

$$\sqrt{141} \cdot (4\sqrt{3} \pm \sqrt{47})^3 \left(\frac{7 \pm \sqrt{47}}{\sqrt{2}}\right)^2 \times$$

$$\left(\sqrt{\frac{18+9\sqrt{3}}{4}} - \sqrt{\frac{14+9\sqrt{3}}{4}}\right)^{12}$$

$$\sqrt{345} \cdot \left(\frac{3\sqrt{3}-\sqrt{23}}{2}\right)^{12} \left(\frac{7\sqrt{23} \pm 15\sqrt{5}}{\sqrt{2}}\right)^4 (\sqrt{5} \pm 2)^8 (2 \pm \sqrt{3})^6$$

$$\sqrt{289} \cdot \left\{ \sqrt{\frac{17+\sqrt{17} + (5+\sqrt{17})\sqrt[3]{17}}{16}} \right.$$

$$\left. - \sqrt{\frac{1+\sqrt{17} + (5+\sqrt{17})\sqrt[3]{17}}{16}} \right\}^{48}$$

$$\sqrt{357} \cdot \left(\frac{\sqrt{7}-\sqrt{3}}{2}\right)^{24} (8 \pm 3\sqrt{7})^6 \left(\frac{11 \pm \sqrt{119}}{\sqrt{2}}\right)^4 \left(\frac{\sqrt{21} \pm \sqrt{17}}{2}\right)^6$$

$$\sqrt{385} \cdot (10 - 3\sqrt{11})^6 (6 \pm \sqrt{35})^6 \left(\frac{\sqrt{11} \pm \sqrt{7}}{2}\right)^{12} (\sqrt{5} \pm 2)^8$$

$$\sqrt{445} \cdot (\sqrt{5}-2)^{12} \left(\frac{\sqrt{445}-21}{2}\right)^6 \left(\sqrt{\frac{13+\sqrt{89}}{8}} \pm \sqrt{\frac{5+\sqrt{89}}{8}}\right)^{24}$$

$$\sqrt{505} \cdot (\sqrt{5}-2)^{14} (\sqrt{101}-10)^6 \left(\frac{5\sqrt{5} + \sqrt{101}}{4} - \sqrt{\frac{105+5\sqrt{105}}{8}}\right)^{12}$$

$$\sqrt{441} \cdot \left(\frac{\sqrt{4+\sqrt{7}} - \sqrt[3]{7}}{2}\right)^{24} \left(\frac{\sqrt{7}-\sqrt{3}}{2}\right)^{12} (2-\sqrt{3})^4 \times$$

$$\sqrt{553} \cdot \left(\frac{\sqrt{3+\sqrt{7}} - \sqrt[3]{6\sqrt{7}}}{\sqrt{3+\sqrt{7}} + \sqrt[3]{6\sqrt{7}}}\right)^{12}$$

$$\left(\sqrt{\frac{143+16\sqrt{79}}{2}} - \sqrt{\frac{141+16\sqrt{79}}{2}}\right)^{12} \left(\sqrt{\frac{100+11\sqrt{79}}{4}} \pm \sqrt{\frac{96+11\sqrt{79}}{4}}\right)^{12}$$

$$\sqrt[294]{117} \cdot \left(\frac{\sqrt{13}-3}{2}\right)^6 \cdot (\sqrt{13}-2\sqrt{3})^4 \cdot \left(\frac{\sqrt{4+\sqrt{3}} \pm \sqrt[3]{3}}{2}\right)^{24}$$

$$\sqrt{133} \cdot (8-3\sqrt{7})^6 \left(\frac{5\sqrt{7} \pm 3\sqrt{19}}{2}\right)^6$$

$$\sqrt{153} \left(\frac{\sqrt{5+\sqrt{17}}}{8} - \frac{\sqrt{\sqrt{17}-3}}{8}\right)^{48} \left(\frac{\sqrt{37+9\sqrt{17}} \pm \sqrt{\frac{33+9\sqrt{17}}{4}}}{4}\right)^8$$

$$\sqrt{145} (\sqrt{5}-2)^6 \cdot \left(\frac{\sqrt{29}-5}{2}\right)^6 \left(\frac{\sqrt{17+\sqrt{145}} \pm \sqrt{\frac{9+\sqrt{145}}{8}}}{8}\right)^{12}$$

$$\sqrt{177} \left(\frac{3\sqrt{59} \pm 23}{\sqrt{2}}\right)^4 (2-\sqrt{3})^{18}$$

$$\sqrt{213} \left(\frac{59 \pm 7\sqrt{71}}{\sqrt{2}}\right)^2 \left(\frac{5\sqrt{3} \pm \sqrt{71}}{2}\right)^3 \left(\frac{\sqrt{21+12\sqrt{3}}}{2} - \frac{\sqrt{19+2\sqrt{3}}}{2}\right)^4$$

$$\sqrt{217} \left(\frac{\sqrt{11+4\sqrt{7}}}{2} - \frac{\sqrt{9+4\sqrt{7}}}{2}\right)^{12} \left(\frac{\sqrt{16+5\sqrt{7}} \pm \sqrt{\frac{12+5\sqrt{7}}{4}}}{4}\right)^{12}$$

$$\sqrt{205} (\sqrt{5}-2)^8 \left(\frac{3\sqrt{5}-\sqrt{41}}{2}\right)^6 \left(\frac{\sqrt{7+\sqrt{41}} \pm \sqrt{\frac{\sqrt{41}-1}{8}}}{8}\right)^{24}$$

$$\sqrt{253} (24-5\sqrt{23})^6 \left(\frac{9\sqrt{23} \pm 13\sqrt{11}}{2}\right)^6$$

$$\sqrt{265} \cdot \left(\frac{\sqrt{53} \pm 7}{2}\right)^6 (\sqrt{5} \pm 2)^6 \left(\frac{\sqrt{89+5\sqrt{265}}}{8} - \frac{\sqrt{81+5\sqrt{265}}}{8}\right)^{12}$$

$$\sqrt{147} \cdot \frac{1}{4} \left\{ \frac{1 \pm \left(2\sqrt{\frac{28}{27}} - \sqrt{\frac{7}{3}}\right)}{2} \right\}^{24}$$

$$g_2 = 1; g_6 = (\sqrt{2}-1)^4; g_{10} = (\sqrt{5}-2)^4;$$

$$\sqrt{14} \cdot \left( \sqrt{\frac{3+\sqrt{2}}{4}} - \sqrt{\frac{\sqrt{2}-1}{4}} \right)^{24}$$

$$\sqrt{18} (5-2\sqrt{6})^4 \cdot \sqrt{22} \cdot (\sqrt{2}-1)^{12}$$

$$\sqrt{30} \cdot (\sqrt{5}-2)^4 (\sqrt{10}-3)^4 \cdot \sqrt{58} \cdot \left( \frac{\sqrt{29}-5}{2} \right)^{12}$$

$$\sqrt{70} \cdot (\sqrt{5}-2)^8 (\sqrt{2}-1)^{12} \cdot \sqrt{46} \cdot \left( \sqrt{\frac{5+\sqrt{2}}{4}} - \sqrt{\frac{1+\sqrt{3}}{4}} \right)^{24}$$

$$\sqrt{42} \cdot \left( \frac{5-\sqrt{21}}{2} \right)^6 (2\sqrt{2}-\sqrt{7})^4 \cdot \sqrt{82} \cdot \left( \sqrt{\frac{13+\sqrt{41}}{8}} - \sqrt{\frac{5+\sqrt{41}}{8}} \right)^{24}$$

$$\sqrt{78} \cdot \left( \frac{\sqrt{13}-3}{2} \right)^{12} (\sqrt{26}-5)^4 \cdot 8$$

$$\sqrt{102} \cdot (\sqrt{2}-1)^{12} (3\sqrt{2}-\sqrt{17})^4$$

$$\sqrt{34} \cdot \left( \sqrt{\frac{7+\sqrt{17}}{8}} - \sqrt{\frac{\sqrt{17}-1}{8}} \right)^{24}$$

$$\sqrt{130} \cdot \left( \frac{\sqrt{13}-3}{2} \right)^{12} (\sqrt{5}-2)^{12}$$

$$\sqrt{190} \cdot (\sqrt{5}-2)^{12} (\sqrt{10}-3)^{12}$$

$$\sqrt{142} \cdot \left( \sqrt{\frac{11+5\sqrt{2}}{4}} - \sqrt{\frac{7+5\sqrt{2}}{4}} \right)^{24}$$

$$\sqrt{90} \cdot (\sqrt{5}-2)^4 \cdot (\sqrt{6}-\sqrt{5})^4 \cdot \left( \sqrt{\frac{3+\sqrt{6}}{4}} - \sqrt{\frac{\sqrt{6}-1}{4}} \right)^{24}$$

$$\sqrt{198} \cdot (\sqrt{2}-1)^{12} (4\sqrt{2}-\sqrt{33})^4 \cdot \left( \sqrt{\frac{9+\sqrt{33}}{8}} - \sqrt{\frac{1+\sqrt{33}}{8}} \right)^{24}$$

246  
 $79 \text{th } \sqrt{t^2 - 6} = u$

then  $u^5 - 2u^4 + u^3 + 2u - 3 = 0$

163.  $t^3 - 2t^2 + 3t = \frac{1}{2}$

$$\sqrt{2} \sqrt{\phi(x) \phi(x^7) \phi(x^9) \phi(x^{63}) + \phi(-x) \phi(-x^7) \phi(-x^9) \phi(-x^{63}) + 4x^4 f^2(x^6) f^2(x^{42})}$$

$$= \phi(x) \phi(x^{63}) + \phi(-x) \phi(-x^{63}) + 4x^{16} \psi(x^4) \psi(x^{126})$$

$$\text{If } \phi(x) = 1 + 6\left(\frac{x}{1-x} - \frac{x^2}{1-x^2} + \frac{x^3}{1-x^3} - \frac{x^5}{1-x^5} + \dots\right)$$

then  $\phi(x) + \phi(-x) = 2\phi(x^4)$

$$\phi^2(x) + \phi(x)\phi(-x) + \phi^2(-x) = 3\phi^2(x^2)$$

$$\frac{1}{x^{\frac{3}{7}}} \frac{f(x^2, -x^6)}{f(-x, -x^6)} = 1 - x^{\frac{1}{7}} \frac{f(x^2, -x^6)}{f(x^2, -x^6)} + x^{\frac{6}{7}} \frac{f(-x, x^6)}{f(-x, -x^6)}$$

$$= \frac{1}{2} \left\{ \frac{3f(x^4)}{x^{\frac{1}{7}} f(-x^7)} + \sqrt{\frac{4f^3(x^4)}{x^{\frac{6}{7}} f^3(x^7)} + \frac{21f^4(x^4)}{x^{\frac{5}{7}} f^4(x^7)} + \frac{28f^5(x^4)}{x^{\frac{4}{7}} f^5(x^7)}} \right\}$$

III:—

$$\frac{u^2}{\omega} + \frac{v^2}{u} - \frac{\omega^2}{v} = 8 + \frac{f^4(x)}{x f^4(x^7)}$$

$$\frac{v}{\omega^2} - \frac{u}{v^2} - \frac{\omega}{u^2} = 5 + \frac{f^4(x)}{x f^4(x^7)}$$

$$u = x^{\frac{1}{56}} f(x^2, -x^6) \quad v = x^{\frac{3}{56}} f(x^2, -x^6) \quad \omega = x^{\frac{4}{56}} f(x^2, -x^6)$$

then  $\frac{u^2}{v} - \frac{v^2}{\omega} + \frac{\omega^2}{u} = 0$

$$u v \omega = x^{\frac{8}{56}} f(x) f(-x)$$

$$\frac{v}{u^2} - \frac{\omega}{v^2} + \frac{u}{\omega^2} = \frac{f(-x)}{x^{\frac{1}{7}} f^2(x^7)} \cdot \sqrt{\frac{f^2(x)}{f^2(x^7)}} + 13x + 49x^2 \cdot \frac{f^5(x)}{f^5(x^7)}$$

$$\text{If } \begin{cases} x^3 + ax + b = y \\ y^3 + ay + b = x \end{cases}$$

Then  $(x^3 + a-1)x + b)(x^2 + ax + a^2 + 1 + a)x$   
 $(x^2 + ax + a^2 + 1 + a)(x^2 + rx + r^2 + 1 + a) = 0$

where  $a, r$  and  $r$  are the roots of the equation

$$z^3 + 2z(a+r) + b = 0$$

$$\text{If } p + q + r + a = x$$

$$q/r + p/a = a$$

$$pqr^2 + q/a^2 + r/p^2 + a^2 = b$$

$$p^2 + q/p^2 + r/a^2 = a^2 + c - 3pvr$$

$$r \cdot p^2 + q^2 + r^2 + a^2 = d + 5(qra - pra) + (pq - q^2 - rp)$$

$$\text{Then } x^5 = 5ax^3 + 5bx^2 + 5cx + d$$

$$\frac{\sqrt[3]{\cos 40} + \sqrt[3]{\cos 80}}{\sqrt[3]{3}(\frac{2}{3} - a)} + \frac{\sqrt[3]{\cos 20} + \sqrt[3]{\sec 20}}{\sqrt[3]{6}(\frac{2}{3} - 1)}$$

$$\sqrt[3]{\sec 40} + \sqrt[3]{\sec 80} + \sqrt[3]{\sec 20} = \sqrt[3]{6}(\frac{2}{3} - 1)$$

If  $a, b, r$  be the roots of  $x^3 - ax^2 + bx - 1 = 0$

$$\text{Then } \sqrt[3]{\frac{a}{a}} + \sqrt[3]{\frac{b}{b}} + \sqrt[3]{\frac{r}{r}} + \sqrt[3]{\frac{a}{a}} + \sqrt[3]{\frac{b}{b}} + \sqrt[3]{\frac{r}{r}}$$

and the roots of

$$\begin{cases} (1) \quad \sqrt[3]{2} - \sqrt[3]{2} + a + b + 3 \text{ solve} \\ (2) \quad \sqrt[3]{2} - \sqrt[3]{2} + a + b + 3 \text{ solve} \end{cases}$$

$$(2) \quad y^6 - y^3(a^2 + b^2 + 9) + (a + b + 3)^3 = 0$$

$$\sqrt[3]{a+6} + 3\sqrt[3]{a^2+9} + 3(a+b) + \sqrt[3]{(a+b)^2 - (a^2+b^2)}$$



298

$$\begin{aligned}
 & + \frac{x-1}{-11x} + \frac{x^2-1}{01x} + \frac{x^2-1}{9x} + \frac{x^2-1}{c^2x} + \frac{x-1}{x} = \\
 & + \frac{x^2-1}{2x} + \frac{x^2-1}{-1x} + \frac{x^2-1}{c^2x} + \frac{x-1}{x} =
 \end{aligned}$$

$$-5 + \frac{20}{8 \left( \frac{x}{7-50} \right)} \cdot \left( \frac{x}{11} \right) \cdot (-5/61 + 99) + (5/1 + 9) = \frac{11}{(3\sqrt{+1})8}$$

$$+ 5 + \left( \frac{9 \cdot 5 \cdot 2}{-5 \cdot 5 \cdot 1} \right) \frac{20}{181} + \left( \frac{2 \cdot 2}{1 \cdot 5} \right) \frac{79}{8} + \left( \frac{x}{11} \right) \frac{79}{2} + 5 = \frac{11}{9}$$

$$+ 5 + \left( \frac{2 \cdot 5 \cdot 2}{-5 \cdot 5 \cdot 1} \right) \frac{27}{61} + \left( \frac{2 \cdot 2}{1 \cdot 5} \right) \frac{77}{81} + \left( \frac{x}{11} \right) \frac{7}{2} + 1 = \frac{11}{7}$$

$$\left\{ (x + \frac{1 \cdot 2 \cdot 2}{2} + \frac{1 \cdot 2 \cdot 2}{1}) \cdot 7x - 1 \right\} \frac{x^2-1}{1} =$$

$$+ 5 + \left\{ (x-1) \cdot 7 \right\} \left\{ \left( \frac{2 \cdot 2}{5 \cdot 1} \right) \cdot 2 + (x-1) \cdot 7 \cdot \left( \frac{x}{11} \right) \cdot 7 + 1 \right\}$$

$$+ 5 + \left\{ (x-1) \cdot 7 \right\} \left\{ \left( \frac{2 \cdot 2}{5 \cdot 1} \right) + (x-1) \cdot 7 \cdot \left( \frac{x}{11} \right) + 1 \right\}$$

$$\left\{ (2x - \frac{11x-1}{91x} + \frac{8x-1}{52} - \frac{5x-1}{72}) \cdot 9 + 1 \right\} \cdot 8 = (10x) \phi (12) \phi + (10x) \phi (10) \phi$$

$$(9x) \phi (12) \phi = (12) \phi (12) \phi + (10x) \phi (12) \phi$$

$$(2x + \frac{11x-1}{11x} - \frac{x^2-1}{2x} - \frac{x^2-1}{5x} + \frac{x-1}{x}) \cdot 8 + 1 = (12) \phi \phi - \frac{(9x) \phi}{(12) \phi (12) \phi (12) \phi} \cdot 8$$

$$+ 8x + \frac{(2x-1)(x-1)(x-1) \cdot x^2-1}{(2x-1)(x-1) \cdot 8x}$$

$$+ \frac{(x-1)(x-1) \cdot 7x-1}{1-x} \cdot \frac{1}{2} + \frac{x-1}{2} \cdot \frac{x-1}{2} =$$

$$+ 8x + \frac{x^2-1}{20x} + \frac{x^2-1}{20x} + \frac{x^2-1}{20x} + \frac{x-1}{2}$$

$$x^2 + \frac{(x^2-1)(x^2-1)(x-1)}{(x^2-1)(x^2-1)(x-1)} \cdot \frac{x-1}{1} =$$

$$x^2 \frac{x-1}{1} + \frac{(x^2-1)(x-1)}{(x^2-1)(x-1)} \cdot \frac{x-1}{1} + \frac{x-1}{x-1} \cdot \frac{x-1}{1} =$$

$$x^2 + \frac{x^2-1}{x-1} + \frac{x^2-1}{x-1} + \frac{x-1}{x-1} + \frac{x-1}{x-1} =$$

$$x^2 + \frac{x^2-1}{x-1} - \frac{x^2-1}{x-1} + \frac{x-1}{x-1} - \frac{x-1}{x-1} =$$

$$\left\{ x^2 + \frac{(x^2-1)(x^2-1)(x-1)}{x^2} + \frac{(x^2-1)(x-1)}{x^2} \right\} (x, x) \text{ II}$$

$$\dots + \frac{x^2-1}{x^2} + \frac{(x^2-1)(x-1)}{x^2} - \frac{x-1}{x^2} \cdot \frac{x-1}{x^2} =$$

$$x^2 + \frac{x^2-1}{x^2} + \frac{x^2-1}{x^2} + \frac{x-1}{x^2} + \frac{x-1}{x^2}$$

$$+ x^2 + \frac{(x^2-1)(x^2-1)(x-1)(x-1)(x-1)}{x^2(x^2+1)(x^2+1)(x+1)} +$$

$$\frac{(x^2-1)(x^2-1)(x-1)(x-1)}{x^2(x^2+1)(x+1)} + \frac{(x^2-1)(x-1)}{x(x+1)} + 1 = \frac{(x, x) \text{ II}}{(x, x) \text{ II}}$$

$$x^2 + \left( \frac{x^2+1}{x} + \frac{x^2+1}{x} \right) +$$

$$\left( \frac{x^2+1}{x} + \frac{x^2+1}{x} \right) + \frac{x+1}{1} = \frac{(x, x) f(x) f(x)}{(x, x) f(x) f(x)}$$

$$x^2 + \frac{x^2+1}{x^2} + \frac{x^2+1}{x^2} + \frac{x^2+1}{x^2} +$$

$$x^2 + \frac{x^2+1}{x^2} + \frac{x^2+1}{x^2} + \frac{x^2+1}{x^2} + 1$$



If an  $n$ th degree series can be expressed in terms of  $M$  and  $N$  only, then,

$$x \frac{d^2 u}{dx^2} - m \frac{du}{dx} - \frac{1}{2} u \text{ can be expressed in}$$

terms of  $M$  and  $N$  only.

Dem. Let  $u = M^{\frac{1}{2}} f\left(\frac{N}{M^3}\right)$ . Then  $x \frac{du}{dx} =$

$$M^{\frac{1}{2}} \frac{d}{dx} \left( M^{\frac{1}{2}} f\left(\frac{N}{M^3}\right) \right) = \frac{N^2}{M^2} \left( M^2 - N^2 \right) \frac{df}{d\left(\frac{N}{M^3}\right)}$$

Cor.  $\frac{d^4 u}{dx^4} = \frac{2L^3}{3M}$  and  $\frac{d^2 u}{dx^2} = \frac{2L^5 M}{2N^2}$

The set of simultaneous equations are useful to find the conditions for as well as the method for expressing a function as the sum of a given set of systems. (Areas and other also).

$$e^x \phi(x) = e^x \phi(x) \cdot e^{\frac{D^2}{2} x + \frac{D}{2} x^2 + bx} = e^x \phi(x) + \frac{D}{2} \phi(1) + \frac{D^2}{2} \phi(2) + \frac{D^3}{6} \phi(3) + \dots$$

$$= e^x \left\{ \frac{\sqrt{1+4x}-1}{2\sqrt{1+4x}} \phi\left(x + \frac{1+\sqrt{1+4x}}{2}\right) + \frac{\sqrt{1+4x}+1}{2\sqrt{1+4x}} \phi\left(x + \frac{1-\sqrt{1+4x}}{2}\right) \right\}$$

$$= e^x \left\{ \frac{1}{2} \phi(x) + \frac{\sqrt{1+4x}-1}{\sqrt{1+4x}} \phi\left(x + \frac{1+\sqrt{1+4x}}{2}\right) + \frac{6\sqrt{1+4x}}{\sqrt{1+4x}+1} \phi\left(x + \frac{1-\sqrt{1+4x}}{2}\right) \right\}$$

If  $n$  is any odd positive integer

$$\text{and } h = e^{-\frac{\pi n}{2}}$$

then all the real roots of  $x$  are included in the following formula and all the imaginary roots can be found by multiplying all the real roots by  $\omega$  and  $\omega^2$  ( $\omega = \sqrt[n]{1}$ ).

$$x = \frac{\sqrt{3}}{2} \left\{ k^2 + \frac{13}{13} k^4 + \frac{38.31}{15} k^6 + 49.52.57 k^8 + \frac{76.79.84.93}{19} k^{10} + 8c \right\}$$

$$+ (-1)^{\frac{n-1}{2}} \left\{ k + \frac{7}{7} k^3 + \frac{19.21}{15} k^5 + \frac{37.39.43}{17} k^7 + 61.63.67.73 k^9 + \frac{91.93.97.103.111}{11} k^{11} + 8c \right\}$$

this series is convergent if  $k < e^{\frac{2}{\pi n}}$  and the greatest value of  $h = e^{-\frac{\pi n}{2}}$  which is  $< \frac{1}{13}$

If  $a, b, c$  are real roots of  $1 - \frac{x^3}{3} + \frac{x^6}{6} - 8c$

$$\text{then } 1 + \frac{x^3}{3} + \frac{x^6}{6} + \frac{x^9}{9} + \frac{x^{12}}{12} + 8c$$

$$= \left(1 + \frac{x^3}{3}\right) \left(1 + \frac{x^6}{6}\right) \left(1 + \frac{x^9}{9}\right) \left(1 + \frac{x^{12}}{12}\right) + 8c$$

the nature of roots and to check the product

The above theorem is very useful & known

$$= c \int_{\infty}^a e^{-\frac{x}{a}} \frac{dx}{x(1+x)}$$

when  $a$  is very great

$$a^{\frac{1}{2}} \left\{ 1 + \left(\frac{\phi(x)}{a}\right)^n \right\} + \left\{ 1 + \left(\frac{\phi(x)}{a}\right)^m \right\} + \left\{ 1 + \left(\frac{\phi(x)}{a}\right)^p \right\} \dots$$

$$= a^{\frac{1}{2}} \int_{\infty}^a \frac{dx}{x(1+x)}$$

when  $a$  is very great.

$$a \left\{ 1 + \frac{\phi(x)}{a} \right\} + \left\{ 1 + \frac{\phi(x)}{a} \right\} + \left\{ 1 + \frac{\phi(x)}{a} \right\} \dots$$

The series is logarithmic when  $x$  is less than  $\frac{1}{2} \text{Limit } \frac{1}{a}$  (inclusive), and convergent when  $\frac{1}{2} \text{Limit } \frac{1}{a}$  & to  $\infty$ .

The series is convergent or divergent according as  $|e^{-\frac{x}{a}} \sin \frac{x}{a}| < \frac{1}{n}$  or  $> \frac{1}{n}$ .

$$e^{-\frac{x}{a}} \sin \frac{x}{a} = 1 + \frac{1}{2} e^{-\frac{x}{a}} \sin \frac{x}{a} + \frac{1}{6} e^{-\frac{x}{a}} \sin \frac{x}{a} + \frac{1}{24} e^{-\frac{x}{a}} \sin \frac{x}{a} + \frac{1}{120} e^{-\frac{x}{a}} \sin \frac{x}{a} + \dots$$

$$u_1 + u_2 + \dots + u_n = 13(u_1 + u_2) \\ u_1 + u_2 + \dots + u_n = 7(u_1 + u_2)$$

$$\left[ \frac{858}{7} = 506 \left\{ \phi(2) + \phi(20) \right\} + 931 \left\{ \phi(11) + \phi(22) \right\} + \dots + \phi(11) \right]$$

$$u_1 + u_2 + \dots + u_{11} = \frac{289}{11} (161u_3 + 252u_{11}) + 161u_{20}$$

$$u_1 + u_2 + \dots + u_{13} = \frac{25}{13} (7u_2 + 11u_7 + 7u_{12})$$

Examplon...

Let  $\alpha = 3m^2 - 13$  and  $\beta = \sqrt{\frac{5}{7}(6m^4 - 45m^2 + 16)}$

$$= \left( \frac{1}{2} - \frac{m^2 - 16}{6\beta} \right) \left\{ \phi(x + \sqrt{\frac{7}{\alpha + \beta}}) + \phi(x - \sqrt{\frac{7}{\alpha + \beta}}) \right\} + \left( \frac{1}{2} + \frac{m^2 - 16}{6\beta} \right) \left\{ \phi(x + \sqrt{\frac{7}{\alpha - \beta}}) + \phi(x - \sqrt{\frac{7}{\alpha - \beta}}) \right\}$$

$$\frac{2(3m^2 - 7)}{5(m^2 - 1) \left\{ \phi(x + \sqrt{\frac{5}{3m^2 - 7}}) + \phi(x - \sqrt{\frac{5}{3m^2 - 7}}) \right\} + 8(m^2 - 4) \phi(x)}$$

$\phi(x)$  as the first approximation and the 2nd

$$\frac{1}{2} \left\{ \phi(x - n + 1) + \phi(x - n + 3) + \dots + \phi(x + n - 1) \right\}$$

$e^{ax}$  can be expanded in ascending powers of  $e^{ax}$  and consequently  $e^{ax}$  can be expanded in ascending powers of  $e^{ax} \sin x$  and hence many transcendental equations can be solved.

$$\frac{x+a_1}{a_1} + \frac{(x+a_1)(x+a_2)}{a_1 a_2} + \frac{(x+a_1)(x+a_2)(x+a_3)}{a_1 a_2 a_3} + \dots$$

$$\begin{aligned}
 &= \frac{x}{1} - \frac{x(1+\frac{a_1}{x})(1+\frac{a_2}{x}) \dots (1+\frac{a_n}{x})}{1} \\
 &+ \binom{a_0 - a_n}{a_1} + \binom{a_0 - a_n}{a_2} + \dots + \binom{a_0 - a_n}{a_n} \\
 &= a_0 - \binom{a_1 a_2 \dots a_n}{a_1 a_2 a_3 \dots a_n} + \dots
 \end{aligned}$$

Let  $u_n$  be said to be  $c$  when  $u_n - c$  can not be made greater than any arbitrary small quantity  $\epsilon$  by making  $n$  sufficiently great.

The expansion of  $\phi(x)$  is said to be a legitimate convergent series if  $\phi(x) =$

the sum of the first  $n$  terms of the expansion of  $\phi(x)$ .  
 The remaining terms are negligible and converge, negligible divergent and negligible divergent series.



306

$$= \frac{x}{2} + \frac{x}{2^2} + \frac{x}{2^3} + \dots + \frac{x}{2^{n-1}} + \frac{x}{2^n} + \frac{x}{2^{n+1}} + \dots$$

$$= \frac{x}{2} + \frac{x}{2^2} + \frac{x}{2^3} + \dots + \frac{x}{2^{n-1}} + \frac{x}{2^n} + \frac{x}{2^{n+1}} + \dots$$

$$= \frac{x}{2} + \frac{x}{2^2} + \frac{x}{2^3} + \dots + \frac{x}{2^{n-1}} + \frac{x}{2^n} + \frac{x}{2^{n+1}} + \dots$$

$$= \frac{x}{2} + \frac{x}{2^2} + \frac{x}{2^3} + \dots + \frac{x}{2^{n-1}} + \frac{x}{2^n} + \frac{x}{2^{n+1}} + \dots$$

$$= \frac{x}{2} + \frac{x}{2^2} + \frac{x}{2^3} + \dots + \frac{x}{2^{n-1}} + \frac{x}{2^n} + \frac{x}{2^{n+1}} + \dots$$

$$= \frac{x}{2} + \frac{x}{2^2} + \frac{x}{2^3} + \dots + \frac{x}{2^{n-1}} + \frac{x}{2^n} + \frac{x}{2^{n+1}} + \dots$$

If  $a, b, c$  be any three quantities, then

$$1.8 \cdot 9 = 3 \cdot 3 \cdot 12 \quad ; \quad 1.3 \cdot 12 \cdot 20 = 5 \cdot 5 \cdot 15 \cdot 16$$

$$1.4 \cdot 20 \cdot 30 = 6 \cdot 24 \cdot 25$$

The ratio between

$\sqrt{Ax+a} \sqrt{Bx+b} \sqrt{Cx+c} \dots$  to  $m$  factors  
and  $\sqrt{px+p} \sqrt{qx+q} \sqrt{rx+r} \dots$  to  $n$  factors

will tend to a finite limit

(27)  $\frac{A^{a+\frac{1}{2}} B^{b+\frac{1}{2}} C^{c+\frac{1}{2}} \dots}{P^{p+\frac{1}{2}} Q^{q+\frac{1}{2}} R^{r+\frac{1}{2}} \dots}$  only when

If the following conditions are satisfied

(1)  $A+B+C+\dots = P+Q+R+\dots$

(2)  $A \cdot B \cdot C \cdot \dots = P \cdot Q \cdot R \cdot \dots$

(3)  $\frac{A}{m} + a + \frac{1}{2} + c + \dots = \frac{P}{m} + p + \frac{1}{2} + r + \dots$

All should all  $A, B, C \dots$  as well as  $P, Q, R \dots$  should all be positive but  $a, b, c \dots$  and  $p, q, r \dots$  may be any quantity even whatever.

N.B.  $a, b, c \dots$  and  $p, q, r \dots$  are necessarily determined from the condition (3) and  $A, B, C \dots$  and  $P, Q, R \dots$  can be found from

From (2) alone find the quantities first

e.g.  $2^2 \cdot 6^6 = 3^3 \cdot 3^3 \cdot 4^4$

and multiply the result by as many 1's as satisfy (1). e.g.  $1! \cdot 1! \cdot 2! \cdot 6^6 = 3^3 \cdot 3^3 \cdot 4^4$

$1! \cdot 1! \cdot 2! \cdot 3! \cdot 4! \cdot 5! \cdot 6^6 = 3^3 \cdot 3^3 \cdot 3! \cdot 4! \cdot 5! \cdot 5!$

and approximately eq. 2 is  $| + \frac{2x}{2x} + \frac{2x}{2x} + \frac{2x}{2x} + \dots$  when  $x$  is great

then  $\frac{|x+a| |x+b| |x+c|}{|x+a| |x+b| |x+c|} = 1$  when  $x = \infty$   
the sum of any other quantities  $b, c, d, \dots$   
of the sum of quantities  $a, b, c, d, \dots$   
 $\frac{a+b+c+d}{a+b+c+d}$

$$\frac{m^2 - 2^2}{5(2^2 + 2x + 1) + 8x} = \frac{a+b}{2^2 + 2x + 1} + \frac{m^2 - 2^2}{8(2^2 + 2x + 1) + 8x}$$

$$\frac{a}{2^2 + 2x + 1} = \left\{ 1 + \left( \frac{x}{2^2 + 2x + 1} \right) \right\}^3 \dots$$
  
$$\frac{b}{8(2^2 + 2x + 1) + 8x} = \left\{ 1 - \left( \frac{x}{2^2 + 2x + 1} \right) \right\}^3 \dots$$

$$\frac{\sin k \pi n - \sin m \pi n}{2^2 + 2x + 1} = \frac{\sin k \pi n + \sin m \pi n}{8(2^2 + 2x + 1) + 8x}$$

$$\frac{a-b}{2^2 + 2x + 1} = \frac{2x}{8(2^2 + 2x + 1) + 8x} + \frac{x}{2^2 + 2x + 1} + \frac{x}{8(2^2 + 2x + 1) + 8x}$$

$$n = \frac{(\sqrt{x^2 - m^2} + \sqrt{x^2 - m^2})}{(\sqrt{x^2 - m^2})}$$

of  $a = \left\{ 1 + \left( \frac{x}{2^2 + 2x + 1} \right) \right\}^3 \dots$  and

$$\frac{m \tan \frac{\pi}{m} - n \tan \frac{\pi}{n}}{m \tan \frac{\pi}{m} - n \tan \frac{\pi}{n}} = \frac{1}{m} + \frac{1}{(m+1)(n+1)} + \dots$$

$$\frac{m-2}{m} = \frac{x}{(m+1)(n^2+x^2)} + \frac{5x+8x}{3x+(m+1)(n^2+x^2)}$$

$$n = \left\{ 1 + \left( \frac{x+1}{m-x} \right)^2 \right\} + \left\{ 1 + \left( \frac{x+1}{m-x} \right)^2 \right\} + \dots$$

$$m = \left\{ 1 + \left( \frac{x+1}{m+x} \right)^2 \right\} + \left\{ 1 + \left( \frac{x+1}{m+x} \right)^2 \right\} + \dots$$

$$= \frac{1}{1+x} + \frac{1}{1+x} + \frac{1}{1+x} + \dots$$

$$2 \left\{ \frac{1}{(x+1)^2+x^2} + \frac{1}{(x+2)^2+x^2} + \dots \right\}$$

$$\frac{\pi x}{m+1} \cdot \frac{e^{\frac{\pi x}{m+1}}}{1} = 1 + \frac{1}{x^2} + \frac{3}{2^2(2+x^2)} + \frac{5}{2^2(2+x^2)} + \dots$$

$$= \frac{1}{1^2+x^2} + \frac{x}{2^2(2+x^2)} + \frac{5x+8x}{2^2(2+x^2)}$$

$$2 \left\{ \frac{1}{(x+1)^2+x^2} + \frac{1}{(x+2)^2+x^2} + \dots \right\}$$

$$= \frac{1}{1^2+x^2} + \frac{1}{2^2} + \frac{1}{2^2+x^2} + \frac{1}{5^2} + \dots$$

$$\frac{1}{1^2+x^2} - \frac{1}{2^2} + \frac{1}{3^2+x^2} - \frac{1}{3^2} + \dots$$







of  $V$  or all the terms  $V \pm \frac{1}{2} \frac{dV}{dx} \pm \frac{1}{6} \frac{d^2V}{dx^2} \pm \dots$  (Binomial expansion)  
 If  $V = (a_1 - a_2 + a_3 - a_4 + \dots)^p$  then the expansion

- (1)  $\{1 - (a + b + c + d + e) + a(c + d) + b(d + e) + ce\}^p$  - kalcade.
- (2)  $\{1 - (a + b + c + d) + (ac + bd) + cd\}^p$  - kalcad
- (3)  $\{1 - (a + b + c)\}^p$  - kalc in part (a, b, c).

(11)  $1 - a_1 a_2$  is part (a) (if  $a_1, a_2$  tend to 0)  
 (12)  $\{1 - (a + b)\}^p$  - kalc is part (a) (if  $a, b$  tend to 0)  
 (13)  $1 - a_1 a_2 a_3$  is part (a) (if  $a_1, a_2, a_3$  tend to 0)

$\frac{a_1}{1 - a_1} - \frac{1}{a_2} - \frac{1}{a_3} - \dots - \frac{1}{a_n}$  is intelligible when

$$= \lim_{n \rightarrow \infty} (a_1 + a_2 + \dots + a_n + b_0 - \dots)$$

$$(a_1 + b_0 - l_1) + (a_2 + b_1 - l_2) + (a_3 + b_2 - l_3) + \dots$$

- part or divergent

or one limit according as  $\frac{1}{p}$  is convergent

$$\frac{1}{p} + \frac{a_1}{p} + \frac{a_2}{p} + \dots + \frac{a_n}{p} + \dots$$

tends to two limits



$$(a_1 - a_2) + (a_2 - a_3) + (a_3 - a_4) + \dots$$

$$= a_1 - \lim_{n \rightarrow \infty} a_n$$

$$\phi(\infty) = \phi(2) - \{\phi(2) - \phi(1)\} - \{\phi(1) - \phi(0)\} - \dots$$

$$= \phi(2) - \{\phi(2) - \phi(1)\} - \{\phi(1) - \phi(0)\} - \dots$$

where  $a_1, a_2$  are increasing quantities

$$S = a_1 - a_2 + a_3 - a_4 + \dots$$

$$\lim_{n \rightarrow \infty} S_{2m+1} - S_n = \lim_{n \rightarrow \infty} a_n$$

$$= a_1 - (a_1 - a_2) - (a_2 - a_3) - (a_3 - a_4) - \dots$$

$$(1+x)^{\frac{1}{2}} \sin(m \tan^{-1} x) =$$

$$\left\{ 1 - \frac{x^2}{\tan^2 \frac{\pi}{4}} \right\} \left\{ 1 - \frac{x^2}{\tan^2 \frac{3\pi}{4}} \right\} \dots$$

$$\left\{ 1 - \frac{x^2}{\tan^2 \frac{\pi}{4}} \right\} \left\{ 1 - \frac{x^2}{\tan^2 \frac{3\pi}{4}} \right\} \dots$$

$$\sin(m \sin^{-1} x) = mx \left\{ 1 - \frac{x^2}{\tan^2 \frac{\pi}{4}} \right\} \left\{ 1 - \frac{x^2}{\tan^2 \frac{3\pi}{4}} \right\} \dots$$

to be omitted factor only

$$\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} + \frac{1}{a_4} + \dots$$

$$= \frac{1}{a_1} - \frac{1}{a_1} + \frac{1}{a_1} - \frac{1}{a_1 + a_2} + \frac{1}{a_1 + a_2} - \frac{1}{a_1 + a_2} + \frac{1}{a_1 + a_2 + a_3} - \dots$$

is intelligible  
or not according as  $\lim_{n \rightarrow \infty} a_n < \infty$  or  $> \infty$

IV.  $F(x+p) \{ \phi(x) + \psi(x) F(x+q) \} = f(x)$

S. substitute  $F(x) = \frac{F_1(x+q)}{F_1(x+p)}$  then

$\phi(x) F_1'(x+p+q) + \psi(x) F_1'(x+q) = f(x) F_1'(x+p)$

I.  $\frac{x^5-a}{x^5-a} = \frac{y^5-c}{y^5-c} = 5(x+y-1)$   
 II.  $\frac{x^7-a}{x^7-a} = \frac{y^7-c}{y^7-c} = 7(xy-1)$

Substitution:  $x = a+p+z$ ,  $y = a+p+z+d$ ,  $d = a+p$

III.  $x+z+2+x=a$   
 $px+qy+nz+m=b$   
 $p^2x+q^2y+n^2z+m^2=c$   
 $p^3x+q^3y+n^3z+m^3=d$

$\frac{1-cp}{x} + \frac{1-cq}{y} + \frac{1-cn}{z} + \frac{1-cm}{w} = a+bl+e^2+af^2$

Find the sum of the right hand side by converting it into a continued fraction or by using indeterminate coefficients and then split up the result into partial fractions.

IV.  $x^2+ay=l; y^2+cx=d$   
 $x = a+p+z$   
 $y = -\frac{a}{2}(a+p+z)$   
 $d-py = \text{sum of } l$

I.  $aF(x+b) + lF(x+a) + cF(x+q) + b = \phi(x)$

Write  $\phi(x)$  as  $\int_{\beta}^{\alpha} u^x v dx$  where  $v$  is a function of  $x$ , then

$$F(x) = \int_{\beta}^{\alpha} u^x v dx = \int_{\beta}^{\alpha} a u^x p + l u^x q + c u^x r + b dx$$

II.  $\phi(x) F(x+b) + \psi(x) F(x+a) = f(x)$

Find a function  $\chi(x)$  so that  $\frac{\chi(x+b)}{\chi(x)} = \psi(x)$

Now let  $F(x) = F_1(x) \chi(x)$ , then we have

$$\frac{F_1(x+b) + F_1(x+a)}{F_1(x)} = \frac{\phi(x) \chi(x+b)}{\chi(x)}$$

III. If  $a, l, c$  are constants in A.P.

and  $u, v, w$  are functions of  $x$  in G.P.

Solve  $uF(x+a) + vF(x+a) + wF(x) = \phi(x)$

Find  $\chi(x)$  so that  $\frac{\chi(x+a)}{\chi(x)} = \sqrt{\frac{u}{v}}$  or  $\frac{u}{v}$

and substitute  $F(x) =$

$$\chi(x) \left\{ \sqrt{u} F_1(x+\frac{a}{2}) - \sqrt{v} F_1(x+\frac{a}{2}) \right\}$$

If  $\phi(x)$  vanishes for  $a, l, c, d$  etc. of  $x$ , then

(ii) the coeff<sup>s</sup> of  $x^{n-1}$  in the expansion of  $\frac{1}{\phi(x)}$

$$= -\frac{1}{a^n \phi'(a)} - \frac{1}{c^n \phi'(c)} - \frac{1}{d^n \phi'(d)} - \dots - \theta(m)$$

where  $\lim_{m \rightarrow \infty} K^n \theta(m) = 0$  for any value of  $K$ , and

$\theta(m)$  is 0 in many cases.

(21) the expansion of the function

$$\frac{\phi(x)}{\phi(x)} + \frac{(a-x)\phi'(a)}{1} + \frac{(a-x)^2\phi''(a)}{2!} + \dots$$

is convergent for all values of  $x$ .

If  $\phi(x) = 0$  for the values of  $a, l, c, d$  etc. of  $x$ ,

and if  $|a|$  be the nearest to 0, then

(i) the expansion of  $\phi(x)$  is convergent if  $|x| < |a|$

and divergent if  $|x| > |a|$ .

(22) the  $n^{\text{th}}$  coeff<sup>s</sup> of  $x^{n-1}$  in the expansion =  $\left\{ \phi'(a) \right\}^2$

$$= - \left[ \frac{d}{dx} \frac{\phi(x)}{\phi(x)} \right]_{x=a}$$

If  $\phi(x) = a_0 + a_1x + a_2x^2 + \dots$  when the

coeff<sup>s</sup> are positive or least after some terms  
 one of them, then  $\lim_{n \rightarrow \infty} \phi\left(\frac{a}{n}\right) = \infty$ .

and  $(\sqrt{1} - \sqrt{2} + \sqrt{3} - \sqrt{4})^2 = \frac{2}{\pi}$ .

oscillates between  $(\sqrt{1} - \sqrt{2} + \sqrt{3} - \sqrt{4})^2 + \frac{2}{\pi}$

e.g.  $1 - \frac{\sqrt{2}}{2} + (\frac{\sqrt{3}}{2} + \frac{1}{2}) - (\frac{\sqrt{2}}{2} + \frac{1}{2}) + (\frac{\sqrt{3}}{2} + \frac{1}{2}) + (\frac{\sqrt{2}}{2} + \frac{1}{2}) - (\frac{\sqrt{3}}{2} + \frac{1}{2}) + \frac{2}{\pi}$

oscillates between  $(a_1 - a_2 + a_3 - a_4 + a_5 - a_6 + a_7 - a_8 + a_9 - a_{10})^2 + \frac{2}{\pi}$

$a_1^2 - 2a_1a_2 + (2a_1a_3 + a_2^2) - (2a_1a_4 + 2a_2a_3) + 8c$

$$= \int_0^{\infty} \frac{2}{x} \frac{e^{-\frac{2}{\pi x}} + e^{-\frac{2}{\pi x}}}{\phi(x) + \phi(x)} dx$$

$$\phi(1) - \phi(2) + \phi(3) - \phi(4) + \phi(5) - \phi(6) + \dots$$

$$= \int_0^{\infty} \frac{2}{x} \phi(x) dx - \int_0^{\infty} \frac{e^{-\frac{2}{\pi x}}}{e^{-\frac{2}{\pi x}} - \phi(x)} dx$$

$$\phi(1) + \phi(2) + \phi(3) + \phi(4) + \phi(5) + \dots$$

$$= \int_0^{\infty} \frac{e^{-\frac{2}{\pi x}}}{e^{-\frac{2}{\pi x}} - \phi(x)} dx$$

$$\frac{1}{2} \phi(1) - \phi(2) + \phi(3) - \phi(4) + \dots$$

$$= \int_0^{\infty} \frac{e^{-2x} x^{2n-1} dx + 2 \int_0^{\infty} \frac{e^{-2x}}{2^{2n-1} \cos(\frac{\pi}{2} - x)} dx}{e^{2\pi x}}$$

$$\text{Cor. } 1^{n-1} e^{-x} + 2^{n-1} e^{-2x} + 3^{n-1} e^{-3x} + \dots$$

$$= \int_0^{\infty} \phi(x) dx + \int_0^{\infty} \frac{e^{-\frac{2}{\pi x}}}{\phi(x) - \phi(x)} dx$$

$$\frac{1}{2} \phi(1) + \phi(2) + \phi(3) + \phi(4) + \dots$$

is zero, infinite or finite. when  $a$  and  $b$  divergent or oscillating according as  $l_1 - 2a$  does not contain any log. functions.

The product of the two series

$$\int_{-\infty}^{\infty} x^{n-1} \frac{\phi(x) - \phi(-x)}{2} dx = \sin \frac{\pi}{2} \int_{-\infty}^{\infty} x^{n-1} \phi(x) dx$$

$$\int_{-\infty}^{\infty} x^{n-1} \frac{\phi(x) + \phi(-x)}{2} dx = \cos \frac{\pi}{2} \int_{-\infty}^{\infty} x^{n-1} \phi(x) dx$$

$0 = \int_{-\infty}^{\infty} x^{n-1} \phi(x) dx = \phi(0) - \phi(\infty)$  when  $n = 0$

$$= \frac{1}{2} \int_{-\infty}^{\infty} \frac{\phi(x) + \phi(-x)}{\cos \pi x + \cos \pi a} dx$$

$$\phi(l-a) - \phi(l+a) + \phi(l-a) - \phi(l+a) - 2c$$

$$+ \frac{2a(2a+1)}{(l-a)(l-a-1)} \cdot \frac{\sqrt{}}{e^{-(a+2)n} + (l+a+1)(l+a+2)}$$

$$= \frac{1}{2} \cdot \frac{(l-a)}{(l-a-1)} \cdot \frac{(l-a)}{(l-a-1)} \cdot \frac{1}{2a} \cdot \frac{1}{l-a} \cdot \frac{1}{(a+1)} \cdot e^{-(a+1)n}$$

$$\int_{-\infty}^{\infty} \frac{1 + (\frac{x}{a})^2}{1 + (\frac{x}{a+1})^2} \cdot \frac{1 + (\frac{x}{a+1})^2}{1 + (\frac{x}{a+2})^2} \cdot \dots \cdot \cos \pi x dx$$

$$= \frac{2\sqrt{a-1}\sqrt{a}}{\sqrt{a-1}\sqrt{a}} \left\{ e^{-n} - \frac{a-1}{a-1} \cdot e^{-2n} + \frac{(a+1)(a-1)}{(a-1)(a-1)} e^{-3n} \dots \right.$$

$$\int_0^{\infty} \frac{(1 + \frac{1}{x})(1 + \frac{1}{x^2}) \dots (1 + \frac{1}{x^n})}{\cos x dx}$$

then  $u_{n+2} = \frac{1}{x^2} u_n + \frac{d u_n}{dx}$

$$u_n = \left(\frac{1}{x}\right)^{n+1} e^{-\frac{1}{x}} \left\{ 1 - \frac{1}{x} + \frac{1}{2x^2} + \dots \right\}$$

$$\int u_n = \int e^{-\frac{1}{x}} \left\{ e^{-\frac{1}{x}} \left( \frac{1}{x} \right)^{n+2} + \dots \right\} dx$$

$$= \frac{1}{\pi} \frac{\sin 2\pi n - \cos 2\pi n}{\sin 2\pi n} + \dots$$

$$= \frac{1}{\pi} \frac{\sin 2\pi n - \cos 2\pi n}{\sin 2\pi n} + \dots$$

$$= \frac{1}{\pi} \frac{\sin 2\pi n + \cos 2\pi n}{\sin 2\pi n} + \dots$$





Let  $\phi$  be a function defined by the relation  $\phi(x) = f(x) \phi\{f(x)\}$  where  $f$

is a known function.

Then  $\int_a^x \phi(x) dx$  is always constant

whenever be the value of a. Let this C.

Denote  $f f(x)$  by  $f^2(x)$ ,  $f f f(x)$  by  $f^3(x)$  etc.

then (1)  $f^m(x) - f^{m+1}(x)$

(2)  $\int_{f^m(x)}^{f^{m+1}(x)} \phi(x) dx = (m-m)C$

(3)  $\int_a^x \phi(x) dx$  is of an order lower than  $f^m(x)$ .

$\int_a^x f^n(x) dx = \psi^n(x) + \frac{1}{n} \cdot \psi'(x) + \frac{1}{n^2} \psi''(x) + \dots$

then (1)  $\psi^n(x) = f^n(x) = x$

(ii)  $\frac{d}{dx} f^n(x) \cdot \psi'(x) = \frac{d}{dx} x$

(3)  $\psi^n(x) = \psi'(x) \cdot \frac{dx}{d\psi^n(x)}$

(4)  $\psi'(x) = \frac{\phi(x)}{C}$

If  $\psi$  be a function defined by the relation

$\psi^n(x) = x$  and  $\psi'(x) \phi(x) = \frac{dx}{d\psi^n(x)}$  where  $\phi$  is a known function, then  $\int_a^x \phi(x) dx$

then  $\psi = \psi^n(x) + \frac{1}{n} \psi'(x) + \frac{1}{n^2} \psi''(x) + \dots$

$$x^2(2x+1) = 11$$

$$7/8(1-a) = 64 \cdot 9 \cdot \frac{1+27a}{8} \text{ and}$$

$$\text{Then } 7/8(1-a) = 64 \cdot 9 \cdot \frac{1+27a}{8}$$

of  $a$  and  $B$  be of the I or IX degree

$$\{x^2 + 2x + 1\}^2 + \frac{1+27a}{8} \cdot 64 \cdot 9 \cdot (x^2 + 1) = 0$$

$$\{x^2 + 2x + 1\}^2 + \frac{1+27a}{8} \cdot 64 \cdot 9 \cdot (x^2 + 1) = 0$$

$$x^2 + 2x + 1 + \frac{1+27a}{8} \cdot 64 \cdot 9 \cdot (x^2 + 1) = 0$$

$$\text{Then } \frac{3\sqrt{a}}{8} + \sqrt{(1-a)(1-a)} = 1$$

$$\text{of } a = \frac{b(2+b)^2}{(1+b)^3} \text{ and } a = b^2(2+b)$$

$$\text{Then } p^2 + 1 + \frac{p}{q} = \left(\frac{p}{q}\right)^2 + \left(\frac{p}{q}\right)^3$$

$$\text{of } p = \frac{f(x^2)f(x^2)f(x^2)}{f(x^2)f(x^2)} \text{ and } q = \frac{x^2 f(x^2) f(x^2)}{f(x^2) f(x^2)}$$

$$\text{Then } x^2 - 3x^2 + 2x^2 = 90$$

$$\frac{f(x^2)f(x^2)f(x^2)}{f(x^2)f(x^2)} = x^2 \text{ and } \frac{x^2 f(x^2) f(x^2)}{f(x^2) f(x^2)} = x^2$$

$$\text{Then } p^2 + \frac{p}{q} = \left(\frac{p}{q}\right)^2 + \frac{p}{q} + \left(\frac{p}{q}\right)^3$$

$$\text{of } p = \frac{f(x^2)f(x^2)f(x^2)}{f(x^2)f(x^2)} \text{ and } q = \frac{x^2 f(x^2) f(x^2)}{f(x^2) f(x^2)}$$

$$\text{Then } p^2 + \frac{p}{q} = \left(\frac{p}{q}\right)^2 + \left(\frac{p}{q}\right)^3 + 4$$

$$\text{of } p = \frac{f(x^2)f(x^2)f(x^2)}{f(x^2)f(x^2)} \text{ and } q = \frac{x^2 f(x^2) f(x^2)}{f(x^2) f(x^2)}$$

is understood by interchanging a and c and at the same time  
and d. It is further to be observed that if a and x  
and in one of opposite signs

$$+ c^2 d^3 f(a^2, b^2, c^2, d^2) + c^2 d^2 f(a^2, b^2, c^2, d^2) + \dots$$

$$f(a, b) + c f(a, b, c) + d f(a, b, c, d) + \dots$$

$$\left\{ x + \frac{1-x^{2n}}{1-x^2} - \frac{1-x^{2n-1}}{1-x} + \frac{1-x^{2n-2}}{1-x^2} - \dots \right\} (1-x)^n f =$$

$$(1-x) + 2x(1-x^2) + 3x^2(1-x^4) + 4x^3(1-x^6) + \dots$$

$$\left\{ \begin{aligned} (1-x)^n \psi\left(\frac{2n}{2}\right) f_{2n} &= (2^{2n} - 2^{2n-1}) f(1-x) - (2^{2n-1} - 2^{2n-2}) f(1-x^2) + \dots \\ (1-x)^n \psi\left(\frac{2n-1}{2}\right) f_{2n-1} &= (2^{2n-1} - 2^{2n-2}) f(1-x) - (2^{2n-2} - 2^{2n-3}) f(1-x^2) + \dots \end{aligned} \right.$$

$$f(a, b) + c f(a, b, c) + d f(a, b, c, d) + \dots$$

$$f(a, b) + c f(a, b, c) + d f(a, b, c, d) + \dots$$

$$f(a, b) + c f(a, b, c) + d f(a, b, c, d) + \dots$$

$$f(a, b) + c f(a, b, c) + d f(a, b, c, d) + \dots$$

$$f(a, b) + f(a, c) + f(a, d) + f(a, e) = (m + n + p) f$$

$$(2x + \frac{x-1}{x} - \frac{x-1}{x} + \frac{x-1}{x} - \frac{x-1}{x}) 9 + 1 =$$

$$(2x) f(x) + 4x f(x) + 4x f(x)$$

$$(p-a)f + (q-a)f + (r-a)f =$$

$$f(a, b) + f(a, c) + f(a, d) + f(a, e) =$$

$$\{ (p-a)f + (q-a)f - (p-a)f \} =$$

$$(r-a)f + f(a, b) - f(a, c) - f(a, d) =$$

$$(r-a)f + \frac{2}{1-i} + (r-a)f = (r-a)f$$

$$\frac{f(a, b) - f(a, c) - f(a, d)}{x - x^2} = (r-a)f - (r-a)f$$

$$2y = x + \sqrt{x^2 - x^2}$$

$$f x = y + \sqrt{x^2 - y^2}$$

where  $\alpha\beta = 1$  and  $\alpha - \beta = 11 + \frac{y}{2}$

where  $\gamma\delta = 1$  and  $\gamma - \delta = 11 + 12\frac{x}{2}$

$$\sqrt{5} - \sqrt{3} = 1 + \frac{y}{x}$$

$$\sqrt{5} - \sqrt{3} = \frac{5}{2} + 1$$

$$\sqrt{\frac{f(a, b)}{f(a, c)}} = \frac{f(a, b) - f(a, c)}{f(a, c)}$$

$$f x = y + \sqrt{\frac{y}{x} - y^2}$$

$$\text{then } 3y = x + \sqrt{9x^2 - x^2}$$





$$p^2q - spq = p^3 - 2p^2q - 2pq^2 + q^3$$

$$* \int p = \int \frac{f(x) f'(x)}{f(x)^2} = \frac{1}{f(x)} + \frac{f'(x)}{f(x)^2}$$

$$\int p = \frac{f(x)}{f'(x)} = \frac{f(x)}{f'(x)} + \frac{f'(x)}{f(x)^2}$$

$$* \int p = \frac{f(x)}{f'(x)} = \frac{f(x)}{f'(x)} + \frac{f'(x)}{f(x)^2}$$

$$p^2 + \frac{p}{25} = \left(\frac{p}{5}\right)^3 - 4\left(\frac{p}{5}\right)^2 + \left(\frac{p}{5}\right)$$

$$\int p = \frac{f(x)}{f'(x)} = \frac{f(x)}{f'(x)} + \frac{f'(x)}{f(x)^2}$$

and  $\sqrt{a} + \sqrt{a} + \sqrt{a} = \sqrt{6+6+6}$   
 $\sqrt{a} + \sqrt{a} + \sqrt{a} = \sqrt{a+6+6}$   
 $a^2 + 6a - 1 = 0$

So a, b are roots of the equation

$$= (n+\frac{1}{2}) \left\{ \frac{1}{2n^2+2n+1} + \frac{1}{2n^2+2n+1} + \frac{1}{6} + \frac{1}{6} \right\}$$

$$= (n+\frac{1}{2}) \left\{ \frac{1}{n^2+n} + \frac{1}{n^2+n} + \frac{5(n+n)}{12} + \frac{1}{2} + \frac{1}{2} \right\}$$

$$\frac{1}{n^2+n} + \frac{1}{n^2+n} + \frac{5(n+n)}{12} + \frac{1}{2} + \frac{1}{2}$$

$$(pQ)^2 + 5 + \frac{(pQ)^2}{9} = \left(\frac{p}{9}\right)^2 - \left(\frac{Q}{9}\right)^3$$

$$\int p = \frac{f(x)}{f'(x)} \text{ and } Q = \frac{x^n f(x+1)}{f'(x+1)} \text{ this}$$

$$= 1+t+t^2+2t^2+5t^4+17t^5+8t \text{ ansmp.}$$

$$2 \left\{ 1 - \left(\frac{1+t}{1-t}\right)^2 + \left(\frac{1+t}{1-t}\right)^4 - \left(\frac{1+t}{1-t}\right)^6 + \left(\frac{1+t}{1-t}\right)^8 - \dots \right\}$$

$$= \frac{1-t^{-2}}{1-t^2} + \frac{1-t^{-4}}{1-t^2} + \frac{1-t^{-6}}{1-t^2} + \frac{1-t^{-8}}{1-t^2} + \dots$$

$$= \frac{1-t^{-2}}{1-t^2} + \frac{1-t^{-4}}{1-t^2} + \frac{1-t^{-6}}{1-t^2} + \frac{1-t^{-8}}{1-t^2} + \dots$$

$$= \frac{1-t^{-2}}{1-t^2} + \frac{1-t^{-4}}{1-t^2} + \frac{1-t^{-6}}{1-t^2} + \frac{1-t^{-8}}{1-t^2} + \dots$$

$$= \frac{1-t^{-2}}{1-t^2} + \frac{1-t^{-4}}{1-t^2} + \frac{1-t^{-6}}{1-t^2} + \frac{1-t^{-8}}{1-t^2} + \dots$$

No. of the form  $p^2 q^3$

$$= 2.1732542\sqrt{2} - 1.458455\sqrt{2}$$

$$= \sqrt[3]{4.72308422} - \sqrt[3]{3.102272}$$

$$= \int_0^{\infty} e^{-ax} \sin kx dx$$

$$= \int_0^{\infty} \frac{e^{-ax} \sin kx}{e^{ax} - 1} dx$$



The no of terms less than according to formula

15	50	14.9
300	62	61.9
1000	168	168.9

$$\int \frac{dx}{x} = \log x + c = \log \log a + c$$

$$\left\{ \frac{1}{\log a} + \frac{1}{\log a} + \frac{1}{\log a} + \dots \right\} - x$$

the  $\int f(x) = \frac{f(x)}{x} = \frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

and  $\frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

the  $\int f(x) = \frac{f(x)}{x} = \frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

and  $\frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

the  $\int f(x) = \frac{f(x)}{x} = \frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

and  $\frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

the  $\int f(x) = \frac{f(x)}{x} = \frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

and  $\frac{f(x)}{x^2} = \frac{f(x)}{x^3} = \dots$

$$\int_0^{\infty} f(x) \cdot p = \int_0^{\infty} \frac{f(x) \cdot x^{a-1}}{x^{a-1}} dx = \int_0^{\infty} \frac{f(x) \cdot x^{a-1}}{x^{a-1}} dx = \int_0^{\infty} f(x) \cdot x^{a-1} dx$$

$$p \cdot \frac{p}{13} = \left(\frac{p}{8}\right)^2 = -2 \cdot \frac{p}{8} - 3 - 3 \cdot \frac{p}{8} + \left(\frac{p}{8}\right)^2$$

Then  $\int_0^{\infty} x^{a-1} F(x) dx = \int_0^{\infty} x^{a-1} f(x) dx$

and  $\int_0^{\infty} F(x) f(x) dx = \frac{a+p}{1}$

$$\int_0^{\infty} \frac{a}{x} = \frac{a-p}{1} = p$$

Then  $\int_0^{\infty} \psi(x) \cdot \frac{f(x) + f(-x)}{2} dx = \frac{3}{\pi} \phi(m)$

$$\int_0^{\infty} \phi(x) \cdot \frac{F(x) + F(-x)}{2} dx = \psi(m)$$

value of p, then

or  $\int_0^{\infty} x^{p-1} F(x) dx = \int_0^{\infty} x^{p-1} f(x) dx = \frac{\sin p \pi}{\pi}$  for all values of a and b.

$$\frac{1}{x^2} + \frac{1}{x^3} + \frac{1}{x^4} + \dots = \frac{1}{x^2} \left( 1 + \frac{1}{x} + \frac{1}{x^2} + \dots \right) = \frac{1}{x^2} \cdot \frac{1}{1 - \frac{1}{x}} = \frac{1}{x^2} \cdot \frac{x}{x-1} = \frac{1}{x(x-1)}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{A}{x} + \frac{B}{x-1}$$

$$1 = A(x-1) + Bx$$

$$1 = Ax - A + Bx = (A+B)x - A$$

$$\begin{cases} A+B = 0 \\ -A = 1 \end{cases} \Rightarrow \begin{cases} A = -1 \\ B = 1 \end{cases}$$

$$f(x) = \frac{-1}{x} + \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$f(x) = \frac{1}{x(x-1)} = \frac{1}{x} - \frac{1}{x-1}$$

$$\text{Hence, if } \begin{cases} x = \frac{b-2y-2z}{a} \\ y = \frac{c-2x-2z}{b} \\ z = \frac{a-2x-2y}{c} \end{cases} \text{ then, } \begin{cases} x^2+2yz = a^2+2bx \\ y^2+2zx = b^2+2cy \\ z^2+2xy = c^2+2az \end{cases}$$

then  $x^2+2yz = (A^2+2Bc)(P^2+2q/r) + (B^2+2cA)(q^2+2r)$   
 If  $x = Ap + Bq + Cr, y = Bp + Cq + Ar, z = Cp + Aq + Br$

$$\left\{ \begin{aligned} a^2 + b^2 + c^2 - 2abc &= (x^2 + y^2 + z^2 - 2xyz) \\ a + b + c &= (x + y + z) \\ \frac{a}{x} + \frac{b}{y} + \frac{c}{z} &= 0 \end{aligned} \right.$$

then,  $\begin{cases} x - y = \frac{b - c}{a} \\ y - z = \frac{c - a}{b} \\ z - x = \frac{a - b}{c} \end{cases}$  and  $L = (b-a)(c-b)(c-a)$

Let  $\theta = x^2 + y^2 + z^2 + 2xy + 2yz + 2zx$  and  $L = (x-y)(y-z)(z-x)$

$$\begin{cases} x^2 + 2yz = a \\ y^2 + 2zx = b \\ z^2 + 2xy = c \end{cases}$$

Similarly for  $y^2 + 2zx$  and  $z^2 + 2xy$  also

then,  $x^2 + 2yz = (x^2 - 5x^2 + 4) a \theta +$

$$\left\{ \begin{aligned} x &= a + \beta + \gamma(m-1) \\ y &= \beta + \gamma + \alpha(m-1) \\ \text{and } z &= \gamma + \alpha + \beta(m-1) \end{aligned} \right.$$

$$\left. \begin{array}{l} 12n+11 = 42 \\ 12n+7 = 44 \\ 12n+5 = 46 \\ 12n+1 = 48 \end{array} \right\}$$

$$\left. \begin{array}{l} 8n+7 = 43 \\ 8n+5 = 43 \\ 8n+3 = 43 \\ 8n+1 = 37 \end{array} \right\}$$

of the form  $4n+1 = 80$   
 of the form  $6n+1 = 80$   
 of which there

The no of primes was between 1000 and 1000 = 166.

1-2-3-5-6-7-10-11-13-14-15-17	-19+21-22-23-26-29-30-31+33+34+35-37	+38+39-41-42-43+46-47+51-53+55+57+58	-59-61+62+65-66-67+69-70-71-73+74+77	-78-79+82-83+85+86+87-89+91+93+94+95-97	-101-102-103-105+106-107-109-110+111-113-114+115	+118+119+122+123-127+129-130-131+133+134-137-138	-139+141+142+143+145+146-149-151-154+155-157+158	+159+161-163-165+166-167-170-173-174+177+178-179	-181-182+183+185-186+187-190-191-193+194-195-197-199	+201+202+203+205+206+209+210-211+213+214+215+217	-218-219+221-222-223+226-227-229-230-231-233+235	+237-238-239-241-246+247+249-251+253+254-255-257	-258+259+262-263+265-266+267-269-271-273+274-277	+278-281-282-283-285-286+287-290+291-293+295+298+299	-319+321-322+323+326+327+329+330-331+334+335-337	+339+341+345+346-347-349-353-352+355-357+358-359	-362+363-366-367-370+371-373-374+377-379+381+382	-383-385+386-387-389+390+391+393+394+395-397+398-399
-------------------------------	--------------------------------------	--------------------------------------	--------------------------------------	---	--	--	--	--	--	--	--	--	--	--	--	--	--	--

$$\int_{-\infty}^{\infty} x^2 \phi\left(\frac{x}{\sigma}\right) dx = 1 - \frac{\sigma}{1} + \left(\frac{\sigma}{2}\right)^2 - \left(\frac{\sigma}{3}\right)^3 + \left(\frac{\sigma}{4}\right)^4 - \dots$$

$$-\phi(-1) + 2\phi(-2) - 3\phi(-3) + (1) \phi(1) + 2\phi(2) - 3\phi(3) + \dots$$

$$\int_{-\infty}^{\infty} \phi(x) dx = \phi(1) + \phi(2) + \frac{1}{\phi(1)} + \frac{1}{\phi(2)} + \frac{1}{\phi(3)} + \frac{1}{\phi(4)} + \dots$$

then  $n = (p + \frac{1}{2}) \log n - n$  very nearly

$$(1 + \frac{1}{2})^n + \frac{1}{2} (1 + \frac{1}{2})^{n-1} + \dots + \frac{1}{n} (1 + \frac{1}{2})^1 = \log p$$

$$\int_{-\infty}^{\infty} e^{2\pi i x} dx = \int_{-\infty}^{\infty} \frac{e^{2\pi i x}}{(1+x)^2} dx$$

$$\int_{-\infty}^{\infty} \frac{e^{2\pi i x}}{(1+x)^2} dx = \int_{-\infty}^{\infty} \frac{e^{2\pi i x}}{(1+x)^2} dx$$

where  $e = .5772\dots$

$$\int_{-\infty}^{\infty} x dx = c + \log \log x + \frac{1}{\log x} + \frac{1}{(\log x)^2} + \frac{1}{(\log x)^3} + \dots$$

where  $n - \log x = \delta$

$$\left\{ \frac{1}{8} \frac{(\log x)^2}{28(1-\delta)} - \frac{1}{155} \frac{1}{8(1-\delta)^2(2-3\delta^2)} \right\} + \left\{ \frac{1}{4} \frac{1}{8^2(1-\delta)^2} - \frac{1}{135} \frac{1}{8^2(1-\delta)^2} \right\} + \dots$$

where  $n = 1.45136380$

$$\int_{-\infty}^{\infty} x dx = x \left\{ \frac{1}{\log x} + \frac{1}{(\log x)^2} + \frac{1}{(\log x)^3} + \dots \right\} + \dots$$

and  $\psi(1) = \phi(1) - \phi(2) = (1) \psi(1) - \psi(2)$

$\psi(2) + (\psi(3)) \psi(2) + (\psi(4)) \psi(3) + (\psi(5)) \psi(4) + (\psi(6)) \psi(5) = (1) \psi(1) - \psi(2)$

$\psi(2) + (\psi(3)) \psi(2) + (\psi(4)) \psi(3) + (\psi(5)) \psi(4) = \psi(2) - (\psi(3)) \psi(2) + (\psi(4)) \psi(3) - (\psi(5)) \psi(4)$

$\psi(2) - (\psi(3)) \psi(2) + (\psi(4)) \psi(3) - (\psi(5)) \psi(4) - (\psi(6)) \psi(5) = (1) \psi(1) - \psi(2)$

$\psi(1) = \psi(2) + (\psi(3)) \psi(2) + (\psi(4)) \psi(3) + (\psi(5)) \psi(4) + (\psi(6)) \psi(5)$

$$= \frac{1}{2} \left\{ \frac{1}{2} B_2 \left( \log P \right)^2 + \frac{5}{6} B_4 \left( \log P \right)^4 + \frac{7}{8} B_6 \left( \log P \right)^6 + \frac{9}{10} B_8 \left( \log P \right)^8 + \frac{11}{12} B_{10} \left( \log P \right)^{10} \right\}$$

$$n = \int_P \frac{1}{x} dx - \frac{1}{2} \int_P \frac{1}{x^2} dx + \frac{1}{6} \int_P \frac{1}{x^3} dx - \frac{1}{30} \int_P \frac{1}{x^5} dx + \frac{1}{42} \int_P \frac{1}{x^7} dx - \frac{1}{30} \int_P \frac{1}{x^9} dx + \dots$$

where  $1$  and  $P$  are any two numbers

$$\int_0^{\infty} (e^{-ax} + e^{-a^2x} + e^{-a^3x} + \dots) \log x dx = \frac{1}{a}$$

If  $P$  is a function of  $n$  such that

Let  $f(x) = 0$  and  $\phi(x) = 0$ . Then  $\phi(x) \phi'(x) = 0$  and the roots of  $f(x) = 0$

Hence  $\frac{dP}{dn} = \frac{P \log P}{1} (P - \frac{1}{P} - \frac{2}{\sqrt{P}} - \frac{3}{\sqrt[3]{P}} - \frac{4}{\sqrt[4]{P}} + \frac{5}{\sqrt[5]{P}} + \frac{6}{\sqrt[6]{P}} - \dots)$

$\int \log \frac{P}{P} dn = \int \frac{dP}{P} - \frac{1}{2} \int \frac{dP}{P^{3/2}} - \frac{1}{3} \int \frac{dP}{P^{4/3}} + \frac{1}{4} \int \frac{dP}{P^{5/4}} - \dots$

From which we infer

Hence  $\log \frac{1}{2} + \log \frac{3}{2} + \log \frac{5}{2} + \log \frac{7}{2} + \dots = \frac{1}{2} - \frac{1}{2^{n-1}} - \frac{1}{2^{n-1}} - \frac{1}{2^{n-1}} - \frac{1}{2^{n-1}} - \frac{1}{2^{n-1}} + \dots$

$\frac{1}{2} - \frac{1}{2^{n-1}}$  nearly.

$\log 2 + \log 3 + \log 5 + \log 7 + \dots + \log 11 + \dots + \dots$

$= e^{-x} \log 1 - e^{-2x} \log 2 + e^{-3x} \log 3 - e^{-4x} \log 4 + \dots$

whence  $\phi(x) - \phi(2x) + \phi(3x) - \phi(4x) + \dots + \dots$

$= \log 2 (2e^{-2x} + 4e^{-4x} + 8e^{-8x} + 16e^{-16x} + \dots) + \phi(x)$

+  $\dots$

+  $\log 5 (e^{-5x} + e^{-25x} + e^{-125x} + \dots)$

+  $\log 3 (e^{-3x} + e^{-9x} + e^{-27x} + \dots)$

+  $\log 2 (e^{-2x} + e^{-4x} + e^{-8x} + \dots)$

{ being finite now  
2, 3, 5, 7 &c



$$+ \log x^{-2} + \log x^{-2} - 1 \log x^{-2} +$$

$$\left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) \log x =$$

$$\left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) \log x +$$

$$\left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) \log x +$$

$$\left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) \log x +$$

$$\left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) \log x$$

$$\log \left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) +$$

$$\log \left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) +$$

$$\log \left( \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots \right) =$$

$$x \log x + 2 \log x + 3 \log x + \dots$$

$$w^4 = w^3 + 2w^2 + 9w$$

$$f(x) = \frac{f(x) f(x)}{x^2 f(x) f(x)} \text{ and } w = \frac{f(x) f(x)}{x^2 f(x) f(x)}$$

according to the above app: the parameter of  $\chi(a+b)$  is  $3.99944(a+b)$  for  $\chi(a+b)$ .

$$\frac{(a+b)^2}{3x} = 2x - \frac{2 + \sqrt{1-3x}}{3x}$$

If the parameter of an ellipse =  $\pi(a+b)(1+k)$ , the

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) + \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) + \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) - \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) - \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) + \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) + \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) - \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) - \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) + \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) + \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) - \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) - \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) + \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) + \frac{\pi}{2}$$

$$\frac{1}{\sqrt{1-k^2}} \left\{ \frac{\pi}{2} \phi(x) - \frac{\pi}{2} \right\} = \frac{\pi}{2} \phi(x) - \frac{\pi}{2}$$

Then  $pQ + \frac{pQ}{25} = (\frac{p}{Q})^2 + (\frac{Q}{p})^2 - 2(\frac{p}{Q} + \frac{Q}{p} + 2)$

$$f(x) = \frac{f(x)f(x)}{f(x)f(x)}$$

$$Q = \frac{f(x)f(x)}{f(x)f(x)}$$

If  $U_1 + U_2 + U_3 + \dots + U_n$  and  $V_1 + V_2 + V_3 + \dots + V_n$  are nearly equal and also  $\frac{1}{U_n}$  and  $\frac{1}{V_n}$  are nearly equal for all  $n$  then the two series are nearly equal and also divergent but the diff<sup>er</sup> between the two series from

is derivable by G.

Given the pth power of any integer  $\neq 1$

If G be the G.C.M. of any one of  $2^p-1, 2^p, 3^p+1$  and any one of  $3^p-1, 3^p, 3^p+1$

$x^2 + 15y^2$	$80n + 49$	$80n + 1$
$x^2 - 15y^2$	$60n + 49$	$60n + 1$
$5x^2 + 3y^2$	$80n + 17$	$80n - 7$
$5x^2 - 3y^2$	$60n + 17$	$60n - 7$
$x^2 + 6y^2$	$24n + 7$	$24n + 1$
$x^2 - 6y^2$	$24n + 19$	$24n + 1$
$2x^2 + 3y^2$	$24n + 11$	$24n + 5$
$2x^2 - 3y^2$	$24n - 1$	$24n + 5$

$$\left\{ \frac{27^{\frac{1}{3}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) + \frac{\pi^{\frac{1}{2}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) - \frac{1}{(10)^{\frac{1}{2}}} \left\{ \frac{11}{17} \right\} \right\} \frac{11}{17} =$$

$$\frac{27^{\frac{1}{3}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) + \frac{\pi^{\frac{1}{2}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) - \frac{1}{(10)^{\frac{1}{2}}} \left\{ \frac{11}{17} \right\} \frac{11}{17} =$$

$$\frac{27^{\frac{1}{3}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) + \frac{\pi^{\frac{1}{2}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) - \frac{1}{(10)^{\frac{1}{2}}} \left\{ \frac{11}{17} \right\} \frac{11}{17} =$$

$$\frac{27^{\frac{1}{3}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) + \frac{\pi^{\frac{1}{2}}}{(17)^{\frac{1}{2}}} \left( \frac{1}{17} \right) - \frac{1}{(10)^{\frac{1}{2}}} \left\{ \frac{11}{17} \right\} \frac{11}{17} =$$

with the condition  $a/b = 1$

$$= \psi(1) - \frac{1}{2} \psi\left(\frac{2}{3}\right) - \frac{1}{3} \psi\left(\frac{2}{3}\right) - \frac{1}{5} \psi\left(\frac{2}{3}\right) + 8c$$

$$\left\{ \frac{2}{3} \psi\left(\frac{2}{3}\right) - \frac{1}{3} \psi\left(\frac{2}{3}\right) - \frac{1}{5} \psi\left(\frac{2}{3}\right) + 8c \right\} \frac{2}{3} \text{ from}$$

$$\int_0^{\infty} f(x) \cos x \, dx = \psi(x)$$

$$= \sqrt{\frac{1}{\pi}} \left( e^{-\frac{1}{\pi}} - \frac{1}{\pi} e^{-\frac{2}{\pi}} - \frac{1}{\pi^2} e^{-\frac{3}{\pi}} - \frac{1}{\pi^3} e^{-\frac{4}{\pi}} - \dots \right)$$

$$= \frac{1}{\pi} \left( e^{-\frac{1}{\pi}} - \frac{1}{\pi} e^{-\frac{2}{\pi}} - \frac{1}{\pi^2} e^{-\frac{3}{\pi}} - \frac{1}{\pi^3} e^{-\frac{4}{\pi}} - \dots \right)$$

$$\frac{1}{2} + \frac{1}{3} + \frac{1}{5} + \frac{1}{6} + \frac{1}{9} + \frac{1}{10} + \dots - 8c$$

If a prime no of the form  $A_1 + B$  can be  
 expressed as  $a^2 - cy^2$  then a prime no of the form  
 $A_1 - B$  can be expressed as  $bx - ay^2$   
 All nos can be expressed as the sum of 4 squares  
 All nos except of the form  $(8n-1)$  can be expressed as the  
 sum of 3 perfect squares  
 All nos of the form  $2^p \cdot 3^{2q} \cdot 5^r \cdot 7^s \cdot 11^t \cdot 13^u \cdot 17^v$   
 can be expressed as the sum of 2 squares.  
 $p, q, r, s, t, u, v$  may have all integral values including 0.

$x^2 + y^2$	$4n + 1$
$x^2 + 2y^2$	$8n + 1, 8n + 3$
$x^2 + 3y^2$	$6n + 1$
$x^2 + 3y^2$	$12n + 1$
$x^2 + 5y^2$	$n + 1, 20n + 9$
$x^2 + 5y^2$	$5n + 1, 10n + 9$
$x^2 + 7y^2$	$14n + 1, 14n + 9, 14n + 25$
$x^2 + 7y^2$	$28n + 1, 28n + 9, 28n + 25$

A prime no of the form \_\_\_\_\_ can be expressed as \_\_\_\_\_

$$\frac{f(x_1)}{3} + \frac{f(x_2)}{3} + \frac{f(x_3)}{3} = \frac{f(x)}{3}$$

$$\frac{f(x_1)}{3} + \frac{f(x_2)}{3} + \frac{f(x_3)}{3} = \frac{f(x)}{3}$$

$\sqrt{\frac{1}{N} \sum_{i=1}^N x_i^2}$  is the root mean square

$$= \frac{1}{n} \sum_{i=1}^n x_i^2$$

and that of the sum of the divisors of  $n$

The average value of the most divisors of  $x$  is  $2c + \log x$  exactly

then the average value of  $a_n = 11n$  exactly

$$= \int_0^\infty e^{-nx} 11n dx + (a_1 + a_2 + a_3 + \dots) + 2c$$

$$\text{If } a_1 e^{-x} + a_2 e^{-2x} + a_3 e^{-3x} + \dots$$

- I If  $n$  continuous numbers at least  $5/6$  of them can be expressed as the sum of 3 sqs.
- II If  $n$  even numbers at least  $11/12$  of them can be expressed as the sum of 3 sqs.
- III If  $n$  odd numbers at least  $7/8$  of them can be expressed as the sum of 3 sqs.

$$1 = x \cdot \cos x = \frac{\sin \frac{x}{2}}{\frac{x}{2}} = \frac{f(-x) + f(x)}{2} = \frac{1}{2} \log a \log b$$

$$f(m+n) = p+q = r$$

$$\frac{dx}{x} = \frac{p}{\log n} \log a + \frac{q}{\log n} \log b + \frac{r}{\log n} \log c$$

$$\int \frac{dx}{x} = \int \frac{p}{\log n} \log a + \frac{q}{\log n} \log b + \frac{r}{\log n} \log c$$

If the magd. no of each nos = x

$$= \frac{p \log a}{1} + \frac{q \log b}{1} + \frac{r \log c}{1}$$

$$1 + \frac{p}{a} + \frac{q}{b} + \frac{r}{c}$$

sol.

$$\frac{1}{2} \log a \log b$$

$$= (1 + \frac{1}{2})(1 + \frac{1}{4})(1 + \frac{1}{8})$$

$$\sqrt{2(1 - \frac{1}{2})(1 - \frac{1}{4})(1 - \frac{1}{8})}$$

$$f(x) = \frac{\phi(x)}{(x-m)^2}$$

Series  $m+n$

When  $x$  becomes unity  
and the rest are equal  
the least —  $2n+1$

- 24n+1
- 24n+5
- 24n+7
- 24n+11
- 24n+13
- 24n+17
- 24n+19
- 24n+23

the least —  $12n+1$   
The rest are equal

- 12n+1
- 12n+5
- 12n+7
- 12n+11
- 12n+1

the least —  $10n+1$   
less than }  $10n+3$   
                  }  $10n+9$

- 10n+1
- 10n+3
- 10n+7
- 10n+9
- 10n+1

the least —  $8n+1$   
The rest are equal

- 8n+1
- 8n+3
- 8n+5
- 8n+7
- 8n+1

Hence the one of previous nos  
of the form  $4n+1$  is less.  
Similarly for  $6n+1$ .

- 4n+1
- 4n-1
- 6n+1
- 6n-1

If a number  
is a factor of the form



$e^{\frac{1}{2} \log x} = e^{\frac{1}{2} \log x}$   
 $= e^{\frac{1}{2} \log x}$

The area of the region bounded by the lines  $x=0$ ,  $y=0$ ,  $x=1$ , and  $y=1$  is  $\frac{1}{2}$ .  
 $\int_0^1 \int_0^1 dx dy = \frac{1}{2}$

$x^2 = a + y^2 = a + 2$  and  $e^x = a + x$   
 $x^2 + y^2 = a + 2$  and  $e^x = a + x$

$\frac{2x \log x + \log x}{(e + \log x)^2} + \frac{e \log x + \log x}{(e + \log x)^2}$   
 $\frac{2x \log x + \log x}{(e + \log x)^2} + \frac{e \log x + \log x}{(e + \log x)^2}$

$e^x + a^x = e^x + e^{ax} = e^x + e^{ax}$   
 $e^x + a^x = e^x + e^{ax}$

$e^{-x} + e^{-ax} + e^{-bx} = e^{-x} + e^{-ax} + e^{-bx}$   
 $e^{-x} + e^{-ax} + e^{-bx} = e^{-x} + e^{-ax} + e^{-bx}$





$$p = \frac{f(-z)}{f(z)} = \frac{f(-z)}{z^2 f(z)} = \frac{f(-z)}{z^2} \cdot \frac{1}{f(z)}$$

$$p = \frac{f(-z)}{z^2} \cdot \left( \frac{1}{z^2} - \frac{1}{z} + \frac{1}{2} - \frac{1}{6}z + \frac{1}{24}z^2 - \frac{1}{720}z^4 + \dots \right)$$

$$p = \frac{f(-z)}{z^4} \left( 1 - z + \frac{1}{2}z^2 - \frac{1}{6}z^3 + \frac{1}{24}z^4 - \frac{1}{720}z^5 + \dots \right)$$